

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/00</b>	<b>A2</b>	(11) International Publication Number: <b>WO 00/29561</b>
		(43) International Publication Date: 25 May 2000 (25.05.00)

(21) International Application Number: PCT/DK00/00144  
(22) International Filing Date: 27 March 2000 (27.03.00)

(30) Priority Data:  
PA 1999 00427 29 March 1999 (29.03.99) DK  
60/128,558 9 April 1999 (09.04.99) US

(71) Applicant (for all designated States except US): STATENS  
SERUM INSTITUT [DK/DK]; Artillerivej 5, DK-2300  
Copenhagen S (DK).

(72) Inventor; and

(75) Inventor/Applicant (for US only): FOMSGAARD, Anders  
[DK/DK]; Hostrups Have 5, DK-1954 Frederiksberg C  
(DK).

(74) Agent: PLOUGMANN, VINGTOFT & PARTNERS A/S;  
Sankt Annæ Plads 11, P.O. Box 3007, DK-1021 Copen-  
hagen K (DK).

(81) Designated States: AE, AG, AL, AM, AT, AT (Utility model),  
AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU,  
CZ, CZ (Utility model), DE, DE (Utility model), DK, DK  
(Utility model), DM, DZ, EE, EE (Utility model), ES, FI,  
FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID,  
IL, IN, IS, JP, KE, KG, KP, KR, KR (Utility model), KZ,  
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,  
MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,  
SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US,  
UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS,  
MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ,  
BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,  
GN, GW, ML, MR, NE, SN, TD, TG).

**Published**


*Upon the request of the applicant, before the expiration of  
the time limit referred to in Article 21(2)(a).*

*Without international search report and to be republished  
upon receipt of that report.*

*Without classification; title and abstract not checked by the  
International Searching Authority.*

(54) Title: METHOD FOR PRODUCING A NUCLEOTIDE CONSTRUCT WITH OPTIMISED CODONS FOR AN HIV GENETIC  
VACCINE BASED ON A PRIMARY, EARLY HIV ISOLATE AND SYNTHETIC ENVELOPE BX08 CONSTRUCTS

**Synthetic BX08 Env**  
**Strategy for building the full-length gp160 and derived truncated forms**

**(57) Abstract**

The present invention relates to a method for producing a nucleotide sequence construct with optimized codons for an HIV genetic vaccine based on a primary, early HIV isolate. Specific such nucleotide sequence construct are the synthetic envelope BX08 constructs. The invention further relates to the medical use of such constructs for the treatment and prophylaxis of HIV through DNA vaccine and for diagnostics.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Method for producing a nucleotide sequence construct with optimised codons for an HIV genetic vaccine based on a primary, early HIV isolate and synthetic envelope BX08 constructs.

## 5 Field of the invention

The invention relates to a DNA vaccine against HIV, which is designed from a clinical primary isolate. One aspect of the invention relates to a method of producing a nucleotide sequence construct, in a preferred aspect based on a cassette system, the nucleotide sequence construct being used as a DNA vaccine. The method can, for example, lead to the  
10 disclosed synthetic BX08 HIV-1 envelope vaccine nucleotide sequence construct, designed to generate suitable DNA vaccines against HIV, specifically HIV-1. Furthermore, the invention can be used for the production of recombinant protein antigens.

## Background of the invention

15 There is an urgent need for new vaccine strategies against HIV. One such new promising strategy is called genetic immunisation or DNA vaccine (Webster et al 1997). Some of the advantages of a DNA vaccine against HIV is the induction of Th cell activation, induction of antibodies also against conformational dependent epitopes, and the induction of cellular immunity. So far, most DNA vaccine envelope genes tried, have been from tissue culture  
20 adapted virus strains (Boyer et al 1997) that often differs in several aspects from primary clinical isolates (such as early isolates) e.g. in co-receptor usage (Choe et al 1996, Dragic et al 1997).

One disadvantage in HIV envelope based DNA vaccines may be the intrinsic relatively low  
25 expression which is regulated by the Rev expression. This may prevent an optimal investigation of the vaccines in small animal models like mice where Rev is functioning suboptimally. Recently it has been shown using the tissue culture adapted HIV-1 MN strain, that an exchange of the HIV codon usage to that of highly expressed mammalian genes greatly improves the expression in mammalian cell lines and renders the HIV expression Rev  
30 independent (Haas et al 1996). Additionally, it is known that rare codons cause pausing of the ribosome, which leads to a failure in completing the nascent polypeptide chain and a uncoupling of transcription and translation. Pausing of the ribosome is thought to lead to exposure of the 3' end of the mRNA to cellular ribonucleases.

The world-wide spread of HIV-1 has presently resulted in 8,500 new infections daily and AIDS is now the number 1 cause of death among US males (and number 3 among US females) aged 25-40 years. The epidemic hot-spots now include Eastern Europe, India and South East Asia and southern Africa. The attempts to solve this world-wide problem involve education, prevention, treatment and vaccine development. Affordable protective vaccines represent the best solution to the world-wide problem of infection with HIV-1. Induction of virus neutralising antibodies is one of the key components in vaccine development. Several recombinant envelope vaccines have been tested in humans and animals, however, they seem unable to induce sufficient protection. In this respect DNA vaccination may provide a different and more natural mode of antigen presentation. It is hoped that the immune responses induced by such DNA vaccines could aid in limiting virus replication, slowing disease progression or preventing occurrence of disease. Unfortunately many HIV envelope vaccines induce only moderate levels of antibodies. This could in part be due to limitations in expression, influenced by regulation by the Rev protein and by a species-specific and biased HIV codon usage. Also the virus variability is considered a barrier for development of antibody based vaccines and thus a tool for more easy producing of closely related vaccine variants is needed.

It has been suggested to improve the immunogenicity and antigenicity of epitopes by certain mutations in the envelope gene. An elimination of certain immune dominant epitopes (like V3) could render less immune dominant but more relevant, conserved, or hidden epitopes more immunogenic (Bryder et al 1999). Also elimination of certain N-linked glycosylation sites could improve the exposure of relevant epitopes and increase the immunogenicity of those epitopes. Thus, it is possible that elimination of the glycosylation sites in V1 and V2 may in a more favourable way expose neutralising epitopes (Kwong et al 1998, Wyatt et al 1998). The HIV envelope contains putative internalisation sequences in the intracellular part of gp41 (Sauter et al 1996). Thus it would be relevant to eliminate and/or mutate the internalisation signals in a membrane bound HIV envelope vaccine gene to increase the amount of surface exposed vaccine derived HIV glycoproteins as gp150. Since the antibody response, that is measured and calculated in titers, is improved by adding the secreted gp120 as opposed to adding the membrane bound form (Vinner et al 1999), it could be advantageous to express the vaccine as a secreted gp120 or a secreted gp140. This would include important parts of gp41, such as the 2F5 neutralising linear epitope (Mascola et al 1997).

## Summary of the invention

Our suggested solution to the problems described above is to design DNA envelope vaccines from a clinical primary isolate with Rev-independent high expression in mammals, that is built as a cassette for easy variant vaccine production.

5

- A method of producing a nucleotide sequence construct with codons from highly expressed mammalian proteins based on a cassette system coding for an early, primary HIV envelope is described. The method comprises the steps of direct cloning of an HIV gene, derived from a patient within the first 12 months of infection, thereby obtaining a first nucleotide sequence;
- 10 designing a second nucleotide sequence utilising the most frequent codons from mammalian highly expressed proteins to encode the same amino acid sequence as the first nucleotide sequence; redesigning the second nucleotide sequence so that restriction enzyme sites surround the regions of the nucleotide sequence encoding functional regions of the amino acid sequence and so that selected restriction enzyme sites are removed, thereby obtaining
- 15 a third nucleotide sequence encoding the same amino acid sequence as the first and the second nucleotide sequence; redesigning the third nucleotide sequence so that the terminals contain convenient restriction enzyme sites for cloning into an expression vehicle; producing snuts between restriction enzyme sites as well as terminal snuts and introducing snuts into an expression vehicle by ligation. The nucleotide sequence construct obtained by this
- 20 method uses the mammalian highly expressed codons (figure 1) and renders the envelope gene expression Rev independent and allows easy cassette exchange of regions surrounded by restriction enzyme sites that are important for immunogenicity, function, and expression.
- 25 The method can, for example, lead to the disclosed synthetic, Rev-independent, clinical (such as early), primary HIV-1 envelope vaccine gene, built as a multi cassette. From the sequence of the envelope of the HIV-1 BX08 isolate (personal communication from Marc Girard, Institute Pasteur, Paris), the present inventors have designed a synthetic BX08 HIV-1 envelope vaccine nucleotide sequence construct.

30

With the great diversity of envelopes in HIV among different patients and within one patient, it would be of advantage to vaccinate with several envelope variants, all being highly expressed. To avoid synthesising several full length envelopes, it is much easier to exchange relevant parts of an envelope cassette to various strains in a multivalent vaccine.

35

Whether it is the disclosed synthetic BX08 nucleotide sequence construct, or any of the nucleotide sequence constructs obtained by the method, they are designed to generate suitable DNA vaccines against HIV, specifically HIV-1. In this case the mammal, preferably a human being, is inoculated with the nucleotide sequence construct in an expression vehicle  
5 and constitutes a host for the transcription and translation of the nucleotide sequence construct. The nucleotide sequence constructs of the present invention can furthermore be used for the production of recombinant protein antigens. In this case the nucleotide sequence construct is placed in an expression vehicle and introduced into a system (e.g. a cell-line), allowing production of a recombinant protein with the same amino acid sequence.  
10 The recombinant protein is then isolated and administered to the mammal, preferably a human being. The immune system of the mammal will then direct antibodies against epitopes on the recombinant protein. The mammal, preferably a human being, can thus be primed or boosted with DNA and/or recombinant protein obtained by the method of the invention.

15 A relevant HIV DNA vaccine can potentially be used not only as a prophylactic vaccine, but also as a therapeutic vaccine in HIV infected patients, e.g. during antiviral therapy. An HIV specific DNA vaccine will have the possibility to induce or re-induce the wanted specific immunity and help the antiviral therapy in limiting or even eliminating the HIV infection. The  
20 immunogenicity and antigenicity of epitopes in the envelope can be improved by certain mutations in the envelope gene. The cassette system allows for easy access to the relevant parts of the envelope gene, and thereby eased efforts in the process of genetic manipulation. Some suggested mutations are: an elimination of certain immune dominant epitopes (like V3); elimination of certain N-linked glycosylation sites (like glycosylation sites around V2);  
25 elimination and/or mutation of the nucleotide sequence encoding the internalisation signals in the cytoplasmic part of a membrane bound HIV envelope to increase the amount of surface exposed vaccine derived HIV glycoproteins; elimination or mutation of the cleavage site between gp120 and gp41; with introduced mutations in gp41 for preserving conformational epitopes.

30 Table 1 below, lists the nucleotide sequence constructs of the invention by the names used herein, as well as by reference to relevant SEQ ID NOs of DNA sequences, and the amino acid sequence encoded by the DNA sequence in the preferred reading frame. It should be noted, that the snut name consist of the number of the approximate position for the end of  
35 the snut and the restriction enzyme used to cleave and/or ligate that end of the snut.

**Table 1 List of names of nucleotide sequence constructs (Snuts (S) and Pieces (P)) with reference to SEQ ID NO for the nucleotide sequence and protein sequence.**

Name	Nucleotide SEQ ID NO:	Protein SEQ ID NO:
S <sub>0-N-Lang</sub>	1	2
S <sub>235EcoRV</sub>	3	4
S <sub>375PstI</sub>	5	6
S <sub>495ClaI</sub>	7	8
S <sub>650-720EcoRI</sub>	9	10
S <sub>900XbaI</sub>	11	12
S <sub>990SacI</sub>	13	14
S <sub>1110SpeI</sub>	15	16
S <sub>1285XhoI</sub>	17	18
S <sub>1285gp120</sub>	19	20
S <sub>1285gp160</sub>	21	22
S <sub>1465PstI</sub>	23	24
S <sub>1465PstI cys</sub>	25	26
S <sub>1630XbaI</sub>	27	28
S <sub>1700EagI</sub>	29	30
S <sub>1890HindIII</sub>	31	32
S <sub>2080SacII</sub>	33	34
S <sub>2190ClaI</sub>	35	36
S <sub>2330PstI</sub>	37	38
S <sub>2425ES</sub>	39	40
P <sub>1</sub>	41	42
P <sub>2</sub>	43	44
P <sub>3</sub>	45	46
P <sub>3GV1</sub>	47	48
P <sub>3GV1V2</sub>	49	50
P <sub>3GV2</sub>	51	52
P <sub>4gp160</sub>	53	54
P <sub>4gp150</sub>	55	56
P <sub>4gp140</sub>	57	58
P <sub>5</sub>	59	60
P <sub>8gp160</sub>	61	62
P <sub>8gp150</sub>	63	64
P <sub>8gp140</sub>	65	66
synBX08-140	67	68
synBX08-150	69	70
synBX08-160	71	72
synBX08-120	73	74
synBX08-41	75	76

## Detailed disclosure of the invention

One aspect of the present invention relates to a method for producing a nucleotide sequence construct coding for an HIV gene. The nucleotide sequence construct is produced as a cassette system consisting of snuts. A snut (S) is a nucleotide sequences construct between  
5 restriction enzyme cleavage sites comprising the minimal entity of the cassette system.

First an HIV gene is obtained from a patient within the first 12 months of infection. The term HIV should be understood in the broadest sense and include HIV 1 and HIV 2. It is possible to determine the period in which the infection has taken place with an accuracy depending  
10 on the frequency of the blood tests taken from the patient. For example, patients suffering from various diseases such as lack of certain factors in their blood or hepatitis have their blood tested on a regular basis making it possible to determine the period in which the infection has taken place. Apart from patients with diseases wherein blood tests are used to monitor the course of the disease, other groups of patients have blood tests taken, e.g. blood  
15 donors. Unfortunately, humans are still infected due to transfer of virus in blood samples, medical equipment, etc., making it possible to determine the date where the infection has taken place within the time frame of a few days. The importance of obtaining the virus early in the course of the infection is due to the known fact that many early isolates share the common feature of staying relatively constant in their envelope sequences (Karlsson et al.,  
20 1998). As these early isolates may share cross-reactive antibody- and/or T-cell epitopes a vaccine based on such early isolates would have a better chance of inducing immune response to shared epitopes of the virus. It is believed that an early, directly cloned virus isolate will share immunogenic sites with other early virus isolates seen during an HIV infection, so that if a mammal generates antibodies and/or T-cells directed against these  
25 epitopes, the transferred virus will be eliminated prior to the extensive mutations that may occur after approximately 12 months of infection. Thus, the virus should be isolated as early as possible, that is within the first 12 months of infection, such as 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, or 0.5 month after infection.

30 The HIV gene for genetic vaccine is preferably cloned directly from viral RNA or from proviral DNA. Direct cloning in this application stands for the virus not being multiplied in stable cell lines *in vitro*. It is presently expected that passing the virus through a stable cell line will promote mutation in the virus gene. It is particularly preferred not to pass the virus through cells lines selecting for viruses with CXCR4 receptor usage. Direct cloning also includes  
35 multiplication of virus in e.g. PBMC (peripheral blood mononuclear cells) since all virus can



multiply in PBMC, and this type of multiplication generally does not select for CXCR4 receptor usage. Multiplication of virus is often necessary prior to cloning. Preferably cloning is performed directly on samples from the patient. In one embodiment of the invention, cloning is performed from patient serum. The cloning is then performed directly on the HIV virus, that is on RNA. In another embodiment of the invention cloning, is performed from infected cells. The cloning is then performed on HIV virus incorporated in the genes in an infected cell (e.g. a lymphocyte), that is on DNA. In the latter case the virus might be a silent virus, that is a non-replicating virus. To evaluate if the virus is silent, capability of multiplication in e.g. PBMC is tested.

10

Cloning is a technique well known to a person skilled in the art. A first nucleotide sequence is hereby obtained. In another aspect of the invention, the first nucleotide sequence, sharing the properties mentioned with direct cloning, is obtained by other means. This could be from a database of primary isolates or the like.

15

Based on the first nucleotide sequence, the amino acid sequence encoded by said nucleotide sequence is determined. A second nucleotide sequence encoding the same amino acid sequence is then designed utilising the most frequent codons from highly expressed proteins in mammals (e.g. figure 1 presenting the most frequent codons from

20 highly expressed proteins in humans).

Presently, it appears that the usage of the most frequent codons from mammalian highly expressed proteins has two advantages: 1) the expression is Rev independent; 2) the level of expression is high. The Rev independence is especially advantageous when performing experiments in mice where the Rev systems is functioning sub-optimally. For the use in human vaccine, Rev independence and high expression are important to increase the amount of antigen produced. The determination of the codons for high expression is in this context based on the statistics from human highly expressed proteins (Haas, Park and Seed, 1996 hereby incorporated by reference). It is contemplated that the expression of a protein can be even higher, when current research in binding between codon (on the mRNA) and anticodon (on the tRNA) reveals codons with optimal binding capabilities, and when interactions in-between codons and/or in-between anticodons are known.

The second nucleotide sequence designed utilising optimised codons is then redesigned to obtain a third nucleotide sequence. The purpose of the redesigning is to create unique restriction enzyme sites around the nucleotide sequence encoding functional regions of the

- amino acid sequence. By having unique restriction enzyme sites around the nucleotide sequence encoding functional regions of the amino acid sequence, the nucleotide sequence encoding functional regions of the amino acid sequence can easily be isolated, changed, and re-inserted. Examples of functional regions of the amino acid sequence are transmembrane
- 5 spanning regions, immunodominant regions, regions with antibody cross reacting domains, fusion domains and other regions important for immunogenicity and expression such as variable region 1 (V1), variable region 2 (V2), variable region 3 (V3), variable region 4 (V4) and variable region 5 (V5).
- 10 It is important to select the restriction enzymes sites with care. By changing the second nucleotide sequence to insert restriction enzyme sites around the nucleotide sequence encoding functional regions of the amino acid sequence, the third nucleotide sequence must still code for the same amino acid sequence as the second and first nucleotide sequence do. Thus, if necessary, the second nucleotide sequence is redesigned by changing from
- 15 optimised codons to less optimal codons. It is understood, that the restriction enzyme sites around the nucleotide sequence encoding functional regions of the amino acid sequence should preferably be placed in the terminal region of the nucleotide sequence encoding functional regions of the amino acid sequence. That is preferably outside the nucleotide sequence encoding functional regions of the amino acid sequence, such as 90 nucleotides
- 20 away, e.g. 81, 72, 63, 54, 45, 36, 27, 21, 18, 15, 12, 9, 6, 3 nucleotides away, but could also be inside the nucleotide sequence encoding functional regions of the amino acid sequence, such as 54, 45, 36, 27, 21, 18, 15, 12, 9, 6, 3 nucleotides inside the nucleotide sequence encoding the functional region of the amino acid sequence.
- 25 The type of restriction enzyme sites allowed is determined by the choice of expression vector. In certain cases, the number of restriction enzyme sites is limited and it is hard, if not impossible, to place unique restriction enzyme sites around all the nucleotide sequences coding for functional regions of the amino acid sequence. This problem can be solved by dividing the entire nucleotide sequence into pieces, so that each piece comprises only
- 30 unique restriction enzyme sites. Modifications to each of the piece is performed separately prior to assembly of the pieces. It is preferred that the nucleotide sequence is divided into 9 pieces. In another aspect, the nucleotide sequence is divided into 8 pieces, or 7, or 6, or 5, or 4, or 3, or 2 pieces. It is especially preferred that the nucleotide sequence is divided into 3 pieces.

Thus, the redesign of the second nucleotide sequence is an interaction between the choice of cloning vector, expression vector, selection of restriction enzyme sites, dividing into pieces, and exchange of codons to insert restriction enzyme sites. In a preferred embodiment of the present invention the cloning vector is Bluescript allowing the restriction enzyme sites chosen from the group consisting of: *EagI*, *MluI*, *EcoRV*, *PstI*, *Clal*, *EcoRI*, *XbaI*, *SacI*, *SpeI*, *XhoI*, *HindIII*, *SacII*, *NotI*, *BamHI*, *SmaI*, *Sall*, *DraI*, *KpnI*. If other cloning vectors are chosen, other restriction enzyme sites will be available as known by the person skilled in the art.

- 10 As a part of the redesigning of the second nucleotide sequence, selected restriction enzyme sites may be removed. The selected restriction enzyme sites to be removed are those sites that are sites of the same type as the ones already chosen above and that are placed within the same piece. The removal of these restriction enzyme sites is performed by changing from optimised codons to less optimal codons, maintaining codons for the same amino acid sequence.

- The third nucleotide sequence is redesigned so that the terminal snuts contain convenient restriction enzyme sites for cloning into an expression vehicle. The expression "vehicle" means any nucleotide molecule e.g. a DNA molecule, derived e.g. from a plasmid, bacteriophage, or mammalian or insect virus, into which fragments of nucleic acid may be inserted or cloned. An expression vehicle will contain one or more unique restriction enzyme sites and may be capable of autonomous replication in a defined host or vehicle organism such that the cloned sequence is produced. The expression vehicle is an autonomous element capable of directing the synthesis of a protein. Examples of expression vehicles are mammalian plasmids and viruses, tag containing vectors and viral vectors such as adenovirus, vaccinia ankara, adenoassociated virus, canarypox virus, simliki forest virus (sfv), Modified Vaccinia Virus Ankara (MVA), and simbis virus. In one embodiment of the invention, the expression vector contains tag sequences. In another embodiment of the invention a bacteria is transformed with an expression plasmid vector and the bacteria is then delivered to the patient. Preferred expression vehicles are simliki forest virus (sfv), adenovirus and Modified Vaccinia Virus Ankara (MVA).

- The snuts are produced by techniques well known by the person skilled in the art. The preferred method for synthesising snuts, is herein referred to as "the minigene approach" wherein complementary nucleotide strands are synthesised with specific overhanging sequences for annealing and subsequent ligation into a vector. This can be performed with

two sets of complementary nucleotide strands, or with three sets of complementary nucleotide strands. The minigene approach minimises the known PCR errors of mismatches and/or deletions, which may occur due to hairpins in a GC rich gene with mammalian highly expressed codons. In figures 10-21, the production of a representative selection of snuts is  
5 illustrated.

For the production of long snuts, that is snuts of more than about 240 nucleotides, the technique of overlapping PCR is preferred as illustrated in figure 8. Herein two nucleotide strands about 130 nucleotides long with an overlap are filled to obtain a double strand, which  
10 is subsequently amplified by PCR.

For the production of multiple snuts with a length of less than about 210 nucleotides, one preferred technique is normal PCR. In a preferred production technique the snuts are synthesised with the same 5' flanking sequences and with the same 3' flanking sequences,  
15 as illustrated in figure 9. The advantages of this approach is, that the same PCR primer set can be used for amplification of several different snuts.

As known by the person skilled in the art, special conditions have to be used for each individual PCR reaction and it should be optimised to avoid inherent problems like deletions  
20 mismatches when amplifying GC rich genes from synthetic ssDNA material. Whichever of the above mentioned techniques are used, it is well known by the person skilled in the art, that it will be necessary to correct unavoidable mismatches produced either due to the nucleotide strand synthesis material and/or the PCR reaction. This can be performed by site directed mutagenesis techniques.

25

After the various snuts have been produced, they are assembled into pieces and subsequently into the complete gene. Methods for assembly (such as ligation) are well known by the person skilled in the art.

30

In a preferred embodiment of the present invention the HIV gene encodes the entire HIV envelope. It is understood that the HIV envelope can be the full length envelope gp160 as well as shorter versions such as gp150, gp140, and gp120 with or without parts of gp41.

35

As will be known by the person skilled in the art, the HIV is divided into several groups. These groups presently include group M, group O, and group N. Further, the HIV is divided

into subtypes A, B, C, D, E, F, G, H, I, and J. In the present invention subtype B is preferred due to the high prevalence of this subtype in the Western countries.

The determination of groups and subtypes is based on the degree of nucleotide sequence identity in the envelope gene is presently defined as follows: If the sequence identity is more than 90% the viruses belong to the same subtype; If the sequence identity is between 80% and 90% the viruses belong to the same group. If the sequence identity is less than 80% the viruses are considered as belonging to different groups.

10 One aspect of the invention relates to a nucleotide sequence construct in isolated form which has a nucleotide sequence with the general formula (I), (II), (III), or (IV)

(I)  $P_1-S_{495\text{ClaI}}-S_{650-720\text{EcoRI}}-P_2-S_{1265\text{gp120}}$

(II)  $P_1-S_{495\text{ClaI}}-S_{650-720\text{EcoRI}}-P_2-S_{1265\text{XhoI}}-S_{1465\text{PstI}}-P_{4\text{gp140}}$

(III)  $P_1-S_{495\text{ClaI}}-S_{650-720\text{EcoRI}}-P_2-S_{1265\text{XhoI}}-S_{1465\text{PstI}}-P_{4\text{gp150}}$

15 (IV)  $P_1-S_{495\text{ClaI}}-S_{650-720\text{EcoRI}}-P_2-S_{1265\text{XhoI}}-S_{1465\text{PstI}}-P_{4\text{gp160}}-S_{2060\text{SacII}}-P_5$

wherein  $P_1$  designates the nucleotide sequence SEQ ID NO:41, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 90% thereto;

wherein  $S_{495\text{ClaI}}$  designates the nucleotide sequence SEQ ID NO: 7, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 95% thereto;

wherein  $S_{650-720\text{EcoRI}}$  designates the nucleotide sequence SEQ ID NO: 9, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 95% thereto;

25 wherein  $P_2$  designates the nucleotide sequence SEQ ID NO: 43, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 85% thereto;

wherein  $S_{1265\text{gp120}}$  designates the nucleotide sequence SEQ ID NO: 19, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 70% thereto;

wherein  $S_{1265\text{XhoI}}$  designates the nucleotide sequence SEQ ID NO: 17, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 80% thereto;

wherein  $S_{1465\text{PstI}}$  designates the nucleotide sequence SEQ ID NO: 23, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 90% thereto;

wherein P<sub>4gp140</sub> designates the nucleotide sequence SEQ ID NO: 57, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 85% thereto;

wherein P<sub>4gp150</sub> designates the nucleotide sequence SEQ ID NO: 55, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 85% thereto;

wherein P<sub>4gp160</sub> designates the nucleotide sequence SEQ ID NO: 53, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 85% thereto;

10 wherein S<sub>2080SacII</sub> designates the nucleotide sequence SEQ ID NO: 33, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 98% thereto; and

wherein P<sub>5</sub> designates the nucleotide sequence SEQ ID NO: 59, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 85% thereto.

The design of the parent synthetic BX08 gp160 envelope cassette gene with its variant length genes gp150, gp140, gp120 is outlined in figure 2.

20 The nucleotide sequence construct with the formula (I)

(I) P<sub>1</sub>-S<sub>495</sub>Clal-S<sub>650-720</sub>EcoRI-P<sub>2</sub>-S<sub>1265</sub>gp120

(visualised in figure 3) (SEQ ID NO: 73) codes for the amino acid sequence of gp120 (SEQ ID NO: 74). This amino acid sequence is the part of the HIV envelope that is secreted. Thus, it contains the immunogenic epitopes without being bound to the cell membrane. This is of particular advantage if the nucleotide sequence construct is used for production of recombinant antigens or for a DNA vaccine as the antibody immune response may be higher to secreted versus membrane bound HIV antigens.

The nucleotide sequence construct with the formula (II)

30 (II) P<sub>1</sub>-S<sub>495</sub>Clal-S<sub>650-720</sub>EcoRI-P<sub>2</sub>-S<sub>1265</sub>XhoI-S<sub>1465</sub>PstI-P<sub>4gp140</sub>

(visualised in figure 4) (SEQ ID NO: 67) codes for the amino acid sequence of gp140 (SEQ ID NO: 68). This amino acid sequence encodes the gp120 and the extracellular part of the gp41 protein. The amino acid sequence is secreted due to the lack of the transmembrane spanning region. This is of particular advantage if the nucleotide sequence construct is used for production of recombinant antigens as the immunogenic and/or antigenic epitopes in the extracellular part of gp41 are included and is of particular advantage for a DNA vaccine as

the antibody immune response may be higher to secreted gp120 versus membrane bound HIV antigens.

The nucleotide sequence construct with the formula (III)

5 (III)  $P_1-S_{495Clal}-S_{850-720EcoRI}-P_2-S_{1285XhoI}-S_{1485PstI}-P_{4gp150}$

(visualised in figure 5) (SEQ ID NO: 69) codes for the amino acid sequence of gp150 (SEQ ID NO: 70). This amino acid sequence contains all of the envelope protein gp160 except the c-terminal tyrosin containing internalisation signals in the intracellular part of gp41. The membrane bound surface expression of the amino acid sequence is thereby maintained and enhanced. Mimicking the organisation of the native epitope conformation may be expected, making this nucleotide sequence construct of particular advantage if the nucleotide sequence construct is used as a vaccine.

The nucleotide sequence construct with the formula (IV)

15 (IV)  $P_1-S_{495Clal}-S_{850-720EcoRI}-P_2-S_{1285XhoI}-S_{1485PstI}-P_{4gp160}-S_{2060SacII}-P_5$

(visualised in figure 6) (SEQ ID NO: 71) codes for the amino acid sequence of gp160 (SEQ ID NO: 72) i.e. the entire envelope.

The nucleotide sequence construct designated  $P_1$  comprises the nucleotide sequence encoding the amino acid sequence in the first variable region (V1) and the amino acid sequence in the second variable region (V2). In one embodiment of the invention the first variable region is surrounded by EcoRV and PstI restriction enzyme sites, and the second variable region is surrounded by PstI and ClaI restriction enzyme sites but as stated above, the choice of restriction enzyme sites can alter.

25

The nucleotide sequence construct designated  $S_{850-720EcoRI}$  comprises the nucleotide sequence encoding the amino acid sequence in the third variable region (V3). In one embodiment of the present invention  $S_{850-720EcoRI}$  is characterised by the restriction enzyme sites EcoRI and XbaI in the terminals.

30

The nucleotide sequence construct designated  $P_2$  comprises the nucleotide sequence encoding the amino acid sequence of the fourth variable and constant region (V4 and C4). In one embodiment of the present invention the fourth variable region is surrounded by SacI and XhoI restriction enzyme sites.

35

The nucleotide sequence construct designated S<sub>1285gp120</sub> comprises the nucleotide sequence encoding amino acid sequence of the fifth variable and constant region (V5 and C5).

S<sub>1285gp120</sub> further comprises a nucleotide sequence encoding a C-terminal stop codon.

- 5 The nucleotide sequence construct designated P<sub>4gp140</sub> comprises the nucleotide sequence encoding amino acid sequence of the transmembrane spanning region. P<sub>4gp140</sub> further comprises a nucleotide sequence encoding a C-terminal stop codon prior to the transmembrane spanning region.
- 10 The nucleotide sequence construct designated P<sub>4gp160</sub> comprises the nucleotide sequence encoding amino acid sequence of the transmembrane spanning region (trans membrane spanning domain: TMD). In a preferred embodiment of the present invention the transmembrane spanning region is surrounded by HindIII and SacII restriction enzyme sites.
- 15 The term "sequence identity" indicates the degree of identity between two amino acid sequences or between two nucleotide sequences calculated by the Wilbur-Lipman alignment method (Wilbur et al, 1983).

The nucleotide sequence constructs with the formula (I), (II), (III), or (IV) illustrates the

- 20 flexibility in the present invention. By producing a gene with the described method enables the production of a plethora of antigens with various immunogenic epitopes and various advantages for production and vaccine purposes. To further illustrate the flexibility of the invention, other changes and mutations are suggested below.
- 25 In order to improve the immunogenicity of the nucleotide sequence constructs of the invention it is suggested to change the nucleotide sequence such that one or more glycosylation sites are removed in the amino acid sequence. By removal of shielding glycosylations, epitopes are revealed to the immunesystem of the mammal rendering the construct more immunogenic. The increased immunogenicity can be determined by an
- 30 improved virus neutralisation. Changes in the nucleotide sequence such that one or more N-linked glycosylation sites are removed in the amino acid sequence is well known by the person skilled in the art. Potential glycosylation sites are N in the amino acid sequences N-X-T or N-X-S (wherein X is any amino acid besides P). The glycosylation site can be removed by changing N to any amino acid, changing X to a P, or changing T to any amino acid. It is
- 35 preferred that N is changed to Q by an A to C mutation at the first nucleotide in the codon, and a C to G mutation at the third nucleotide in the codon. This is preferred to increase the



GC content in the nucleotide sequence construct. As an alternative N is changed to Q by an A to C mutation at the first nucleotide in the codon, and a C to A mutation at the third nucleotide in the codon. Preferred mutations in the synthetic BX08 envelope gene to remove potential N-linked glycosylation sites in V1 and/or V2 are A307C + C309A and/or A325C + C327G and/or A340C + C342A and/or A385C + C387A and/or A469C + C471A. Examples of such changes is illustrated in SEQ ID NOs: 47, 49, and 51.

For historical reasons the HIVs have been divided into syncytia inducing strains and non syncytia inducing strains. The assay to determine whether a strain is syncytia inducing is described in Verrier et al 1997, hereby incorporated by reference. It is presently known, that viruses utilising the CXCR4 co-receptor are syncytia inducing strains. It is also, at the present, known that the binding site for the CXCR4 involves the third variable region (V3). In a preferred embodiment the nucleotide sequence construct is changed to create a binding site for the CXCR4 co-receptor. It is presently performed in the third variable regions, preferably by the mutation G865C + A866G.

It is well established that the HIV envelope comprises immunodominant epitopes. An immunodominant epitope is an epitope that most antibodies from the mammal are directed against. The antibodies directed against these immunodominant epitopes may have little effect in elimination of the virus. It is therefore anticipated that modification of the immunodominant epitopes will induce antibodies directed against other parts of the envelope leading to a better elimination and neutralisation of the virus. By modification is understood any change in the nucleotide sequence encoding an immunodominant epitope in the amino acid sequence such that said amino acid sequence no longer contains an immunodominant epitope. Thus, modification includes removal of the immunodominant epitope and decrease of immunogenicity performed by mutagenesis. In a preferred embodiment of the present invention an immunodominant epitope in the third variable region (V3) is modified, such as deleted or altered. In a much preferred embodiment the nucleotides 793-897 are deleted. In yet another preferred embodiment of the present invention an immunodominant epitope has been removed from gp41, such as deleted. This is performed in P<sub>7</sub> or P<sub>8</sub> by elimination of the nucleotides 1654-1710.

It is anticipated that when gp120 is dissociated from gp41 in a vaccine or antigen, two immunodominant epitopes, one on each protein, are exposed and antibodies are directed against these in the mammal. In the infectious virus, gp120 is coiled on top of gp41 and the gp120/gp41 is most likely organised in a trimer, so that these immunodominant epitopes are

hidden and therefore less elimination of virus is observed. By removing the cleavage site between gp41 and gp120 a full length gp160, gp150, or gp140 can be obtained with a covalent binding between gp41 and gp120. Removal of the cleavage site between gp41 and gp120 is preferably performed by a mutation at position C1423A. An example of such a  
 5 mutation is illustrated in the mutation of S<sub>1265XhoI</sub> (SEQ ID NO: 17) to S<sub>1265gp160</sub> (SEQ ID NO: 21).

In order to stabilise the full length gp160, gp150, and gp140 for example when the cleavage site between gp41 and gp120 has been removed as described above, cysteins can be  
 10 inserted, preferably inside the gp41 helix creating disulphide bounds to stabilise a trimer of gp41s. In a preferred embodiment of the present invention the cysteins are inserted by the mutation 1618:CTCCAGGC:1625 to 1618:TGCTGCGG:1625. An example of such a change is illustrated in SEQ ID NO: 25.

15 The above mentioned decrease in immunodominant epitopes combined with the increase in immunogenicity of the other epitopes is expected to greatly enhance the efficacy of the nucleotide sequence construct as a vaccine.

During the production of the nucleotide sequence construct, it is convenient to ligate the  
 20 snuts into pieces. The pieces, as described above, are characterised by their reversible assembly as there are no duplicate restriction enzyme sites. In a preferred embodiment one piece (herein designated P<sub>3</sub>) contains P<sub>1</sub>, S<sub>495</sub>Clal, S<sub>650-720</sub>EcoRI, and P<sub>2</sub>. Another piece (herein designated P<sub>6</sub>) contains S<sub>1265</sub>XhoI, S<sub>1465</sub>PstII, and P<sub>4gp160</sub>. Yet another piece (herein designated P<sub>7</sub>) contains S<sub>1265</sub>XhoI, S<sub>1465</sub>PstII, P<sub>4gp160</sub>, S<sub>2060</sub>SacII, and P<sub>5</sub>.

25 One advantage of the present nucleotide sequence construct is the easy access to exchange and alterations in the content and function of the nucleotide sequence and the encoded amino acid sequence. In one embodiment the nucleotide sequence coding for a functional region or parts thereof of the amino acid sequence is repeated. The repeat could  
 30 be back-to-back or a functional region or parts thereof could be repeated somewhere else in the sequence. Repeated could mean two (one repetition) but could also be three, six, or nine repeats. In a much preferred embodiment the repetition nucleotide sequence codes for amino acids in the third variable region.

35 In order to improve the protective capabilities of the invention against infections with HIV, one embodiment of the invention relates to the combination of epitopes. The present

nucleotide sequence construct allows insertion of one or more new nucleotide sequences isolated from another group and/or subtype of HIV and/or isolated from another patient. Hereby a vaccine or antigen with two or more epitopes from two or more HIVs is obtained. In a preferred embodiment, the V3 is replaced by the new nucleotide sequence. In a much  
5 preferred embodiment, the new nucleotide sequence codes for amino acids in the third variable region of a different HIV isolate.

In order to improve the efficacy of the vaccine, aiming at raising cellular immunity, a nucleotide sequence coding for a T-helper cell epitope is included in the nucleotide  
10 sequence construct. The nucleotide sequence coding for a T-helper cell epitope or a T-helper cell epitope containing amino acid sequence can be put in anywhere in the nucleotide sequence construct as long as it does not interact with the function of the envelope molecule. However, it is preferably placed in the tail of the nucleotide sequence construct or between the leader sequence and the envelope gene. The T-helper epitopes are preferably selected  
15 from core proteins such as P24gag or from a non-HIV pathogen such as virus, bacteria, e.g. BCG antigen 85. For a therapeutic vaccine an HIV helper epitope is preferred since the patient is already primed by the HIV infection. For a prophylactic vaccine, a T-helper cell epitope from a frequently occurring non HIV pathogen such as Hepatitis B, BCG, CMV, EBV is preferred. Also, since the synthetic BX08 envelope genes may contain T-helper cell  
20 epitopes in addition to important antibody epitopes, the synthetic BX08 vaccine genes can be mixed with other DNA vaccines to improve the efficacy of the other DNA vaccine.

One aspect of the present invention relates to individualised immunotherapy, wherein the virus from a newly diagnosed patient is directly cloned, the envelope or subunits  
25 corresponding to snuts or pieces is produced with highly expressed codons, inserted into any of the nucleotide sequence constructs described above and administered to the patient as a vaccine. Hereby a therapeutic DNA vaccine is obtained, that will help the patient to break immunetolerance or induce/reinduce an appropriate immune response. In one embodiment the variable regions of the virus are produced with highly expressed codons and exchanged  
30 into any of the nucleotide sequence constructs described above.

In one embodiment of the invention, the nucleotide sequence construct as described above satisfies at least one of the following criteria:  
a) serum extracted from a Macaque primate which has been immunised by administration of  
35 an expression vector containing the nucleotide sequence construct is capable of eliminating SHIV as determined by quantitative PCR and/or virus culturing.

b) serum extracted from a primate which has been immunised by administration of an expression vector containing the nucleotide sequence construct is capable of neutralising HIV-1 BX08 and /or other HIV-1 strains *in vitro*.

- 5 c) serum, extracted from a mouse which has been immunised by administration of an expression vector containing the nucleotide sequence construct four times in intervals of three weeks and boosted after 15 weeks, is capable of decreasing the concentration of HIV-antigen in a culture of HIV, serum or PBMCs by at least 50%. An example of such procedure is shown in example 9.

- 10 In one embodiment of the invention, the nucleotide sequence construct of the invention, is used in medicine. That is, it is used as a vaccine, for the production of a recombinant protein, such that the recombinant protein is used as a vaccine, or the nucleotide sequence construct or the recombinant protein is used in a diagnostic composition.

- Thus, the nucleotide sequence construct of the invention can be used for the manufacture of  
15 a vaccine for the prophylactics of infection with HIV in humans.

- Intramuscular inoculation of nucleotide constructs, i.e. DNA plasmids encoding proteins have been shown to result in the generation of the encoded protein *in situ* in muscle cells and dendritic cells. By using cDNA plasmids encoding viral proteins, both antibody and CTL  
20 responses were generated, providing homologous and heterologous protection against subsequent challenge with either the homologous or cross-strain reaction, respectively. The standard techniques of molecular biology for preparing and purifying DNA constructs enable the preparation of the DNA therapeutics of this invention. While standard techniques of molecular biology are therefore sufficient for the production of the products of this  
25 invention, the specific constructs disclosed herein provide novel therapeutics which can produce cross-strain protection, a result heretofore unattainable with standard inactivated whole virus or subunit protein vaccines.

- The amount of expressible DNA to be introduced to a vaccine recipient will depend on the strength of the transcription and translation promoters used in the DNA construct, and on the  
30 immunogenicity of the expressed gene product. In general, an immunologically or prophylactically effective dose of about 10 µg to 300 µg is administered directly into muscle tissue. Subcutaneous injection, intradermal introduction, impression through the skin, inoculation by gene gun preferably DNA coated gold particles, and other modes of administration such as intraperitoneal, intravenous, peroral, topic, vaginal, rectal, intranasal  
35 or by inhalation delivery are also contemplated. It is also contemplated that booster

vaccinations are to be provided. It is further contemplated that booster vaccinations with recombinant antigens are to be provided, administered as described above.

The DNA may be naked, that is, unassociated with any proteins, adjuvants or other agents which impact on the recipients immune system. In this case, it is desirable for the DNA to be in a physiologically acceptable solution, such as, but not limited to, sterile saline or sterile buffered saline. Alternatively, the DNA may be associated with surfactants, liposomes, such as lecithin liposomes or other liposomes, such as ISCOMs, known in the art, as a DNA-liposome mixture, (see for example WO93/24640) or the DNA may be associated with and adjuvant known in the art to boost immune responses, such as a protein or other carrier. Agents which assist in the cellular uptake of DNA, such as, but not limited to, calcium ions, detergents, viral proteins and other transfection facilitating agents may also be used to advantage. These agents are generally referred to as transfection facilitating agents and as pharmaceutically acceptable carriers.

15

Those skilled in the field of molecular biology will understand that any of a wide variety of expression systems may be used. A wide range of suitable mammalian cells are available from a wide range of sources (e.g. the American Type Culture Collection, Rockland, Dm; also, see e.g. Ausubel et al. 1992). The method of transformation or transfection and the choice of expression vehicle will depend on the host system selected. Transformation and transfection methods are described e.g. in Ausubel et al 1992; expression vehicles may be chosen from those provided e.g. in P.H. Pouwels et al. 1985.

20

In one embodiment of the present invention the protein encoded by the nucleotide sequence construct is produced by introduction into a suitable mammalian cell to create a stably-transfected mammalian cell line capable of producing the recombinant protein. A number of vectors suitable for stable transfection of mammalian cells are available to the public e.g. in *Cloning Vectors: A Laboratory manual* (P.H. Pouwels et al. 1985); methods for constructing such cell lines are also publicly available, e.g. in Ausubel et al. 1992.

30

Standard reference works describing the general principles of recombinant DNA technology include Watson, J.D. et al 1987; Darnell, J.E. et al 1986; Old, R.W. et al, 1981; Maniatis, T. et al 1989; and Ausubel et al. 1992.

## Figure legends

The invention is further illustrated in the following non-limiting examples and the drawing wherein

5 Figure 1 provides the codon preference of highly expressed proteins in human cells.

Figure 2 illustrates the outline of gp120, gp140, gp150, and gp160 encoding synthetic genes derived from the wild type sequence at the top. Variable (V) and constant (C) regions are shown together with the leader peptide (LP) and the transmembrane spanning domain  
10 (TMD). The approximate nucleotide positions of the restriction enzyme sites are shown. The approximate position of the three restriction enzyme sites dividing the full-length gp160 gene into the three pieces each containing only unique restriction enzyme sites are shown in bold.

15 Figure 3 building of the synthetic gp120 gene. Variable (V) and constant (C) regions are shown together with the leader peptide (LP) and the transmembrane spanning domain (TMD). The approximate nucleotide positions of the restriction enzyme sites are shown.

Figure 4 building of the synthetic gp140 gene. Variable (V) and constant (C) regions are  
20 shown together with the leader peptide (LP) and the transmembrane spanning domain (TMD). The approximate nucleotide positions of the restriction enzyme sites are shown.

Figure 5 building of the synthetic gp150 gene. Variable (V) and constant (C) regions are shown together with the leader peptide (LP) and the transmembrane spanning domain (  
25 TMD). The approximate nucleotide positions of the restriction enzyme sites are shown.

Figure 6 building of the synthetic gp160 gene. Variable (V) and constant (C) regions are shown together with the leader peptide (LP) and the transmembrane spanning domain (  
30 TMD). The approximate nucleotide positions of the restriction enzyme sites are shown.

Figure 7 illustrates the codons coding amino acids in general

Figure 8 illustrates how overlapping PCR is performed.

35 Figure 9 illustrates how PCR using conserved flanking ends is performed.

Figure 10 illustrates how  $S_{1265XhoI}$  is produced using complementary strands (minigene-approach) technology. The  $S_{1265XhoI}$  is ligated from three sets of complementary strands into the vector pBluescript KS<sup>+</sup> between restriction enzyme sites *XhoI* and *PstI*.

5

Figure 11 illustrates how  $S_{1465PstI}$  is produced. The same approach, as the approach used for the production of  $S_{1265XhoI}$ , was used except that only two sets of complementary strands were used.

- 10 Figure 12 illustrates the assembly of  $P_1$ . The  $S_{O-N-Lang}$  and  $S_{235EcoRV}$  are ligated into the *XbaI* and *PstI* site of the  $S_{375PstI}$  containing plasmid.

Figure 13 illustrates the assembly of  $P_2$ . The  $S_{900XbaI}$  was excerted by *HindIII* and *SacI* from its plasmid and ligated with  $S_{990SacI}$  (*SacI-SpeI*) into the  $S_{110SpaI}$  plasmid that was opened at the *HindIII* and *SpeI* sites.

15

Figure 14 illustrates the assembly of  $P_3$ .  $S_{495ClnI}$  (*ClnI-EcoRI*) and  $S_{650-720EcoRI}$  (*EcoRI-XbaI*) and  $P_2$  (*XbaI-XhoI*) were ligated simultaneously into the  $P_1$  plasmid opened at the *ClnI* and *XhoI* sites to obtain the  $P_3$  plasmid.

20

Figure 15 illustrates the assembly of  $P_{4gp160}$ .  $S_{1890HindIII}$  (*SacI-HindIII*) and  $S_{1700EagI}$  (*HindIII-EagI*) were ligated simultaneously into the  $S_{1630XbaI}$  plasmid opened by *SacII* and *EagI*.

Figure 16 illustrates the assembly of  $P_5$ .  $S_{2190ClnI}$  (*ClnI-PstI*) and  $S_{2330PstI}$  (*PstI-EcoRI*) were ligated into the  $S_{2425E8}$  plasmid opened by *ClnI* and *EcoRI*.

25

Figure 17 illustrates the assembly of  $P_{8gp160}$ .  $S_{1465PstI}$  (*XbaI-PstI*) and  $S_{1265XhoI}$  (*PstI-XhoI*) were ligated into the  $P_{4gp160}$  plasmid opened by *XbaI* and *XhoI*.

- 30 Figure 18 illustrates the assembly of  $P_{8gp150}$ .  $S_{1465PstI}$  (*XbaI-PstI*) and  $S_{1265XhoI}$  (*PstI-XhoI*) were ligated into the plasmid containing  $P_{4gp150}$  with the stop codon.  $P_{4gp150}$  plasmid was opened at the *XbaI* and *XhoI* sites for the ligation.

Figure 19 illustrates the assembly of  $P_{8gp140}$ .  $S_{1465PstI}$  (*XbaI-PstI*) and  $S_{1265XhoI}$  (*PstI-XhoI*) were ligated into the plasmid containing  $P_{4gp140}$  with a stop codon.  $P_{4gp140}$  plasmid was opened at the *XbaI* and *XhoI* sites for the ligation.

35

Figure 20 illustrates the assembly of  $P_{8gp41}$ . Two complementary nucleotide strands 1265gp41S and 1265gp41AS designed with overhang creating a 5' *XhoI* and a 3' *PstI* restriction enzyme site were annealed and ligated into the piece 8 which is already opened at the *XhoI* and *PstI* sites whereby  $S_{1265}$  is deleted.

Figure 21 illustrates the assembly of  $P_7$ .  $P_8$  (*XhoI*-*SacII*) and  $S_{2060SacII}$  (*SacII*-*ClaI*) were ligated into  $P_5$  plasmid opened at *XhoI* and *ClaI*.

- 10 Figure 22a SDS PAGE of  $^{35}S$ -labelled HIV-1 BX08 envelope glycoproteins radio-immuno precipitated from transiently transfected 293 cells using the indicated plasmids. Cell pellet (membrane bound antigens) or cell supernatant (secreted antigens) were precipitated by a polyclonal anti-HIV-1 antibody pool. Lane 1: untransfected cells. Lane 2: supernatant from syn.gp120<sub>MN</sub> transfected cells. Lane 3: cell pellet from wt.gp160<sub>BX08</sub> transfected cells.
- 15 Lane 4: cell pellet from cells co-transfected by wt.gp160<sub>BX08</sub> and pRev. Lane 5: Mwt. marker. Lane 6: cell pellet from syn.gp160<sub>BX08</sub> transfected 293 cells. Lane 7: cell pellet from syn.gp150<sub>BX08</sub> transfected 293 cells. Lane 8: supernatant from syn.gp140<sub>BX08</sub> transfected cells. Lane 9: supernatant from syn.gp120<sub>BX08</sub> transfected cells.
- 20 Figure 22b is an SDS-PAGE of  $^{35}S$ -labeled HIV-1 BX08 envelope glycoproteins radio-immune precipitated from transiently transfected 293 cells as cell pellet (membrane bound) or cell supernatant (secreted antigens) by anti-HIV-1 antibody pool using the indicated plasmids. Lane 1: untransfected 293 cells. Lane 2: cell pellet from syn.gp160MN transfected 293 cells as positive control (Vinner et al 1999). Lane 3: Cell supernatant from syn.gp120MN transfected 293 cells as positive control (Vinner et al 1999). Lane 4: Cell supernatant from syn.gp120BX08 transfected 293 cells demonstrating a glycoprotein band of 120 kDa. Lane 5: Cell supernatant from syn.gp140BX08 transfected 293 cells demonstrating a glycoprotein band of 120 kDa. Lane 6: Mwt. marker. Lane 7 at two different exposure times: Cell pellet from syn.gp150BX08 transfected 293 cells
- 30 demonstrating a glycoprotein band of 120 kDa (lower gp30 band is not well seen in this exposure). Lane 8: Cell supernatant from syn.gp150BX08 transfected 293 cells showing no secreted proteins (all protein is membrane bound, see lane 7).

Figure 22c show fluorescent microscopy of U87.CD4.CCR5 cells transfected with BX08 gp160 genes plus pGFP. Panel A: cells transfected with empty WRG7079 vector plus pGFP showing no syncytia. Panel B: cells transfected with wild type BX08gp160 gene



plus pGFP showing some syncytia. Panel C: cells transfected with synBX08gp160 plus pGFP showing extreme degree of syncytia formation. This demonstrates expression, functionality, and tropism of the expressed BX08 glycoprotein with much more expressed functionally active gp160 from the synthetic BX08 gene.

5

Figure 23 shows the anti-Env-V3 BX08 antibody titers (IgG1). Panels show individual mice DNA immunized with syn.gp140BX08 plasmid either i.m. (left panel) or by gene gun (right panel), respectively. Immunization time points are indicated by arrows.

- 10 Figure 24 shows a Western Blotting of (from left to right) one control strip, followed by sera (1:50) from 2 mice i.m. immunized with synBX08gp120, 2 mice i.m. immunized with synBX08gp140, 2 mice i.m. immunized with synBX08gp150, and 2 mice immunized with synBX08gp160, followed by 2 mice gene gun immunized with synBX08gp120, 2 mice gene gun immunized with synBX08gp140, 2 mice gene gun immunized with synBX08gp150, and 2 mice gene gun immunized with synBX08gp160 respectively. Strip 5 is a mouse 5.1 DNA immunized i.m. with synBX08gp140 plasmid (same mouse as in figure 23). Plasma was examined at week 18. The positing of gp160 (spiked with four coupled gp51), gp120 and gp41 is indicated at the right. A positive reaction to HIV glycoproteins further demonstrates the mouse anti-HIV immunoglobulin reacting to HIV of a strain (IIIB) different from BX08 to illustrate cross-strain reactivity.
- 15
- 20

Figure 25 Theoretical example of calculation of the 50% inhibitory concentration ( $IC_{50}$ ) values.  $IC_{50}$  for each mouse serum is determined by interpolation from the plots of percent inhibition versus the dilution of serum.

25

- Figure 26 CTL responses were measured at week 18 to the mouse H-2D<sup>d</sup> restricted BX08 V3 CTL epitope (IGPGGRAFYTT) for BALB/c mice (H-2D<sup>d</sup>) i.m. immunized at week 0, 9, and 15 with the synthetic vaccine genes: syn.gp120<sub>BX08</sub>, syn.gp140<sub>BX08</sub>, syn.gp150<sub>BX08</sub>, and syn.gp160<sub>BX08</sub>, respectively, and median values of different E:T ratios for groups of mice are shown (26A). Intramuscular DNA immunization with syn.gp150<sub>BX08</sub> induced a higher CTL response when injected i.m. in high amounts versus gene gun inoculation of skin (26B).
- 30

- Figure 27 Summary of western immuno blotting assay of mice sera (1:40) collected at week 0, 9, and 18 from mice genetically immunized with syn.gp120<sub>BX08</sub>, syn.gp140<sub>BX08</sub>,
- 35

syn.gp150<sub>BX08</sub>, syn.gp160<sub>BX08</sub>, wt.gp160<sub>BX08</sub>, and wt.gp160<sub>BX08</sub> plus pRev, respectively.  
Percent responders in groups of 17-25 mice against gp120 and gp41 are shown.

Figure 28 IgG anti-rgp120 (IIIB) antibody titers of individual mice inoculated at week 0, 9, 15  
5 (28A), or gene gun immunized at week 3, 6, 9, and 15 (28B) with the syn.gp150<sub>BX08</sub> DNA  
vaccine.

Figure 29 IgG antibody titers to HIV-1 rgp120<sub>IIIB</sub>. Median titers are shown from groups of  
mice i.m. inoculated at week 0, 9, and 15 (29A), or gene gun immunized at week 0, 3, 6,  
10 9, and 15 (29B) with the synthetic genes syn.gp120<sub>BX08</sub>, syn.gp140<sub>BX08</sub>, syn.gp150<sub>BX08</sub>,  
and syn.gp160<sub>BX08</sub>, respectively.

## Examples

### Example 1: Designing the nucleotide sequence construct

Initially the overall layout of the nucleotide sequence construct is decided. The overall layout comprises the various derivatives the gene will be expressed as. For BX08 these include, but  
5 are not restricted to gp160, gp150, gp140, gp120, and gp41.

Next, the vehicle of expression (plasmid or virus) is to be determined: Preparation for a suitable vector determines both need for leader sequence, terminal restriction enzyme sites and whether or not an N- or C-terminal protein tag is to be considered (Poly-his, Myc-antibody-epitop, etc.). For BX08 a plasmid expression vehicle was chosen. All native wild  
10 type HIV codons are systematically exchanged with the codons most frequently represented in a pool of highly expressed human genes (figure 1). By this exchange the amino acid sequence is conserved while the nucleotide sequence is dramatically altered. Thus, gene structures like overlapping reading frames (e.g. *vpu*, *rev*, and *tat*) or secondary structures (e.g. RRE) are most likely destroyed whereas protein cleavage sites, and glycosylation sites  
15 are maintained. The 100% amino acid identity between wtBX08 and synthetic BX08 in the present examples should be calculated after the initial Ala-Ser amino acid sequence, as that sequence is a part of the 6 amino acid sequence long *NheI* restriction enzyme site.

Depending on the restriction enzyme sites located in the expression vector it is decided  
20 which restriction enzyme sites can be present (tolerated) throughout the finished gene construct. The terminal restriction enzyme sites of the synthetic gene must remain unique to enable cloning into the vector chosen for expression. General requirements for restriction enzyme sites of choice: Preferably creating cohesive ends facilitating ligation, creating no compatible ends with adjacent restriction enzyme sites (e.g. *BamHI/BglII*), and being efficient  
25 cutters. For BX08 the restriction enzyme sites accepted were the ones present in the polylinker of the pBluescript cloning vectors (*EagI*, *MluI*, *EcoRV*, *PstI*, *ClaI*, *EcoRI*, *XbaI*, *SacI*, *SpeI*, *XhoI*, *HindIII*, *SacII*, *NotI*, *BamHI*, *SmaI*, *Sall*, *DraI*, *KpnI* with the exception of *BglII* and *NheI*). This was decided to satisfy the original cloning strategy using individual cloning of snuts in pBluescript with restriction enzyme cleaved (trimmed) ends after PCR  
30 amplification, which is not necessary when blunt-end cloning and assembling of complementary oligonucleotides are employed. All locations at which the selected restriction enzyme sites can be introduced by silent mutations (keeping 100% loyal to the amino acid sequence) are identified using the SILMUT software or equivalent.

From these possible restriction enzyme sites, a selection of restriction enzyme sites are introduced by silent nucleotide substitutions around functional regions of choice of the corresponding gene (e.g. RRE) or gene products (e.g. variable region 1 (V1), V2, V3, CD4 binding area, transmembrane domain, and regions of immunological significans, etc.).

- 5 Restriction enzyme sites are located at terminal positions of subcloned snuts (building entities) but additional restriction enzyme sites may be present within subunits. For BX08 the construct was initially to be cloned in the WRG7079 vector containing a tPA-leader sequence. Cloning sites were 5'-*NheI* → *BamHI*-3'. The entire humanised BX08 sequence was divided into thirds: 5'-*NheI* → *XhoI* → *SacII* → *BamHI*-3'. These sites were chosen in
- 10 this particular order because it resembles the polylinker of pBluescript (KS<sup>+</sup>) enabling successive ligations of the assembled thirds in this cloning vector. Within these thirds restriction enzyme sites were kept unique. Next, restriction enzyme sites were placed to flank the functional regions chosen as follows:
  - A. (5'-V1): *EcoRV*-235: Between C1 and V1. Alternatives: 3×*HindIII* (already excluded
  - 15 because exclusive use at position 1890) or *EcoRV*.
  - B. (V1-3'): Only alternative *PstI* 375.
  - C. (5'-V2): as B.
  - D. (V2-3'): Alternatives: *SpeI*, *Clal* 495. *Clal* chosen because it is closer to V2.
  - E. *EcoRI* 650 placed because next possible site was too far away.
  - 20 F. (5'-V3): *BglII* 720 was the alternative closest to the V3 region and further more unique.
  - G. (V3-3'): Alternatives *XhoI* (excluded) and *XbaI* 900 located very close to the V3 loop.
  - H. (5'-V4) *SacI* 990: alternatively *EcoRI* or *BamHI* (both excluded)
  - I. (V4-3'): Alternatives *SpeI* 1110, *KpnI* 1145, *PstI* 1135. *PstI* already used, *SpeI* chosen because of distance to previous site (*SacI* 990).
  - 25 J. (5'-V5): *XhoI* initially determined.
  - K. (Fusion peptide-3') *PstI* 1465 was the closest alternative to *XhoI* 1265.
  - L. (5'-Immunodominant region): *XbaI* 1630 chosen among *EcoRV* (blunt end), *PstI* and *XhoI* (both already used).
  - M. (Immunodominant region-3'): *EagI* 1700 perfect location.
  - 30 N. (C34 and C43 -3' (Chan, Fass, et al. 1997), and 5'-trans membrane domain): *SacII*. No alternatives.
  - O. (Trans membrane domain -3'): *SacII* 2060 already present.
  - P. *Clal* 2190 perfect position in relation to previous RE-site.
  - Q: *PstI* perfect position in relation to previous RE-site.
  - 35 R. *EcoRI* 2400 introduced to facilitate later substitution of terminal snut.
  - S. *BamHI* 2454 determined by the WRG7079 vector.

- Remove undesired restriction enzyme sites by nucleotide substitutions (keeping loyal to the amino acid sequence). Nucleotide substitution should preferably create codon frequently used in highly expressed human genes (figure 1). If that is not possible, the codons should
- 5 be the selected from the regular codons (figure 7). The substitutions made to the second nucleotide sequence to obtain desired restriction enzyme sites are shown in Table 2.

**Table 2 lists silent nucleotide substitutions in the humanised BX08 envelope sequence. Substitutions were made to create or delete restriction enzyme sites.**

Position:	substitution	Remarks:
138	c → g	creates Mlu I site on pos. 134-139
240	c → t	creates EcoRV site on pos. 238-243
501	c → a	creates Cla I site on pos. 501-506
502	a → t	do
503	g → c	do
504	c → g	do
657	c → a	creates EcoRI site on pos. 656-661
660	c → t	do
724	c → a	creates Bgl II site on pos. 724-729
726	c → g	do
727	a → t	do
728	g → c	do
729	c → t	do
840	c → t	EagI site is eliminated
904	a → t	creates Xba I site on pos. 904-909
905	g → c	do
906	c → t	do
907	c → a	do
909	c → a	do
994	a → t	creates Sac I site on pos. 990-995
995	g → c	do
1116	c → t	creates SpeI site on pos. 1114-1119
1119	c → t	do
1273	a → t	creates XhoI site on pos. 1272-1277
1274	g → c	do
1275	c → g	do
1293	c → t	Bgl II site is eliminated
1443	c → t	BstXI site is eliminated
1452	g → c	do
1467	c → t	PstI site on pos. 1466-1471
1470	c → a	do
1590	g → c	PstI site on pos. 1588-1593 is eliminated
1620	g → c	PstI site on pos. 1618-1623 is eliminated
1638	c → t	creates XbaI site on pos. 1638-1643
1641	g → a	do
1653	g → c	PstI site is eliminated
1687	a → t	PstI site is eliminated
1688	g → c	PstI site is eliminated
1710	c → g	creates EagI site on pos. 1709-1714
1758	c → t	Bgl II site is eliminated
1875	g → c	PstI site is eliminated
1893	c → a	Hind III on pos. 1893-1898
1897	c → t	do
1944	c → t	Bgl II site is eliminated

Position:	substitution	Remarks:
2199	c → t	Cla I site on pos. 2198-2203
2202	c → t	do
2203	c → t	do
2253	c → g	SacII site is eliminated
2292	g → c	PstI site is eliminated
2320	a → t	PstI site on pos. 2321-2326 is eliminated
2321	a → t	do
2322	g → t	do
2325	c → a	do
2430	c → a	creates EcoRI site on pos. 2429-2434
2433	c → t	do

### Example 2: synthesis of oligos

- In order to clone the individual snuts, nucleotide strands were synthesised or purchased. In total 28 synthetic nucleotide strands were synthesised. Nucleotide strands were synthesised
- 5 by standard 0.2  $\mu$ mol  $\beta$ -cyanoethyl-phosphoramidite chemistry on an Applied Biosystems DNA synthesiser model 392, employing 2000 Å CPG columns (Cruachem, Glasgow, Scotland), acetonitrile containing less than 0.001% water (Labscan, Dublin, Ireland) and standard DNA-synthesis chemicals from Cruachem, including phosphoramidites at 0.1 M and Tetrahydrofuran/N-methylimidazole as cap B solution. The nucleotide strands O-N-C
- 10 and 119MS-RC (for cloning of snut O-N-Lang), 650-E-BG and 720-XBAC-31 (for cloning of snut 650-720-EcoRI), 2425esup and 2425ESdo (for cloning of snut 2425-E-S) were synthesised with 5' end "trityl on" and purified on "Oligonucleotide Purification Cartridges" (Perkin Elmer, CA, USA) as described by the manufacturer. Other nucleotide strands (235-ECO5, 375-pst1.seq, 495-Cla1.seq, 900-Xba1, 990-sac1, 1110-SPE, 1630-Xba.seq, 1700-
- 15 Eag.seq, 17-Eag.seq, 1890-Hind.MPD, 2060-sac, 2190-cla, 2330-pst) were synthesised with 5' end "trityl off" and purified by standard ethanol precipitation. Oligoes 1265-1UP, 1265-1DO, 1265-2UP, 1265-2DO, 1265-3UP, 1265-3DO, 1465-1UP, 1465-1DO, 1465-2UP, 1465-2DO were purchased from Pharmacia.

### Example 3: Cloning of snuts

- 20 The nucleotide sequence construct was designed in 17 DNA small pieces called snuts (Table 3) encompassing important structures like variable and constant regions each flanked with restriction enzyme (RE) sites to facilitate cassette exchange within each third of the gene: *NheI*-*XhoI*, *XhoI*-*SacII*, *SacII*-*BamHI*.
- 25 Each snut was cloned individually in a commercial vector (pBluescriptKS or pMOSblue) and kept as individual DNA plasmids, named after the snut which gives the nucleotide position of the RE in the BX08.

Table 3 list the snuts by their name and cloning vector.

Name	Cloning vector:
S <sub>O-N-Lang</sub>	pMOSblue
S <sub>235EcoRV</sub>	pMOSblue
S <sub>375PstI</sub>	pBluescriptSK
S <sub>495ClaI</sub>	pMOSblue
S <sub>650-720EcoRI</sub>	pMOSblue
S <sub>900XbaI</sub>	pMOSblue
S <sub>990SacI</sub>	pMOSblue
S <sub>1110SpeI</sub>	pMOSblue
S <sub>1265XhoI</sub>	pBluescriptSK
S <sub>1465PstI</sub>	pBluescriptSK
S <sub>1630XbaI</sub>	pBluescriptSK
S <sub>1700EagI</sub>	pBluescriptSK
S <sub>1890HindIII</sub>	pBluescriptSK
S <sub>2060SacII</sub>	pMOSblue
S <sub>2190ClaI</sub>	pMOSblue
S <sub>2330PstI</sub>	pMOSblue
S <sub>2425ES</sub>	pBluescriptSK

Three principally different methods were used to obtain the dsDNA corresponding to each of  
 5 the 17 snuts needed to build the synthetic BX08 genes.

- 1) "Overlapping" PCR: is based on the use of two ssDNA template nucleotide strands (forward and reverse) that complement each other in their 3'-end (figure 8). During the first PCR cycle, both templates annealed to each other at the 3'-ends allowing the full-length  
 10 polymerisation of each complementary strand during the elongation step. The newly polymerized dsDNA strand are then amplified during the following cycles using an adequate forward and reverse primers set (figure 8).

**Snut O-N-LANG:** (S<sub>O-N-Lang</sub>) two ng of the forward template nucleotide strands O-N-C and 2  
 15 ng of the reward template-nucleotide strand 119MS-RC were mixed together with 50pmoles of the forward primer O-N-LANG-5 (5'-CTAGCTA-GCGCGGCCGACCGCCT -3') and 50pmoles of the reverse primer O-N-LANG-3 (5'-CTCGATATCCTCGTGCATCTGCTC -3') in a 100µl PCR reaction volume containing 0.2mM dNTP's, 1x ExpandHF buffer with MgCl<sub>2</sub> (1.5mM) and 2.6 units of enzyme mix (Expand™ High fidelity PCR system from Boehringer  
 20 Mannheim). The PCR was performed with the PE Amp 9600 thermocycler (Perkin Elmer) using the following cycle conditions: initial denaturation at 94°C for 30 sec., followed by 30 cycles of 94°C for 15 sec., 65°C for 30 sec., 72°C for 45 sec., with a final elongation at 72°C for 5 min., and cooling to 4°C.

**Snut 650-720-EcoRI:** ( $S_{650-720\text{EcoRI}}$ ) PCR amplification was performed as described for snut O-N-LANG. One  $\mu\text{g}$  of the forward ssDNA template-oligonucleotide 650-E-BG and 1  $\mu\text{g}$  of the reverse ssDNA template-oligonucleotide 720-XBAC were mixed with 40 pmoles of the forward primer 650-E-5 (5'-CCGGAATT-CGCCCCGTGGTGAGCA-3') and 40 pmoles of the reverse primer 720-X-3 (5'-CTGCTCTAGAGATGTTGCAGTGGGCCT-3').

2) "Normal" PCR amplification: Eleven nucleotide strands: 235-ECO5, 375-pst1, 900-xba1, 990-sac1, 1110-SPE, 1630-XBA, 1700-EAG, 1890-HIN, 2060-sac, 2190-cla, and 2330-pst, were designed with common 5' and 3' flanking sequences which allowed PCR amplification with the same primer set (Forward primer : BX08-5 (5'-AGCGGATAACAATTTACACAGGA-3') and reverse primer : BX08-3 (5'-CGCCAGGGTTTTCCAGTCACGAC-3') (Figure 9). The 495-Cla1 oligonucleotide was designed without a common flanking sequence and was therefore amplified with a specific set of primers 495-5N/495-3N (5'-GAATCGAT-CATCACCCAG-3' and 5'-GACGAATTCGCTGGGTGCACT-3'). Each oligonucleotide was resuspended in 1 ml of water and kept as a stock solution (approximately 0.2 mM). PCR amplification was performed with the Expand™ High Fidelity PCR System from Boehringer Mannheim (Cat. No. 1759078). Four concentrations of template nucleotide strand were systematically used: undiluted stock solution, stock solution  $10^{-1}$ , stock solution  $10^{-2}$ , stock solution  $10^{-3}$ . One to 5  $\mu\text{l}$  of synthetic ssDNA template was amplified using the following conditions: BX08-5 (0.5  $\mu\text{M}$ ), BX08-3 (0.5  $\mu\text{M}$ ), 4 dNTP's (0.2 mM), 1x ExpandHF buffer with  $\text{MgCl}_2$  (1.5 mM) and 2.6 units of enzyme mix. The PCR was performed using the PE Amp 9600 thermocycler (Perkin Elmer) using the following cycle conditions: initial denaturation at 94°C for 15 sec., followed by 30 cycles of 94°C for 15 sec., 65°C for 30 sec. and 72°C for 45 sec., with a final elongation at 72°C for 7 min., and cooling to 4°C.

25

3) *Minigene approach* : This method was used to synthesise  $S_{1265\text{XhoI}}$ ,  $S_{1465\text{XbaI}}$  and  $S_{2425\text{ES}}$ . **Snut 2425-E-S** ( $S_{2425\text{ES}}$ ): 100 picomoles of each oligonucleotide 2425ES-up (35-mer ; 5'-AATTCGCCAGGGCTTCGAGCGCGCCCTGCTGTAAG-3') and 2425ES-do (35-mer ; GATCCTTACAGCAGGGCGCGCTCGAAGC-CCTGGCG-3') were mixed together in a 100  $\mu\text{l}$  final volume of annealing buffer containing NaCl 25 mM, Tris 10 mM and 1 mM EDTA. After denaturation at 94°C for 15 min., the mixed oligonucleotides were allowed to anneal at 65°C during 15 min.. The annealing temperature was allowed to slowly decrease from the 65°C to room temperature (22°C) during overnight incubation. The resulting double-strand dsDNA fragments harbored EcoRI- and BamHI-restriction sites overhangs that allowed direct cloning in pBluescript KS(+) vector using standard cloning techniques (Maniatis 1996).

35



**Snut 1265-XhoI** ( $S_{1265XhoI}$ ): This snut was built according to the strategy depicted in figure 10. Three minigenes were constructed following the same method described for snut 2425-E-S. These minigenes are named 1265-1, 1265-2 and 1265-3. The minigene 1265-1 results from the annealing of the oligonucleotides 1265-1up (68-mer ; 5'-TCG AGC AGC GGC AAG GAG ATT  
 5 TTC CGC CCC GGC GGC GGC GAC ATGC GCG ACA ACT GGC GCA GCG AGC T-3') and 1265-1do (68-mer ; 5'-GTA CAG CTC GCT GCG CCA GTT GTC GCG CAT GTC GCC GCC GCC GGG GCG G AAA ATC TCC TTG CCG CTG C-3'). 1265-2 results from the annealing of 1265-2up (61-mer ; 5'-GTA CAA GTA CAA GGT GGT GAA GAT CGA GCC CCT GGG CAT CGC CCC CAC CAA GGC CAA GCG C-3') and 1265-2do (63-mer ; 5'-CAC GCG GCG CTT GGC CTT GGT GGG GGC GAT GCC CAG GGG CTC GAT  
 10 CTT CAC CAC CTT GTA CTT-3'). Finally, 1265-3 results from the annealing of 1265-3up (69-mer ; 5'-CGC GTG GTG CAG CGC GAG AAG CGC GCC GTG GGC ATC GGC GCT ATG TTC CTC GGC TTC CTG GGC GCT GCA-3') and 1265-3do (59-mer ; 5'-GCG CCC AGG AAG CCG AGG AAC ATA GCG CCG ATG CCC ACG GCG CGC TTC TCG CGC TGC AC-3'). Each minigene were designed in order to present single strand overhangs at their 5' and 3'- ends that allow easy ligation and XhoI-PstI  
 15 direct cloning into pBlueScript KS+ vector.

**Snut 1465-PstI** ( $S_{1465PstI}$ ): Two minigenes were constructed following the same methode described for snut 2425-E-S. These minigenes are named 1465-1 and 1465-2. The minigene 1465-1 was obtained after annealing of 1465-1up (90-mer : 5'-GGC AGC ACC ATG GGC GCC  
 20 GCC AGC CTG ACC CTG ACC GTG CAG GCC CGC CAG CTG CTG AGC GGC ATC GTG CAG CAG CAG AAC AAC CTG CTG-3') and 1465-1do (98-mer : 5'-CGC GCA GCA GGT TGT TCT GCT GCT GCA CGA TGC CGC TCA GCA GCT GGC GGG CCT GCA CGG TCA GGC TGG CGG CGC CCA TGG TGC TGC CTG CA-3'), whereas minigene 1465-2 results from the annealing of 1465-2up (78-mer ; 5'-CGC GCC ATC GAG GCC CAG CAG CAC CTG CTC CAG CTGA CCG TGT GGG GCA TCA AGC AGC TCC  
 25 AGG CCC GCG TGC TGG CT-3') and 1465-2do (78-mer ; 5'-CTA GAG CCA GCA CGC GGG CCT GGA GCT GCT TGA TGC CCC ACA CGG TCA GCT GGA GCA GGT GCT GCT GGG CCT CGA TGG-3'). Each minigene were designed in order to present single strand overhangs at their 5' and 3'- ends that allow easy ligation and PstI-XbaI direct cloning into pBlueScript KS+ vector using standard cloning techniques (Maniatis) (see figure 11).

#### 30 Example 4: assembly of snuts to pieces.

The snut genes were then assembled into pieces (Table 4) so that unique restriction enzyme sites or mutagenesis can be used within each of these. This strategy will require fewer assemblings for optimal use of the cassette system. The following piece clones were made and kept individually for construction of the synBX08 gp160 gene (Figure 6):

**Table 4 lists pieces by their name and their snut composition.**

Piece name	snut composition	vector
P <sub>1</sub>	S <sub>O-N-LANG</sub> -S <sub>235EcoRV</sub> -S <sub>375PstI</sub>	pBluescriptSK
P <sub>2</sub>	S <sub>900XbaI</sub> -S <sub>990SacI</sub> -S <sub>1110SpeI</sub>	pMOSblue
P <sub>3</sub>	P <sub>1</sub> -S <sub>495ClaI</sub> -S <sub>850</sub> -S <sub>720EcoRI</sub> -P <sub>2</sub>	pBluescriptSK
P <sub>4gp160</sub>	S <sub>1630XbaI</sub> -S <sub>1700EagI</sub> -S <sub>1890HinfI</sub>	pBluescriptSK
P <sub>5</sub>	S <sub>2190ClaI</sub> -S <sub>2330PstI</sub> -S <sub>2425ES</sub>	pBluescriptKS
P <sub>7</sub>	P <sub>8gp160</sub> -S <sub>2060SacI</sub> -P <sub>5</sub>	pBluescriptKS
P <sub>8gp160</sub>	S <sub>1265XhoI</sub> -S <sub>1465PstI</sub> -P <sub>4gp160</sub>	pBluescriptKS

**Piece 1:** The building strategy is shown in figure 12.

**Preparation of the insert DNA:** Five to 15µg of each plasmid O-N-LANG-cl7 and 235-EcoRV-cl5N, respectively, were double-digested by XbaI/EcoRV, and PstI/EcoRV, according to classical RE digestion procedure (Maniatis). The RE digestion products, were agarose gel purified according classical method (Maniatis). All RE digests were loaded on a 3% Nusieve 3:1(FMC), TBE 0.5X agarose gel and submitted to electrophoresis (7 Volts/mm during 2-3hours) until optimal fragment separation. The agarose-band containing the DNA fragments that correspond to the snut's sequence sizes (243-bp for O-N-LANG and 143-bp for 235-EcoRV) were excised from the gel. The DNA was extracted from agarose by centrifugation 20min at 5000g using a spin-X column (Costar cat#8160). **Preparation of the vector:** The snut 375-PstI klon1 was used as plasmid vector. Five µg were digested with XbaI and PstI. Removal of the polylinker XbaI/PstI fragment was performed by classical agarose gel purification, using a 0.9% Seakem-GTG agarose, TBE 0.5X gel. The linearised plasmid DNA was extracted from the agarose by filtration through spin-X column. All purified DNA fragments were quantified by spectrophotometry. **Ligation:** All three DNA fragments O-N-LANG (XbaI/EcoRV), 235-EcoRV (PstI/EcoRV) and 375-PstI(XbaI/PstI), were ligated together by classical ligation procedure, using an equimolar (vector:insert1:insert2) ratio of 1:1:1. Thus for, 200 ng (0.1 pmole) of XbaI/PstI-linearised 375-PstI-cl1 were mixed with 16 ng of O-N-LANG (XbaI/EcoRV) and 10 ng of 235-EcoRV (PstI/EcoRV) in a final reaction volume of 20µl of 1X ligation buffer containing 10U of T4 DNA ligase (Biolabs, cat#202S). The ligation was allowed overnight at 16°C. **Transformation:** Competent XL1-Blue bacteria (Stratagene cat#200130, transformation efficiency > 5•10<sup>6</sup> col/µg) were transformed by classical heat-shock procedure : 1/10th of the pre-chilled ligation reaction was mixed with 50µl of competent bacteria. The mixture was allowed to stand in ice during 30 min. Bacteria were heat-shocked at 42°C during 45 sec. and then left 2 min. on ice before being resuspended in 450 µl of SOC medium. Transformed bacteria were incubated 1 hour at 37°C under shaking (250rpm) and plated on LB-ampicilin agar plates. The recombinant clones

- were allowed to grow 16 hours at 37°C. Colony screening: 10 to 50 recombinant colonies were screened by direct PCR screening according to the protocole described into the pMOSBlue blunt-ended cloning kit booklet (RPN 5110, Amersham). Each colony was picked and resuspended in 50µl of water. DNA was freed by a boilling procedure (100°C, 5 min). Ten µl of bacterial lysate were mixed to 1 µl of a 10mM solution of premixed 4 dNTP's , 1 µl of M13reverse primer (5pmoles/µl, 5'-CAGGAAACAGCTATGAC-3'), 1 µl of T7 primer (5pmoles/µl, 5'-TAATACGACTCACTATAGGG-3'), 5µl of 10x Expand HF buffer 2 (Boehringer Mannheim, cat#1759078), 0.5µl of Enzyme mix (Boehringer Mannheim, 5U/µl) in a final volume of 50µl. DNA amplification was performed with a thermo-cycler PE9600 (Perkin-Elmer) using the following cycling parameters: 94°C, 2min, 35 cycles(94°C, 30sec; 50°C, 15sec; 72°C,30sec); 72°C, 5min; 4°C hold. Five µl of the PCR products were analysed after electrophoresis on a 0.9% SeakemGTG , 0.5xTBE agarose gel. Nucleotide sequence confirmation: ds-DNA was purified from minicultures of the selected clones with the JETstar mini plasmid purification system ( Genomed Inc.). Sequencing was performed using M13reverse and T7 primers and with the Big Dye<sup>TM</sup> Terminator Cycle Sequencing Ready reaction kit (Perkin-Elmer, Norwalk, Connecticut, P/N4303152) and the ABI-377 automated DNA sequenator (Applied Biosystems, Perkin-Elmer,Norwalk, Connecticut). Data were processed with the Sequence Navigator and Autoassembler softwares (Applied Biosystems, Perkin-Elmer,Norwalk, Connecticut).
- Piece 2:** The strategy for building that piece is depicted in figure 13. RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described above for piece1, except the following:
- The linearised plasmid 1110-SpeI-cl24M1 was used as vector after being digested by HindIII and SpeI, and agarose gel purified.
  - A 166-bp HindIII/SacI, obtained from snut 900-XbaI-cl15, as well as a 130-bp SacI/SpeI fragment, obtained from snut 990-SacI-cl14, were agarose gel purified.
  - Equimolar amount (0.1 pmoles) of the three DNA fragments described above were ligated in an one step ligation.
  - 100µl of competent SCS110 bacteria (Stratagene cat# 200247) were transformed with 1/10th of the ligation products according to the manufactor instruction.
  - Direct colony PCR screening was performed using T7 primer and pMOS-R (5'- GTTGTAACGACGGCCAG-3').

**Piece 3:** The strategy for building that piece is depicted in figure 14. RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described above for piece1, except the following:

- The plasmid piece1-cl33 was linearised by *Clal* and *XhoI*, in order to be used as vector,  
5 and agarose gel purified.
- A 161-bp *Clal*/*EcoRI* fragment, obtained from 495-*Clal*-cl135M1 as well as a 254-bp *EcoRI*/*XbaI* fragment, obtained from 650-720-*EcoRI*-cl39, and a 374-bp *XbaI*/*XhoI* fragment, obtained from piece2-cl4, were agarose gel purified.
- Equimolar amount (0.1 pmole) of each of these 4 DNA fragments were mixed and ligated  
10 together.
- 50µl of competent XL1Blue bacteria were transformed with 1/10th of the ligation products according to the protocole described for piece1.
- Direct colony PCR screening was performed using M13Reverse and T7 primers.

15 **Piece 4 gp160:** The strategy for building that piece is depicted in figure 15. RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described above for piece1, except the following:

- The plasmid 1630-*XbaI*-cl2 was linearised by *SacII*/*EagI* digestion and agarose gel  
20 purified, in order to be used as vector.
- A 190-bp *EagI*/*HindIII* fragment, obtained from snut 1700-*EagI*-cl4, as well as a 177-bp *SacII*/*HindIII* fragment, obtained from snut 1890-*HindIII*-cl8, were agarose gel purified.
- Equimolar amount (0.1 pmoles) of the three DNA fragments described above were ligated in an one step ligation.
- 25 - 50µl of competent XL1Blue bacteria were transformed with 1/10th of the ligation products according to the protocole described for piece1.
- Direct colony PCR screening was performed using M13Reverse and T7 primers.

**Piece 4-gp150:** PCR-based site-directed mutagenesis was performed on double-stranded  
30 plasmid-DNA from piece4-cl4 according an adaptation of the ExSite™ PCR-Based Site-Directed Mutagenesis Kit procedure (Stratagene cat#200502)(Weiner, M.P., Costa, G.L., Schoettlin, W., Cline, J., Marthur, E., and Bauer, J.C. (1994) *Gene* 151:119-123). The mutations introduced are shown in bold letters in the primer sequences below. PCR amplification was performed with the Expand™ High Fidelity PCR System (Boehringer  
35 Mannheim, cat#1759078). Briefly, 1.5 µg, 0.5 µg or 0.1 µg of circular dsDNA was mixed with 1.5 pmoles of P4M2S (5'-TCTGGAAGCTCAGGGGGCTGCATCCCTGGC-3') and 1.5

pmoles of P4M2AS (5'-CCCGCCTGCCCCGTGTGACGGATCCAGCTCC-3') in a final volume of 50µl containing 4 dNTPs (250µM each), 1x Expand HF buffer 2 (Boehringer Mannheim, cat#1759078), 0.75µl of Enzyme mix (Boehringer Mannheim, 5U/µl). The PCR was performed with a PE9600 thermo-cycler (Perkin-Elmer Corporation) under the following cycling parameters : 94°C, 2min ; 15 cycles (94 °C, 45 sec ; 68 °C, 4 min); 72°C, 7min and 4 °C, hold. PCR products were phenol:chloroform extracted and precipitated (Maniatis). Plasmid template was removed from PCR products by DpnI treatment (Biolabs)(Nelson, M., and McClelland, M., 1992) followed by ethanol-precipitation. Amplicons were resuspended in 50 µl steril water, and phosphorylated according the following procedure: 7.5 µl of amplicons were mixed with 0.5µl of 100mM DTT, 1 µl of 10x pk buffer and 1 µl of pk mix enzyme (pMOSBlue blunt-ended cloning kit, Amersham , cat#RPN 5110). DNA kinasing was allowed 5min at 22°C. After heat-inactivation (10min, 75°C) of the pk enzyme, 1µl of ligase (4units, Amersham , cat#RPN 5110) was directly added to the pk reaction. The ligation was allowed overnight at 22°C. 50µl of competent XL1Blue bacteria were transformed with 1/10th of the ligation reaction according to the classical protocol (Maniatis). Insertion of mutations was checked by sequencing.

**Piece 4-gp140:** PCR-based site-directed mutagenesis was performed on piece 4-cl4, according to the procedure described for piece4-gp150 except that the primers P4M1AS (5'-TGTGTGACTGATTGAGGATCCCCAACTGGC-3') and P4S (5'-AGCTTGCCCACTTGTCCAGCTGGAGCAGGT-3') were used.

**Snut 1265-XhoI-gp120 :** PCR-based site-directed mutagenesis was performed on plasmid 1265-XhoI-cl2M1 according to the procedure described for piece4-gp150 except that the primers 1265MAS (5'-CTTCTCGCGCTGCACCACGCGGCGCTTGGC-3') and 1265M2S (5'-CGCGCCTAGGGCATCGGCGCTATGTTCTC-3') were used.

**Snut 1265-XhoI-gp160/uncleaved :** PCR-based site-directed mutagenesis was performed on plasmid 1265-XhoI-cl2M1 according to the procedure described for piece4-gp150 except that the primers 1265MAS (5'-CTTCTCGCGCTGCACCACGCGGCGCTTGGC-3') and 1265M2S (5'-AGCGCCGTGGGCATCGGCGCTATGTTCTC-3') were used.

**Snut 1465-PstI-CCG :** PCR-based site-directed mutagenesis was performed on plasmid 1465-PstI-cl25 according to the procedure described for piece4-gp150 except that the primers 1465MAS (5'-CTGCTTGATGCCCCACACGGTCAGCTG-3') and 1465MS (5'-TGCTGCGGCCGCGTGCTGGCTCTAGA-3') were used.

**Piece 5 :** The strategy for building that piece is depicted in figure 16. RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described above for piece1, except the following:

- 5 - The plasmid 2425-ES-cl2 was linearised by Clal/EcoRI digestion and agarose gel purified, in order to be used as vector.
- A 129-bp PstI/Clal fragment, obtained 2190-Clal-cl6M15, as well as a 114-bp PstI/EcoRI fragment, obtained from 2330-PstI-cl8, were agarose gel purified.
- Equimolar amount (0.1 pmole) of the three DNA fragments described above were ligated
- 10 in an one step ligation.
- 50µl of competent XL1Blue bacteria were transformed with 1/10th of the ligation products according to the protocole described for piece1.
- Direct colony PCR screening was performed using T3 (5'-ATTAACCCTCACTAAAG-3') and T7 primers.

15

**piece 8 :** The strategy for building that piece is depicted in figure 17. RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described above for piece1, except the following:

- The plasmid piece4-cl4 was linearised by XbaI/XhoI and agarose gel purified, in order to be
- 20 used as vector.
- A 200-bp XhoI/PstI fragment, obtained from 1265-XhoI-cl2M1 as well as a 178-bp PstI/XbaI fragment, obtained from 1465-PstI-cl25 were agarose gel purified.
- Equimolar amount (0.1 pmole) of these 3 DNA fragments were mixed and ligated together.
- 50µl of competent XL1Blue bacteria were transformed with 1/10th of the ligation products
- 25 according to the protocole described for piece1.
- Direct colony PCR screening was performed using T3 and T7 primers.

**piece 8-gp150 :** The strategy for building that piece was identical to that of piece 8, except that piece4-cl4M3 was used as vector (figure 18).

30

**piece8-gp150/ uncleaved :** The strategy for building that piece is identical to that of piece 8, except that piece4 gp160-cl4M3 is used as vector and a 200-bp XhoI/PstI fragment, obtained from snut 1265-XhoI-gp160/uncleaved as well as a 178-bp PstI/XbaI fragment, obtained from snut 1465-PstI-CCG are used like inserts.

35

**piece 8-gp140** : The strategy for building that piece was identical to that of piece 8, except that piece4-cl4M5 was used as vector (figure19).

**piece8-gp140/ uncleaved** : The strategy for building that piece is identical to that of piece 8, except that piece4-cl4M5 is used as vector and a 200-bp XhoI/PstI fragment, obtained from snut 1265-XhoI-gp160/uncleaved as well as a 178-bp PstI/XbaI fragment, obtained from snut 1465-PstI-CCG are used like inserts.

**Piece8-gp41** : The strategy for building that piece is depicted in figure 20. RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described for synBX08 gp160 gene. A 63 bp linker is to be made according to the method described for snut 2425-ES, the minigene approach. Thus for 2 complementary oligonucleotides: 1265-gp41S(5'-TCGAGgctagcGCCGTGGGCATCGGCGCTATGTTCTCGGCTTCCTGGGCGctgca-3') and 1265-gp41AS (5'-gCGCCCAGGAAGCCGAGGAAC-ATAGCGCCGATGCCACGGCgtagcC-3' should be annealed together. This synthetic linker will be directly ligated into the XhoI /PstI sites of piece8-klon13 from which the snut 1265-XhoI-cl 2M1 would have been removed.

- piece7** : The strategy for building that piece is depicted in figure 21. RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described above for piece1, except the following:
- The plasmid piece5-cl1 was linearised by ClaI/XhoI and agarose gel purified, in order to be used as vector,.
  - A 798-bp XhoI/SacII fragment, obtained from piece8-cl13 as well as a 140-bp SacII/ClaI fragment, obtained from 2060-SacII-cl21 were agarose gel purified.
  - The ligation of the 3 fragments was performed using a vector:insert ratio of 1:1, 1:2 or 1:5.
  - 50µl of competent XL1Blue bacteria were transformed with 1/10th of the ligation products according to the protocole described for piece1.
  - Direct colony PCR screening was performed using M13Reverse and T7 primers.

#### **Example 5: assembly of genes**

**synBX08 gp160 gene** : The strategy for building that gene is depicted in figure 6. RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described for piece1.

20 µg of the expression plasmid WRG7079 were digested by NheI/BamHI. Plasmid DNA-

ends were dephosphorylated by Calf Intestin Phosphatase treatment (CIP, Biolabs) (Maniatis) to avoid autoligation of any partially digested vector. CIP enzyme was heat-inactivated and removed by classical phenol-chloroforme extraction. A 1277-bp NheI/XhoI fragment, obtained from piece3-cl27, as well as a 1194-bp XhoI/BamHI fragment, obtained from piece7-cl1, were agarose gel purified. The ligation was performed using a vector:insert ratio of 1:1 or 1:2. Fifty µl of competent XL1Blue bacteria were transformed with 1/10th of the ligation product according to the protocole described for piece1. After transformation bacteria were plated on LB-kanamycin agar plates. Direct PCR colony screening was performed using the primer set WRG-F (5'-AGACATAATAGCTGACAGAC-3') and WRG-R (5'-GATTGTATTTCTGTCCCTCAC-3'). The nucleotide sequence was determined according to the methods described above for piece1.

**synBX08 gp150 gene** : The strategy for building that gene is depicted in figure 5. RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described for synBX08 gp160 gene. A 1277-bp NheI/XhoI fragment, obtained from piece3-cl27, as well as a 800-bp XhoI/BamHI fragment, obtained from piece 8-gp150-cl26, were agarose gel purified and then ligated into the NheI/BamHI WRG7079 sites. The ligation was performed using a vector:insert ratio of 1:1 or 1:2.

For construction of the synthetic BX08 gp150, piece4 was mutated to Piece4gp150 whereby a tyrosine -> cysteine was changed and a stop codon was introduced after the transmembrane spanning domaine ( TMD), followed by a BamHI cloning site. A new piece8gp150 was constructed composed of snut1265/snut1465/piece4gp150.

**synBX08 gp150/uncleaved gene** : RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies are performed according the same procedures described for synBX08 gp160 gene. A 1277-bp NheI/XhoI fragment, obtained from piece3-cl27, as well as a 800-bp XhoI/BamHI fragment, obtained from piece 8-gp150/uncleaved, are agarose gel purified and then are ligated into the NheI/BamHI WRG7079 sites. The ligation was performed using a vector:insert ratio of 1:1 or 1:2.

**synBX08 gp 140 gene** : The strategy for building that gene is depicted in figure 4. RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described for synBX08 gp160 gene. A 1277-bp NheI/XhoI fragment, obtained from piece3-cl27, as well as a 647-bp XhoI/BamHI fragment, obtained from piece 8-gp140-cl2, were agarose gel purified



and then ligated into the *NheI*/*Bam*HI sites of WRG7079. The ligation was performed using a vector:insert ratio of 1:1 or 1:2. For construction of the synthetic BX08 gp140, piece4 was mutated to Piece4gp140 whereby a stop codon was introduced just before the *TMD* followed by a *Bam*HI cloning site. A new piece8gp140 was constructed composed of  
5 snut1265/1465/piece4gp140.

**synBX08 gp140/uncleaved gene** : RE digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies are performed according the same procedures described for synBX08 gp160 gene. A 1277-bp *NheI*/*XhoI* fragment, obtained  
10 from piece3-cl27, as well as a 800-bp *XhoI*/*Bam*HI fragment, obtained from piece 8-gp140/uncleaved, are agarose gel purified and then ligated into the *NheI*/*Bam*HI WRG7079 sites. The ligation was performed using a vector:insert ratio of 1:1 or 1:2.

**synBX08 gp 120 gene** : The strategy for building that piece is depicted in figure 3. RE  
15 digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described for synBX08 gp160 gene. A 1277-bp *NheI*/*XhoI* fragment, obtained from piece3-cl27, as well as a 206-bp *XhoI*/*Bam*HI fragment, obtained from 1265-*XhoI*-gp120-clM5, were agarose gel purified and then ligated into the *NheI*/*Bam*HI sites of WRG7079. The ligation was performed  
20 using a vector:insert ratio of 1:1 or 1:2. For construction of the synthetic BX08 gp120, snut 1265 was mutated to S<sub>1265gp120</sub> to introduce a stop codon at the gp120/gp41 cleavage site followed by a *Bam*HI cloning site.

The gp160, gp150, gp140, and gp120 genes are cloned (*NheI*-*Bam*HI) and maintained in an  
25 eucaryotic expression vectors containing a CMV promotor and a tPA leader, but other expression vectors may be chosen based on other criteria e.g. antibiotic resistance selection, other leader sequences like CD5 etc, presence or not of immune stimulatory sequences etc.

**SynBX08 gp41 gene** : The strategy for building that gene is depicted in figure 20. RE  
30 digestion, DNA fragments purification, ligation as well as direct PCR screening of recombinant colonies were performed according the same procedures described for synBX08 gp160 gene. Piece 8-gp41 is ligated with snut 2060-SacII-klon21 and piece 5 as already decribed for the construction of piece 7, creating piece 7-gp41 (P<sub>7gp41</sub>). Subsequently the piece 7-gp41 containing the entire gp41 gene will be cloned in WRG7079 using the *NheI*  
35 and *Bam*HI sites.

**Example 6a: High expression by codon optimization**

To analyse the expression of glycoproteins from the wild type and synthetic BX08 envelope genes RIPA was performed on transfected mammalian cell lines. Both cell membrane associated and secreted HIV-1 glycoproteins from the cell supernatants were assayed. The

5 envelope plasmids were transfected into the human embryonic kidney cell line 293 (ATCC, Rockville, MD) or the mouse P815 (H-2D<sup>b</sup>) cell line using calcium phosphate (CellPfect Transfection kit, Pharmacia). For radio immune precipitation assay (RIPA), transfected cells were incubated overnight, washed twice and incubated for 1 hour with DMEM lacking cysteine and methionine (Gibco). Then the medium was replaced with medium containing 50

10  $\mu$ Ci per ml of [<sup>35</sup>S] cysteine and 50  $\mu$ Ci/ml of [<sup>35</sup>S] methionine (Amersham Int., Amersham, UK) and incubation continued overnight. Cells were centrifuged, washed twice with HBSS and lysed in 1 ml ice-cold RIPA buffer (10 mM Tris, pH 7.4, 150 mM NaCl, 50 mM EDTA, 1% Nonidet P-40, 0.5% sodiumdeoxycholate) to detect *membrane bound* Env glycoproteins. The cell lysates were centrifuged for 15 min. at 100,000  $\times$  g to remove any undissolved particles

15 and 100  $\mu$ l immune precipitated with protein A-sepharose coupled human polyclonal IgG anti-HIV antibodies (Nielsen et al., 1987). For analysis of secreted Env glycoproteins 500  $\mu$ l of the 5 ml supernatants from transfected cells were incubated with protein A-sepharose coupled human polyclonal IgG anti-HIV antibodies. After washing three times in cold RIPA buffer and once in PBS, the immuno precipitates were boiled for 4 min. in 0.05 M Tris-HCl,

20 pH 6.8, 2% SDS, 10% 2-mercaptoethanol, 10% sucrose, 0.01% bromophenol blue and subjected to SDS PAGE (Vinner et al., 1999). Electrophoresis was carried out at 80 mV for 1 hour in the stacking gel containing 10% acrylamide, and at 30 mV for 18 hours in the separating gradient gel containing 5-15% acrylamide. Gels were fixed in 30% ethanol-10% acetic acid for 1 hour, soaked for 30 min. in En3Hance (Dupont #NEF 981), washed 2  $\times$  15

25 min. in distilled water, dried and autoradiography performed on Kodak XAR-5 film.

Transfection of human 293 cells with the syn.gp120<sub>BX08</sub> and syn.gp140<sub>BX08</sub> genes, respectively, resulted in high amounts of only secreted HIV-1 glycoproteins (Fig. 22a, lane 9 and 8). Thus, the synthetic gene in the absence of *rev* expresses the HIV-1 surface glycoprotein of the expected size which is recognised by human anti-HIV-1 antisera. The

30 expression of BX08 gp120 was Rev independent and with roughly the same high amount of gp120 from the syn.gp120MN gene (Fig. 22a, lane 2). Fig. 22a, lane 6 and lane 7 shows the expression of only membrane bound gp160 and gp150 from 293 cells transfected with syn.gp160<sub>BX08</sub> and syn.gp150<sub>BX08</sub> plasmids, respectively. Also transfection with wt.gp160<sub>BX08</sub> plasmid resulted in a significant expression of membrane bound gp160 despite the absence

35 of Rev (Fig. 22a, lane 3). Co-transfection with equimolar amounts of Rev encoding plasmid seemed to increase this expression somewhat (Fig. 22a, lane 4). This is seen despite the

lower transfection effectivity using two plasmids and the use of only half the amount of wt.gp160<sub>BX08</sub> DNA when combined with pRev. The amounts of secreted HIV-1 glycoproteins from gp120 and gp141 accumulating in the cell supernatants seemed higher than the amounts of cell associated glycoproteins at the time of harvesting of the cells. Interestingly, 5 the amounts of gp160 produced from the "humanized" gene were about equal to the amounts produced by the wt.gp160<sub>BX08</sub> + pRev genes, respectively (Fig. 22a, lane 4 and 6). The processing of gp160, gp150 and gp140 into gp120 plus a gp41, or fractions of gp41, produced from wild type or synthetic genes in the 293 cell-line did not function well under these experimental conditions. Same phenomenon was seen in RIPA from 293-CD4 cells 10 and HeLa-CD4 cells infected by HIV-1<sub>MN</sub> (Vinner et al., 1999). Because of the absence of CCR5 these cell-lines could, however, not be infected by HIV-1 strain BX08.

**Example 6b: Radio immuno precipitation assay (RIPA) of synthetic BX08 transfected cells showing expression of glycoproteins from synthetic BX08 env plasmids**

15 The synthetic envelope plasmid DNA were transfected into the human embryonic kidney cell line 293 (ATCC, Rockville, MD) using calcium phosphate (CellPfect Transfection kit, Pharmacia). For immune precipitation analysis, transfected 293 cells were treated and analysed according to the method described in example 6a.

To analyse expression from these genes, an SDS-PAGE of the <sup>35</sup>S-labelled HIV-1 20 envelopes, immune precipitated from the transfected cells is shown in figure 22b. Both cell-membrane associated and secreted HIV-1 envelope glycoproteins in the cell supernatants were assayed. Transfection of 293 cells with the synthetic BX08 gene encoding gp120 (syn.gp120BX08) in lane 4, and syn.gp140BX08 (lane 5) that did not contain rev encoding regions, resulted in abundant amounts of HIV-1 gp120. Thus, the expressions were Rev 25 independent and expressed in roughly same high amounts as the syn.gp120MN and syn.gp160MN genes (lane 3 and 2, respectively) already showed by our group and others to be markedly increased in comparison with HIV MN wild type genes including rev (Vinner et al 1999).

Transfection with syn.gp150 plasmid (lanes 7 and 8) resulted in significant expression of 30 membrane associated gp120 and low detectable amounts of truncated form of gp41 (cell pellet in lane 7) with no detectable HIV-1 glycoprotein in the cell supernatant lane 8. It is concluded that the synthetic BX08 genes express the envelope glycoproteins of expected size which are recognised by human anti-HIV-1 antiserum.

**Exampel 6C FACS**

To quantitate the surface expression of HIV glycoproteins from the wild type and synthetic BX08 envelope genes transfection experiments were done and cell surface expression  
5 examined by FACS (flow cytometer).

10  $10 \mu\text{g}$  of the BX08 envelope plasmid (wild type BX08gp160 or synBX08gp160) plus  $10 \mu\text{g}$  of an irrelevant carrier plasmid pBluescript were used to transfect a 80-90% confluent layer of 293 cells in tissue culture wells ( $25 \text{ cm}^2$ ) using the CellPect kit (Pharmacia). After 48 hours  
15 cells were Versene treated, washed and incubated with a mouse monoclonal IgG antibodies to HIV gp120 (NEA-9301, NEN<sup>TM</sup>, Life-Science Products Inc., Boston) for time 30 min. on wet ice followed by washing in PBS, 3% FCS and incubation with Phyto-Erytrin (PE) labelled rat anti-mouse IgG1 (Cat #346270, Becton Dickinson) according to the manufacture. After washing the cells were fixed in PBS, 1% paraformaldehyd, 3% FCS, and analysed on a  
20 FACS (FACScan, Becton-Dicknsson). Table 5 show in duplicate expression of BX08 gp160 from 11 % of the cells transfected with wild type BX08 (number 1 and 2) compared to the 48 % of cells expression BX08 glycoprotein when transfected with the synthetic gene (number 3 and 4). Thus, a several fold higher expression is obtained using the synthetic BX08 gene.

20 **Table 5** FACS analysis of 293 cells transfected with synBX08gp160 (No 1 and 2) and wt.gp16+BX08 (No3 and 4) and stained with monoclonal antibodies to surface expressed HIV glycoproteins. A higher expression was obtained with the synthetic gene (mean 48%) as compared to the wild type gene (mean 11 %).

	50 ul	45 ul	A	B	C	C - A	C - B
1	syn.gp160BX08 +	pBluescript SK+	2,57	2,85	36,91	34,34	34,06
2	syn.gp160BX08 +	pBluescript SK+	2,83	2,14	58,42	55,59	56,28
3	wt.gp160BX08 +	pBluescript SK+	1,95	1,52	7,51	5,56	5,99
4	wt.gp160BX08 +	pBluescript SK+	2,97	1,42	14,41	11,44	12,99

25 A: No primary antibody added (control for unspecific secondary Ab binding)  
B: Neither Primary Ab nor Secondary Ab added (autoflourescence control)  
C: Primary Ab and secondary Ab added.

**Example 6D Analyses of the surface expression and biological functionality**

To analyse the surface expression and biological functionality from the wild type and  
30 synthetic BX08 envelope genes transfection experiments were done and cell fusion microscopically studied using HIV envelope receptor expressing cells.

10 µg of the BX08 envelope plasmid (wt.BX08gp160 or syn.BX08gp160 or empty WRG7079 vector plasmid) plus 5 µg of a plasmid (pEGFP, Clontech) expressing green fluorescent protein (GFP) were transfected into  $2 \times 10^6$  adherent U87.CD4.CCR5 cells (NIH AIDS Res. & Reference program, catalog #4035) stably expressing CD4 and CCR5, using the CellPfect  
5 transfection kit (Pharmacia). After 48 hours the cells were examined by microscopy and photographed (Fig 22c).

Fig 22c panel A show the negative control (empty WRG7079 plus pGFP) giving no syncytia. Panel B show cells transfected with the wild type BX08 gp160 plasmid where cell-to-cell fusion (syncytia) is seen. Panel C show cells transfected with the same amounts of  
10 synBX08gp160 plasmid and demonstrating a much higher degree of cell-cell fusion. In fact most or all of the cells in the culture plate were fused at this time. This experiment show surface expression of functional HIV gp160 with tropism to the CCR5 receptor, as well as a much higher expression and biological activity from the synthetic BX08 gene as compared to the wild type equivalent.

15

#### Example 7: Gene inoculation of mice for immunization

6-7 weeks old female BALB/c mice were purchased from Bomholdtgaard, Denmark. Microbiological status was conventional and the mice were maintained in groups of 4/5 per cage with food and water ad libitum and artificially lighted 12 hours per day. Acclimatization  
20 period was 2 days. Mice were anaesthetized with 0.2 ml i.p. of rohypnol:stesolid (1:3, v/v) and DNA inoculated by either i.m. injection of 50 µl 2 mg/ml of plasmid DNA in each tibia anterior muscle at week 0, 9, and 15 and terminated week 18; or gene gun inoculated on shaved abdominal skin using plasmid coated gold particles (0.95 µm particles, 2 µg DNA/mg gold, 0.5 mg gold/shot, 50-71% coating efficiency) with the hand held Helios® gene gun  
25 device (BioRad) employing compressed (400 psi) Helium as the particle motive force. Mice were gene gun vaccinated at week 0, 3, 6, 9, 15, and terminated week 18.

#### Example 8: Serological assays

**Western blotting.** The induction of a humoral response to gp120 and gp41 antigens by *in vivo* expression of the encoded glycoproteins from the synthetic BX08 genes was examined  
30 by western immuno blotting (Figure 27). Mouse antisera (1:40) were evaluated in western blotting using the commercial HIV BLOT 2.2 strips (Genelabs Diagnostic). The conjugate was a 1:200 dilution of the alkaline phosphatase-conjugated rabbit anti-mouse IgG (Dakopatts, Glostrup, Denmark). Buffers, incubation condition and colour development were

used according to the manufacturer. In these western blotting strips the gp160 band from HIV-1 IIIB contain of an oligomeric form of gp41 in a higher concentration than the monomeric gp41 band on the strip (Genelabs Diagnostic). HIV-1<sub>IIIB</sub> lysate is used in these commercial strips where the gp160 band is composed by addition of tetrameric gp41. All

5 preimmune sera tested negative in western blotting. Mice inoculated with syn.gp120<sub>BX08</sub> showed antibody response to the heterologous gp120 of HIV-1 IIIB. Inclusion of the extracellular part of gp41 in the gene syn.gp140<sub>BX08</sub> induced antibody reaction to both gp120 and gp41 in all mice. This confirms the *in vivo* expression of BX08 gp120 and the extracellular part of gp41. DNA vaccination with syn.gp160<sub>BX08</sub> encoding the membrane

10 bound glycoprotein induced antibodies to gp120 and gp41 in 50% and 64% of the mice, respectively. DNA vaccination with syn.gp150<sub>BX08</sub> induced detectable antibodies to gp120 and gp41 in 41% and 53%, respectively. Induction of different levels of antibodies could explain the difference in numbers of positive reactive mice sera in this qualitative western blotting.

15 **ELISA.** Mouse anti HIV-1 gp120 antibodies were measured by indirect ELISA. Briefly, wells of polystyrene plates Maxisorb (Nunc) were coated for 2 days at room temperature with HIV-1 IIIB recombinant gp120 (Intracel) at 0.2 µg/100 µl of carbonate buffer, pH 9.6. Before use the plates were blocked 1 hour at room temperature with 150 µl/well of washing buffer (PBS, 0.5 M NaCl, 1% Triton-X-100) plus 2% BSA and 2% skim milk powder. After 3 x 1 min.

20 washings, mouse plasma was added at 100 µl/well diluted in blocking buffer and ELISA plates incubated for 90 min. at room temperature using a microtiter plate shaker. As standard curve we used a mouse monoclonal antibody to a conserved part of gp120 between V5-C5 (MRDNWRSELYKY) (#NEA-9301, NEN™ Life Science Products, Inc., Boston, MA). As calibration control included on each plate we used a plasma pool from 10 mice vaccinated

25 with BX08 gp120. Plates were again washed 5 x 1 min. and incubated 1 hour at room temperature with 100 µl/well of HRP-conjugated rabbit anti-mouse IgG (#P260, Dakopatts, Glostrup, Denmark) diluted 1:1000 in blocking buffer. Colour was developed with 100 µl/well of peroxidase enzyme substrate consisting of 4 mg of o-phenylenediamine in 11 ml water plus 4 µl hydrogen peroxide (30%, w/w). The enzyme reaction was terminated after 30 min.

30 by 150 µl/well of 1M H<sub>2</sub>SO<sub>4</sub>. The optical density (OD) of wells was measured at 492 nm using a microplate photometer (Molecular Devices, Biotech-Line, Denmark). Anti-HIV-gp120 IgG titers were expressed as the reciprocal plasma dilution resulting in an OD<sub>492nm</sub> value of 0.500. Mouse anti-HIV-1 BX08 antibodies were also measured by indirect peptide ELISAs as described above using a BX08 V3 peptide (SIHIGPGRAFYTTGD) (Schafer, Copenhagen,

35 Denmark).

The IgG antibody response to HIV-1<sub>IIIIB</sub> rgp120 quantitated by ELISA is seen in Fig. 28 and Fig. 29. No background activity was observed in preimmune sera or in sera from 4 mice immunized with empty WRG7079 vector in parallel with the BX08 genes. All mice inoculated with the synthetic BX08 genes either by gene gun or by i.m. injection responded and showed a persistent and high titered (about 100-10,000) IgG response to rgp120 as exemplified in Fig. 4. When comparing the median titers for groups of mice (Fig. 29) a moderate antibody response was observed with the wt.gp160<sub>BX08</sub>. Intramuscular and gene gun immunization with a mixture of wt.gp160<sub>BX08</sub> plasmid plus Rev encoding plasmid did not increase this antibody response. This was found even when both plasmids were coated onto the same gold particles to ensure co-transfection of single target cells. However, to ensure inoculation of equal amounts of total DNA only half of the amount of wt.BX08 plasmid was used when mixing with pRev which may have contributed to the lower antibody response when pRev was included. A 5-fold improvement of the antibody response was obtained using the syn.gp160<sub>BX08</sub> gene. This antibody response seemed further improved using the syn.gp150<sub>BX08</sub> gene where the cytoplasmic internalization signals were eliminated but only using gene gun inoculation. For both the gene gun inoculation of skin and i.m injection the highest antibody titers to rgp120 were induced by genes encoding *secreted* gp120/gp140 glycoproteins versus *membrane bound* gp150/gp160 glycoproteins, respectively. In general, equal antibody and ELISA titers to rgp120 were obtained using gene gun and i.m. injection of the BX08 vaccine genes.

#### Example 9: Neutralization assay

Mouse plasma was diluted in culture medium (RPMI-1640 medium (Gibco) supplemented with antibiotics (Gibco), Nystatin (Gibco) and 10% FCS (Bodinco)) and heat inactivated at 60°C for 30 min. Of the HIV-1 strain BX08 (50 TCID<sub>50</sub> per ml propagated in PBMC) 250 µl was incubated for 1 hour at room temperature with 250 µl dilution of mouse serum (four five-fold dilutions of mouse serum, final dilutions 1:20 to 1:2500). After incubation 1 × 10<sup>6</sup> PBMC in 500 µl culture medium was added to the virus-serum mixture and incubated overnight at 37°C in 5% CO<sub>2</sub>. Subsequently, eight replicates of 10<sup>5</sup> PBMC in 200 µl culture medium were cultured in 96-well culture plates (Nunc) at 37°C in 5% CO<sub>2</sub>. After seven days in culture the concentration of HIV antigen in the culture supernatant was quantitated using HIV antigen detection ELISA (Nielsen et al., 1987).

This ELISA is performed using human IgG, purified from high titered patient sera, both as capture antibody and biotin-linked as detecting antibody. In brief, anti-HIV-capture IgG diluted 1:4000 in PBS, 100 µl/well, are coated onto Immunoplates (Nunc) overnight at 4°C. After washing five times in washing buffer 100 µl of supernatants are applied and incubated overnight

at 4°C. Plates are washed 5 times before incubation with 100 µl HIV-IgG conjugated with biotin diluted 1:1000 in dilution buffer, plus 10% HIV-1 sero-negative human plasma for 3 hours at room temperature. Five times 1 min. washing in washing buffer are followed by 30 min. incubation with 100 µl of 1:1000 avidine-peroxidase (Dako P347 diluted in dilution buffer). Six times 1 min. washings, 5 in washing buffer and the last one are done in dH<sub>2</sub>O before colour is developed with 100 µl of peroxidase enzyme substrate consisting of 4 mg of OPD in 11 ml water plus 4 µl hydrogen peroxide (30 %, w/w). The enzyme reaction is terminated after 30 minutes by additional 150 µl of 1M H<sub>2</sub>SO<sub>4</sub>.

The HIV antigen concentration in cultures, preincubated with mouse serum, was expressed relatively to cultures without mouse serum (culture medium), and the percentage inhibitions of the different dilutions of mouse serum were calculated. The 50% inhibitory concentration (IC<sub>50</sub>) for each mouse serum was determined by interpolation from the plots of percent inhibition versus the dilution of serum, and the neutralizing titer of the serum was expressed as the reciprocal value of the IC<sub>50</sub>. In each set-up a human serum pool known to neutralise other HIV-1 strains was included in the same dilutions as the mouse serum as a calibrating control. For assay of neutralization of the heterologous SHIV89.6P the MT-2-cell-killing format was used (Crawford et al., 1999). The assay stock of SHIV89.6P was grown in human PBMC.

The neutralizing IC<sub>50</sub> antibody titers of plasma pools from 10 mice from each group were measured at different time points (week 0, 9, and 18). A positive background in some preimmune sera and thus in all week 0 serum pools was noted even after dilution and heat inactivation that was found earlier to lower this background. In general the neutralizing titers to BX08 virus of such serum pools were transient and low ranging from 1:6-1:150 above background (data not shown). A possible cross-neutralization reaction to a heterologous, primary HIV-1 envelope was tested using the SHIV89.6P which is relevant in macaque models of AIDS and serum pools from mice DNA immunized i.m. with syn.gp140<sub>BX08</sub>. Preimmune serum had a titer of 1:37, which is indicative of a slightly positive background, whereas the 18 week p.i. serum had a positive neutralizing titer of 1:254 above background.

### 30 Example 10: CTL assay

The cellular immune response in mice following gene gun or i.m. genetic immunization with the different vaccine plasmids were examined (Fig. 26). Spleen was removed aseptically and gently homogenised to single cell suspension, washed 3 times in RPMI-1640 supplemented with 10% FCS and resuspended to a final concentration of  $5 \times 10^7$  cell/ml. The cells were then incubated 5 days with mitomycin-C treated (50 µg/ml for 1 hour) mouse P815 (H-2D<sup>d</sup>)



stimulator cells at a ratio of 10:1 in medium supplemented with  $5 \times 10^{-5}$  M  $\beta$ -mercaptoethanol. For assay of CTL response to HIV-1 BX08, P815 stimulator cells and target cells were pulsed with 20  $\mu$ g/ml of the HIV-1 BX08 V3 peptide containing a conserved murine H-2D<sup>d</sup> restricted CTL epitope (IGPGRAFYTT) (Lapham et al., 1996). After stimulation, splenocytes were washed three times with RPMI-1640 supplemented with 10% FCS and resuspended to a final concentration of  $5 \times 10^6$  cells/ml. 100  $\mu$ l of cell suspension was added in triplicate to U-bottom 96-well microtiter plates and a standard 4 hour <sup>51</sup>Cr-release assay performed (Marker et al., 1973).

All synthetic BX08 plasmids induced a high specific CTL response thus confirming the *in vivo* expression and *in vivo* immunogenicity. The highest CTL response was obtained with syn.gp150<sub>BX08</sub> followed by syn.gp120<sub>BX08</sub>-syn.gp140<sub>BX08</sub>, and syn.gp160<sub>BX08</sub>, respectively. Thus, the CTL response induced did not correlate with the antigen being secreted or not. However, i.m. DNA immunization with syn.gp150<sub>BX08</sub> containing six putative CpG motifs induced a higher CTL response than gene gun immunization (Fig. 26). This difference could be explained by the high amount of DNA used in the i.m. injections.

The T-lymphocyte cytokine profile of spleen cells after ConA stimulation as well as serum antibody IgG<sub>2a</sub>/IgG<sub>1</sub> at week 18 were investigated. Neither the IFN $\gamma$ /IL-4 nor the IgG<sub>2a</sub>/IgG<sub>1</sub> ratios, which both reflects a Th1-type of immune response, were significantly higher for the i.m. immunized mice when compared with gene gun immunized mice (student t-test and Mann-Withney U-test). Thus, the CTL response did not correlate with a certain Th-type of response and the DNA immunization technique did not bias the immune response using synthetic BX08 genes.

#### **Example 11: Antibody responses to DNA vaccination with synBX08 env plasmid**

A relatively low and variable antibody response (1 of 10 mice) was obtained with gene gun inoculation of the syn.gp140BX08 plasmid vaccine starting at week 9, figure 23, right panel. A higher numbers of responders 3/10 with high IgG1 antibody responses at an earlier onset (week 3-9) was obtained with the syn.gp140BX08 plasmid using i.m. injection, left panel. Sera from later time points may show more responders and/or higher titers but are not assayed. However, these results show the induction of an antibody response to the BX08 V3 peptide by DNA vaccination using one of the described synthetic BX08 constructs.

## References

- Webster RG, Robinson HL. DNA vaccines: A review of developments. *Biopharmaceuticals* 1997, 4:273-292.
- 5 Rosenberg ES et al. Vigorous HIV-1 specific CD4+ T cell responses associated with control of viremia. *Science* 1997, 278:1447-1450.
- Boyer J et al. *Nature Med* 1997, 3:526-532.
- 10 Choe H et al. The beta-chemokine receptors CCR3 and CCR5 facilitate infection by primary isolates. *Cell* 1996, 85(7):1135-48.
- Dragic T et al. Co-receptors: gateways to the cell. *HIV advances in Research and Therapy* 1997 (9): 2-12.
- 15 Karlsson AC et al. Characterization of the viral population during primary HIV-1 infection. *AIDS* 1998, 12:839-847.
- Haas J, Park EC, Seed B. Codon usage limitation in the expression of HIV-1 envelope glycoprotein. *Curr. Biol.* 1996, 6:315-324.
- 20 André S, Seed B, Eberle J et al. Increased immune response elicited by DNA vaccination with a synthetic gp120 sequence with optimized codon usage. *J. Virol.* 1998, 72: 1497-1503.
- 25 Letvin NL et al. Potent protective anti-HIV immune responses generated by bimodal HIV envelope DNA plus protein vaccination. *PNAS* 1997, 94: 9378-9383.
- Bryder K et al. Improved humoral and cellular immune responses against the gp120 V3 loop of HIV-1 following genetic immunization with a chimeric DNA vaccine encoding the V3 inserted into the hepatitis B surface antigen. *Scand J Immunol* 1998 Apr, 47(4):289-95.
- 30 Kwong PD, et al. Structure of an HIV gp120 envelope glycoprotein in complex with the CD4 receptor and a neutralizing human antibody. *Nature* 1998 393: 648-659.
- Wyatt R et al. The antigenic structure of the HIV gp120 envelope glycoprotein. *Nature* 1998, 35 393: 705-711.

- Sauter MS, et al. An internalization signal in the SIV transmembrane protein cytoplasmic domain modulates expression of envelope glycoproteins on the cell surface. *J. Cell Biol.* 1996, 132: 795-811.
- 5
- Mascola JR, et al. Potent and synergistic neutralization of HIV-1 primary isolates by hyperimmune anti-HIV immunoglobulin combined with monoclonal antibodies 2F5 and 2G12. *J. Virol.* 1997, 71(10): 7198-7206.
- 10
- Molecular Cloning : A Laboratory Manual With the Lab Manual Source Book 1996 - Sambrook, J./Fritsch, E.F./Maniatis, T.
- Nielsen CM; Bygbjerg IC; Vestergaard BF. Detection of HIV antigens in eluates from whole blood collected on filterpaper, *Lancet*, 1987 Mar 7, 1(8532):566-7.
- 15
- Harada-S, Koyanagi Y; Yamamoto N, Infection of HTLV-III/LAV in HTLV-I-carrying cells MT-2 and MT-4 and application in a plaque assay. *Science*, 229(4713):563-6 1985 Aug 9.
- Verrier FC et al. Antibodies to several conformation-dependent epitopes of gp120/gp41
- 20 inhibit CCR-5-dependent cell-to-cell fusion mediated by the native envelope glycoprotein of a primary macrophage-tropic HIV-1 isolate. *Proc Natl Acad Sci USA*, 1997 Aug 19, 94(17):9326-31.
- Chan DC et al. Core structure of gp41 from the HIV envelope glycoprotein; *Cell*, 1997 Apr
- 25 18, 89(2):263-73.
- Weiner, M.P., Costa, G.L., Schoettlin, W., Cline, J., Marthur, E., and Bauer, J.C. *Gene* 1994, 151:119-123.
- 30 Nelson, M., and McClelland, M. *Methods Enzymol.*, 1992, 216: 279-303
- Cloning Vectors: A Laboratory manual*, P.H. Pouwels et al. 1985, supp. 1987.
- Watson, J.D. et al. *Molecular Biology of the Gene*, Volumes I and II, the Benjamin/Cummings
- 35 Publishing Company Inc, Menlo Park, Calif, 1987.

- Darnell, J.E. et al. *Molecular Cell Biology*, Scientific American Books, New York (1986).
- Old, R.W. et al, *principles of Gene Manipulation: An Introduction to Genetic Engineering*, 2nd edition, University of California press, 1981.
- 5 Ausubel et al. *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, 1992.
- Wilbur, W.J. and Lipman, D.J. Rapid similarity searches of nucleic acid and protein data banks, *Proc Natl Acad Sci USA*, 1983, 80:726-730.
- 10 Vinner L, H. V. Nielsen, K. Bryder, S. L. Corbet, C. Nielsen, and A. Fomsgaard. Gene gun DNA vaccination with Rev-independent synthetic HIV-1 gp160 envelope gene using mammalian codons. *Vaccine*. 1999 Apr 23, 17(17):2166-75.
- Marker, O. and Volkert M. Studies on cell-mediated immunity to lymphocyte choriomeningitis virus in mice. *J. Exp. Med.* 1973, 137:1511-1525.
- 15 Lapham, C., B. Golding, J. Inman, R. Blackburn, J. Manischewitz, P. Highet, and H. Golding. *Brucella abortus* conjugated with a peptide derived from the V3 loop of human immunodeficiency virus (HIV) type 1 induces HIV-specific cytotoxic T-cell responses in normal and in CD4+ cell-depleted BALB/c mice. *J. Virol.* 1996, 70(5):3084-3092.
- 20 Crawford, J. M., P. L. Earl, B. Moss, K. A. Reimann, M. S. Wyand, K. H. Manson, M. Bilska, J. T. Zhou, C. D. Pauza, P. W. H. I. Parren, D. R. Burton, J. G. Sodroski, N. L. Letvin, and D. C. Montefiori. Characterization of primary isolate-like variants of simian-human immunodeficiency virus. *J. Virol.* 1999, 73(12):10199-10207.

## Claims

1. A method for producing a nucleotide sequence construct comprising the following steps:
  - a) obtaining a first nucleotide sequence of an HIV gene from a patient within the first 12  
5 months of infection;
  - b) designing a second nucleotide sequence utilising the most frequent codons from mammalian highly expressed proteins to encode the same amino acid sequence as the first nucleotide sequence of a) encodes
  - c) redesigning the second nucleotide sequence of b) so that restriction enzyme sites  
10 surrounds the regions of the nucleotide sequence which encode functional regions of the amino acid sequence and so that selected restriction enzyme sites are removed thereby obtaining a third nucleotide sequence encoding the same amino acid sequence as the first and the second nucleotide sequence of a) and b) encode;
  - d) redesigning the third nucleotide sequence of c) so that the terminal snuts contain  
15 convenient restriction enzyme sites for cloning into an expression vehicle;
  - e) producing the snuts between restriction enzyme sites of c) and terminal snuts of d);
  - f) assembling the snut of step e) to form a nucleotide sequence construct.
2. A method according to claim 1, wherein the HIV gene is the gene encoding the envelope.  
20
3. A method according to claim 1 or 2, wherein the HIV gene encodes one or more Gag proteins.
4. A method according to any of the preceding claims, wherein the HIV in step a) is in group  
25 M, O or N
5. A method according to claim 4, wherein the HIV is a group M virus.
6. A method according to any of the preceding claims, wherein the HIV is subtype A, B, C, D,  
30 E, F, G, H, I, or J.
7. A method according to claim 6, wherein the HIV is subtype B.
8. A method according to any of the preceding claims wherein the first nucleotide sequence  
35 is obtained by direct cloning.

9. A method according to any of the preceding claims, wherein the HIV in step a) is isolated with the first 11 months of infection, such a 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, or 0.5 month after infection.
- 5
10. A method according to any of the preceding claims, wherein the redesigning in step c) is carried out after the second nucleotide sequence of step b) has been divided into pieces, so that each piece comprises only different restriction enzyme sites.
- 10 11. A method according to claim 10, wherein the second nucleotide sequence of step b) is divided into 9 pieces, or 8, or 7, or 6, or 5, or 4, or 3, or 2 pieces.
12. A method according to claim 11, wherein the second nucleotide sequence of step b) is divided into 3 pieces.
- 15
13. A method according to any of the preceding claims, wherein the second nucleotide sequence of step b) is designed utilising the most frequent codons from human highly expressed proteins to encode the same amino acid sequence as the first nucleotide sequence of step a) encodes.
- 20
14. A nucleotide sequence construct obtainable by the method of any of claims 1-13.
15. A nucleotide sequence construct according to claim 14, wherein the nucleotide sequence encoding the amino acid sequence in the first variable region is surrounded by *EcoRV* and *PstI* restriction enzyme sites.
- 25
16. A nucleotide sequence construct according to claims 14 or 15, wherein the nucleotide sequence encoding the amino acid sequence in the second variable region is surrounded by *PstI* and *ClaI* restriction enzyme sites.
- 30
17. A nucleotide sequence construct according to any of claims 14-16, wherein the nucleotide sequence encoding the amino acid sequence in the third variable region is surrounded by *ClaI* and *EcoRI* restriction enzyme sites.

18. A nucleotide sequence construct according to any of claims 14-17, wherein the nucleotide sequence encoding the amino acid sequence in the transmembrane spanning region is surrounded by *HindIII* and *SacII* restriction enzyme sites.
19. A nucleotide sequence construct according to any of claims 14-18, wherein the nucleotide sequence encoding the amino acid sequence on both sites of the cleavage site is surrounded by *PstI* and *XbaI* restriction enzyme sites.
20. A nucleotide sequence construct in isolated form which has a nucleotide sequence with the general formula (I), (II), (III), or (IV) or subsequences thereof
- (I)  $P_1-S_{495ClaI}-S_{650-720EcoRI}-P_2-S_{1265gp120}$
  - (II)  $P_1-S_{495ClaI}-S_{650-720EcoRI}-P_2-S_{1265XhoI}-S_{1465PstI}-P_{4gp140}$
  - (III)  $P_1-S_{495ClaI}-S_{650-720EcoRI}-P_2-S_{1265XhoI}-S_{1465PstI}-P_{4gp150}$
  - (IV)  $P_1-S_{495ClaI}-S_{650-720EcoRI}-P_2-S_{1265XhoI}-S_{1465PstI}-P_{4gp160}-S_{2060SacII}-P_5$
- wherein  $P_1$  designates the nucleotide sequence SEQ ID NO: 41, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 90% thereto;
- wherein  $S_{495ClaI}$  designates the nucleotide sequence SEQ ID NO: 7, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 95% thereto;
- wherein  $S_{650-720EcoRI}$  designates the nucleotide sequence SEQ ID NO: 9, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 95% thereto;
- wherein  $P_2$  designates the nucleotide sequence SEQ ID NO: 43, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 85% thereto;
- wherein  $S_{1265gp120}$  designates the nucleotide sequence SEQ ID NO: 19, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 70% thereto;
- wherein  $S_{1265XhoI}$  designates the nucleotide sequence SEQ ID NO: 17, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 80% thereto;
- wherein  $S_{1465PstI}$  designates the nucleotide sequence SEQ ID NO: 23, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 90% thereto;

- wherein P<sub>4gp140</sub> designates the nucleotide sequence SEQ ID NO: 57, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 85% thereto;
- wherein P<sub>4gp150</sub> designates the nucleotide sequence SEQ ID NO: 55, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 85% thereto;
- wherein P<sub>4gp160</sub> designates the nucleotide sequence SEQ ID NO: 53, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 85% thereto;
- 10 wherein S<sub>2060SacII</sub> designates the nucleotide sequence SEQ ID NO: 33, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 98% thereto; and
- wherein P<sub>5</sub> designates the nucleotide sequence SEQ ID NO: 59, a nucleotide sequence complementary thereto, or a nucleotide sequence with a sequence identity of at least 85% thereto.
- 15
21. A nucleotide sequence construct according to claim 20, with the formula (I)
- (I) P<sub>1</sub>-S<sub>495ClaI</sub>-S<sub>650-720EcoRI</sub>-P<sub>2</sub>-S<sub>1265gp120</sub>
22. A nucleotide sequence construct according to claim 20, with the formula (II)
- (II) P<sub>1</sub>-S<sub>495ClaI</sub>-S<sub>650-720EcoRI</sub>-P<sub>2</sub>-S<sub>1265XhoI</sub>-S<sub>1465PstI</sub>-P<sub>4gp140</sub>
23. A nucleotide sequence construct according to claim 20, with the formula (III)
- (III) P<sub>1</sub>-S<sub>495ClaI</sub>-S<sub>650-720EcoRI</sub>-P<sub>2</sub>-S<sub>1265XhoI</sub>-S<sub>1465PstI</sub>-P<sub>4gp150</sub>
- 25
24. A nucleotide sequence construct according to claim 20, with the formula (IV)
- (IV) P<sub>1</sub>-S<sub>495ClaI</sub>-S<sub>650-720EcoRI</sub>-P<sub>2</sub>-S<sub>1265XhoI</sub>-S<sub>1465PstI</sub>-P<sub>4gp160</sub>-S<sub>2060SacII</sub>-P<sub>5</sub>
25. A nucleotide sequence construct according to claim 20 consisting essentially of the
- 30 subsequence P<sub>1</sub>.
26. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence S<sub>495ClaI</sub>.
27. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence S<sub>650-720EcoRI</sub>.



28. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence P<sub>2</sub>.
- 5 29. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence S<sub>1265gp120</sub>.
30. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence S<sub>1285XhoI</sub>.
- 10 31. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence S<sub>1465PstI</sub>.
32. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence P<sub>4gp140</sub>.
- 15 33. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence P<sub>4gp150</sub>.
34. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence P<sub>4gp160</sub>.
- 20 35. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence S<sub>2060SacII</sub>.
- 25 36. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence P<sub>5</sub>.
37. A nucleotide sequence construct with a sequence identity of more than 85% to the nucleotide sequence construct in any of claims 20-35.
- 30 38. A nucleotide sequence construct according to claim 37, wherein the sequence identity is more than 90% such as more than 95%, 98%, or 99%.
- 35 39. A nucleotide sequence construct according to claim 37, wherein the sequence identity is 100%.

40. A nucleotide sequence construct according to any of claims 14-39, coding for an HIV envelope or parts thereof with an improved immunogenicity obtained by mutating the nucleotide sequence construct of any of claims 14-39 such that one or more glycosylation sites in the amino acid sequence have been removed.
41. A nucleotide sequence construct according to claim 40 with a mutation at positions A307C + C309A and/or A325C + C327G and/or A340C + C342A and/or A385C + C387A and/or A469C + C471A or any combination of those.
42. A nucleotide sequence construct according to any of claims 14-41, coding for an HIV envelope or parts thereof with a binding site for the CXCR4 co-receptor in the third variable region.
43. A nucleotide sequence construct according to claim 42 with a mutation at positions G865C + A866G.
44. A nucleotide sequence construct according to any of claims 14-43, coding for an HIV envelope or parts thereof, wherein an immunodominant epitope has been modified.
45. A nucleotide sequence construct according to claim 44, wherein an immunodominant epitope in the third variable region has been modified.
46. A nucleotide sequence construct according to claim 45 with a deletion of nucleotides 793-897.
47. A nucleotide sequence construct according to claim 44, wherein an immunodominant epitope has been removed from gp41.
48. A nucleotide sequence construct according to any of claims 14-47, coding for an HIV envelope or parts thereof, wherein the cleavage site between gp41 and gp120 is removed.
49. A nucleotide sequence construct according to claim 48 with a mutation at position C1423A.

50. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence P<sub>1</sub>, S<sub>495</sub>ClaI, S<sub>850-720</sub>EcoRI, and P<sub>2</sub>.
51. A nucleotide sequence construct according to claim 20 consisting essentially of the  
5 subsequence S<sub>1265</sub>XhoI, S<sub>1485</sub>PstI, and P<sub>4gp140</sub>.
52. A nucleotide sequence construct according to claim 20 consisting essentially of the subsequence S<sub>1265</sub>XhoI, S<sub>1485</sub>PstI, P<sub>4gp160</sub>, S<sub>2080</sub>SacII, and P<sub>5</sub>.
- 10 53. A nucleotide sequence construct according to any of claims 14-52, further comprising a nucleotide sequence repeat coding for a functional region of the amino acid sequence.
54. A nucleotide sequence construct according to claim 53, wherein the nucleotide sequence repeat codes for amino acids in the third variable region.
- 15 55. A nucleotide sequence construct according to any of claim 14-54, further comprising a nucleotide sequence coding for a T-helper cell epitope containing sequence.
56. An expression vehicle selected from a group of viral vectors consisting of simliki forest virus (sfv), adenovirus and Modified Vaccinia Virus Ankara (MVA), further comprising a  
20 nucleotide sequence construct according to any of claim 14-55.
57. A method of individualised immunotherapy wherein the virus from a newly diagnosed patient is directly cloned, the envelope is produced with highly expressed codons, inserted  
25 into any of the nucleotide sequence constructs of claims 14-55, and administered to the patient.
58. Use of a nucleotide sequence construct according to any of claims 14-55 in medicine.
- 30 59. Use of a nucleotide sequence according to any of claims 14-55 for the manufacture of a vaccine for the prophylactics of infection with HIV in humans.
60. Use of a nucleotide sequence according to any of claims 14-55 for the manufacture of a composition for the treatment of an HIV infection in a human within 24 weeks of primary  
35 infection.

61. Use of the nucleotide sequence according to any of claims 14-55 for the production of a recombinant protein.

1/33

<b>Amino acid</b>	<b>One letter amino acid code</b>	<b>Three letter amino acid code</b>	<b>Codon</b>
<b>Alanine</b>	A	Ala	GCC
<b>Arginine</b>	R	Arg	CGC
<b>Asparagine</b>	N	Asn	AAC
<b>Aspartic acid</b>	D	Asp	GAC
<b>Cysteine</b>	C	Cys	TGC
<b>Glutamine</b>	Q	Gln	CAG
<b>Glutamic acid</b>	E	Glu	GAG
<b>Glycine</b>	G	Gly	GGC
<b>Histidine</b>	H	His	CAC
<b>Isoleucine</b>	I	Ile	ATC
<b>Leucine</b>	L	Leu	CTG
<b>Lysine</b>	K	Lys	AAG
<b>Proline</b>	P	Pro	CCC
<b>Phenylalanine</b>	F	Phe	TTC
<b>Serine</b>	S	Ser	AGC
<b>Threonine</b>	T	Thr	ACC
<b>Tyrosine</b>	Y	Tyr	TAC
<b>Valine</b>	V	Val	GTG

Fig. 1

2/33

# Synthetic BX08 Env Strategy for building the full-length gp160 and derived truncated forms




Fig. 2

3/33

# Building of Synthetic BX08 Env ( gp120 )




Fig. 3

4/33




Fig. 4

Fig. 4



5/33

# Building of Synthetic BX08 Env ( gp150 )




Fig. 5

6/33




Fig. 6

STARTS: ATG

7/33

aa  $\Sigma$  codons

A	Ala	GCX	GCT	GCC	GCG	GCA
C	Cys	TGY	TGT	TGC		
D	Asp	GAY	GAT	GAC		
E	Glu	GAR	GAG	GAA		
F	Phe	TTY	TTT	TTC		
G	Gly	GGX	GGT	GGC	GGG	GGA
H	His	CAY	CAT	CAC		
I	Ile	ATH	ATT	ATC	ATA	
K	Lys	AAR	AAG	AAA		
L	Leu	YTX	TTG	TTA	CTT	CTC
M	Met	ATG	ATG			
N	Asn	AAV	AAT	AAC		
P	Pro	CCX	CCT	CCC	CCG	CCA
Q	Gln	CAR	CAG	CAA		
R	Arg	MGX	CGT	CGC	CGG	CGA
S	Ser	WSX	TCT	TCC	TCG	TCA
T	Thr	ACX	ACT	ACC	ACG	ACA
V	Val	GTX	GTT	GTC	GTG	GTA
W	Trp	TGG	TGG			
Y	Tyr	TAY	TAT	TAC		
.	.	TRR	TGA	TAG	TAA	
X	?					

	5'	2nd				3'
		T	C	A	G	
1st	T	Phe	Ser	Tyr	Cys	T
		Phe	Ser	Tyr	Cys	C
		Leu	Ser	.	.	A
		Leu	Ser	.	Trp	G
	C	Leu	Pro	His	Arg	T
		Leu	Pro	His	Arg	C
		Leu	Pro	Gln	Arg	A
		Leu	Pro	Gln	Arg	G
	A	Ile	Thr	Asn	Ser	T
		Ile	Thr	Asn	Ser	C
		Ile	Thr	Lys	Arg	A
		Met	Thr	Lys	Arg	G
	G	Val	Ala	Asp	Gly	T
		Val	Ala	Asp	Gly	C
		Val	Ala	Glu	Gly	A
		Val	Ala	Glu	Gly	G

3rd

Fig. 7

8/33

## Overlapping PCR Strategy: Snut O-N-LANG (249 bases)




Fig. 8

9/33

# PCR approach using conserved flanking sequences




Fig. 9

10/33

## Snut 1265 by minigene approach




Fig. 10

11/33

# Snut 1465- minigene approach









Fig. 11

12/33

: synBX08-piece1



inserts :



vector :




Fig. 12



13/33

synBX08-piece2




Fig. 13

14/33

synBX08-piece3




Fig. 14

synBX08-piece4gp160





Fig. 15

16/33



17/33

Figure 17 : synBX08-piece8gp160




Fig. 17

synBX08-piece8-gp150




Fig. 18

synBX08-piece8-gp140




Fig. 19

20/33

Figure 20: synBX08-piece8-gp41




Fig. 20



synBX08-piece 7




Fig. 21

22/33




Fig. 22 A

23/33




Fig. 22 B

24/33

Panel A




Fig. 22 C

25/33

Panel B



Fig. 22 C

26/33

Panel C




Fig. 22 C

27/33

IgG1 anti-V3 BX08 from Balb/c mice vaccinated with envelope genes in WRG7079





Fig. 23

Fig. 24



28/33






Fig. 25

30/33

Fig. 26 A CTL activity induced by different synthetic DNA vaccines




Fig. 26 B CTL activity induced by different DNA delivery methods.




Fig. 26

31/33

Percentage of  
antiserum positive to  
gp120<sub>IIIb</sub> and gp41<sub>IIIb</sub>  
in western blot assay

Week					
			gp120	gp41	
syn.gp120 <sub>BX08</sub>	0	0	0	0	18
	9	65	0	0	90
syn.gp140 <sub>BX08</sub>	0	0	0	0	100
	65	95	0	0	100
syn.gp150 <sub>BX08</sub>	0	0	0	0	41
	30	41	0	0	53
syn.gp160 <sub>BX08</sub>	0	0	0	0	50
	32	44	0	0	64
wt.gp160 <sub>BX08</sub>	0	0	0	0	53
	nd	nd	0	0	48
wt.gp160 <sub>BX08</sub> /pRev	0	0	0	0	5
	nd	nd	0	0	55

Fig. 27

32/33

Fig. 28 A




Fig. 28 B




Fig. 28

33/33

Fig. 29 A





Fig. 29 B



## SEQUENCE LISTING

&lt;110&gt; Statens Serum Institut

<120> Method for producing an nucleotide  
sequence construct with optimised codons for an HIV envelope  
based on a primary, clinical HIV isolate and the BX08  
construct.

&lt;130&gt; 21924DK1

&lt;160&gt; 76

&lt;170&gt; FastSEQ for Windows Version 3.0

&lt;210&gt; 1

&lt;211&gt; 243

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)...(243)

&lt;400&gt; 1

gct agc ggc ggc gac cgc ctg tgg gta acc gta tac tac ggc gta ccc  
Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro

gtg tgg aag gac ggc acc acc acc acc acc acc acc acc acc acc acc  
Val Trp Lys Asp Ala Thr Thr Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys

gac tac gac acc gac gta cac aac gta tgg gac acc cac ggc tgc gta  
Ala Tyr Asp Thr Gln Val His Asn Val Trp Ala Thr His Ala Cys Val

ccc acc gac ccc aac ccc cag gac gta gta gta gta gta gta gta gta  
Pro Thr Asp Pro Asn Pro Gln Gln Val Val Leu Gly Asn Val Thr Gln

aac ttc aac atg ggc aag aac aac atg gta gta gta gta gta gta gta gta  
Asn Phe Asn Met Gly Lys Asn Asn Met Val Gln Met His Gln Asp

atc  
Ile

<210> 2

&lt;211&gt; 81

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;400&gt; 2

Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro

<210> 2

&lt;211&gt; 81

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;400&gt; 2

Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro

<210> 2

&lt;211&gt; 81

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;400&gt; 2

Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro

<210> 2

&lt;211&gt; 81

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
                   20                  25                  30  
 Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
                   35                  40                  45  
 Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
                   50                  55                  60  
 Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp  
 65                  70                  75                  80  
 Ile

<210> 3  
 <211> 143  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(143)

<400> 3  
 gat atc atc agc ctg tgg gac cag agc ctg aag ccc tgc gtg aag ctg 48  
 Asp Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu  
   1                  5                  10                  15  
 acc ccc ctg tgc gtg acc ctg aac tgc acc aag ctg aag aac agc acc 96  
 Thr Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr  
                   20                  25                  30  
 gac acc aac aac acc cgc tgg ggc acc cag gag atg aag aac tgc ag 143  
 Asp Thr Asn Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys  
                   35                  40                  45

<210> 4  
 <211> 47  
 <212> PRT  
 <213> Artificial Sequence

<400> 4  
 Asp Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu  
   1                  5                  10                  15  
 Thr Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr  
                   20                  25                  30  
 Asp Thr Asn Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys  
                   35                  40                  45

<210> 5  
 <211> 132  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(132)

<400> 5  
 ctg cag ctt caa cat cag cac cag cgt gcg caa caa gat gaa gcg cga 48  
 Leu Gln Leu Gln His Gln His Gln Arg Ala Gln Gln Asp Glu Ala Arg

1	5	10	15	
gta cgc cct gtt cta cag cct gga cat cgt gcc cat cga caa cga caa				96
Val Arg Pro Val Leu Gln Pro Gly His Arg Ala His Arg Gln Arg Gln				
	20	25	30	
cac cag cta ccg cct gcg cag ctg caa cac atc gat				132
His Gln Leu Pro Pro Ala Gln Leu Gln His Ile Asp				
	35	40		

<210> 6  
 <211> 44  
 <212> PRT  
 <213> Artificial Sequence

1	5	10	15
Leu Gln Leu Gln His Gln His Gln Arg Ala Gln Gln Asp Glu Ala Arg			
Val Arg Pro Val Leu Gln Pro Gly His Arg Ala His Arg Gln Arg Gln			
	20	25	30
His Gln Leu Pro Pro Ala Gln Leu Gln His Ile Asp			
	35	40	

<210> 7  
 <211> 161  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(161)

1	5	10	15	
atc gat cat cac cca gcc ctg ccc caa ggt gag ctt cga gcc cat ccc				48
Ile Asp His His Pro Gly Leu Pro Gln Gly Glu Leu Arg Ala His Pro				
	5	10	15	
cat cca ctt ctg cgc ccc cgc cgg ctt cgc cat cct gaa gtg caa caa				96
His Pro Leu Leu Arg Pro Arg Arg Leu Arg His Pro Glu Val Gln Gln				
	20	25	30	
caa gac ctt caa cgg cac cgg ccc ctg cac caa cgt gag cac cgt gca				144
Gln Asp Leu Gln Arg His Arg Pro Leu His Gln Arg Glu His Arg Ala				
	35	40	45	
gtg cac cca cgg aat tc				161
Val His Pro Arg Asn				
	50			

<210> 8  
 <211> 53  
 <212> PRT  
 <213> Artificial Sequence

1	5	10	15
Ile Asp His His Pro Gly Leu Pro Gln Gly Glu Leu Arg Ala His Pro			
	5	10	15



His Pro Leu Leu Arg Pro Arg Arg Leu Arg His Pro Glu Val Gln Gln  
                   20                  25                  30  
 Gln Asp Leu Gln Arg His Arg Pro Leu His Gln Arg Glu His Arg Ala  
                   35                  40                  45  
 Val His Pro Arg Asn  
                   50

<210> 9  
 <211> 254  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(254)

<400> 9  
 gaa ttc gcc ccg tgg tga gca ccc agc tgc tgc tga acg gca gcc tgg 48  
 Glu Phe Ala Pro Trp \* Ala Pro Ser Cys Cys \* Thr Ala Ala Trp  
   1                  5                  10  
 ccg agg agg agg tgg tga tca gat ctg aga act tca cca aca acg cca 96  
 Pro Arg Arg Arg Trp \* Ser Asp Leu Arg Thr Ser Pro Thr Thr Pro  
   15                  20                  25  
 aga cca tca tgc tgc agc tga acg aga gcg tgg aga tca act gca ccc 144  
 Arg Pro Ser Ser Cys Ser \* Thr Arg Ala Trp Arg Ser Thr Ala Pro  
   30                  35                  40  
 gcc cca aca aca aca ccc gca aga gca tcc aca tgc gcc ctg gcc gcg 192  
 Ala Pro Thr Thr Thr Pro Ala Arg Ala Ser Thr Ser Ala Leu Ala Ala  
   45                  50                  55                  60  
 cct tct aca cca ccg gcg aca tca tgc gcg aca tcc gcc agg ccc act 240  
 Pro Ser Thr Pro Pro Ala Thr Ser Ser Ala Thr Ser Ala Arg Pro Thr  
                   65                  70                  75  
 gca aca tct cta ga 254  
 Ala Thr Ser Leu  
                   80

<210> 10  
 <211> 80  
 <212> PRT  
 <213> Artificial Sequence

<400> 10  
 Glu Phe Ala Pro Trp Ala Pro Ser Cys Cys Thr Ala Ala Trp Pro Arg  
   1                  5                  10                  15  
 Arg Arg Trp Ser Asp Leu Arg Thr Ser Pro Thr Thr Pro Arg Pro Ser  
                   20                  25                  30  
 Ser Cys Ser Thr Arg Ala Trp Arg Ser Thr Ala Pro Ala Pro Thr Thr  
                   35                  40                  45  
 Thr Pro Ala Arg Ala Ser Thr Ser Ala Leu Ala Ala Pro Ser Thr Pro  
   50                  55                  60  
 Pro Ala Thr Ser Ser Ala Thr Ser Ala Arg Pro Thr Ala Thr Ser Leu  
   65                  70                  75                  80

<210> 11  
 <211> 92  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(92)

<400> 11  
 tct aga acc aac tgg acc aac acc ctg aag cgc gtg gcc gag aag ctg 48  
 Ser Arg Thr Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu  
                   5                  10                  15  
 cgc gag aag ttc aac aac acc acc atc gtg ttc aac cag agc tc 92  
 Arg Glu Lys Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser  
                   20                  25                  30

<210> 12  
 <211> 30  
 <212> PRT  
 <213> Artificial Sequence

<400> 12  
 Ser Arg Thr Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu  
   1                  5                  10                  15  
 Arg Glu Lys Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser  
                   20                  25                  30

<210> 13  
 <211> 130  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(130)

<400> 13  
 gag ctc cgg cgg cga ccc cga gat cgt gat gca cag ctt caa ctg cgg 48  
 Glu Leu Arg Arg Arg Pro Arg Asp Arg Asp Ala Gln Leu Gln Leu Arg  
   1                  5                  10                  15  
 cgg cga gtt ctt cta ctg caa cac cac cca gct gtt caa cag cac ctg 96  
 Arg Arg Val Leu Leu Leu Gln His His Pro Ala Val Gln Gln His Leu  
                   20                  25                  30  
 gaa cga gac caa cag cga ggg caa cat cac tag t 130  
 Glu Arg Asp Gln Gln Arg Gly Gln His His \*  
                   35                  40

<210> 14  
 <211> 42  
 <212> PRT  
 <213> Artificial Sequence

<400> 14

Glu Leu Arg Arg Arg Pro Arg Asp Arg Asp Ala Gln Leu Gln Leu Arg  
 1 5 10 15  
 Arg Arg Val Leu Leu Leu Gln His His Pro Ala Val Gln Gln His Leu  
 20 25 30  
 Glu Arg Asp Gln Gln Arg Gly Gln His His  
 35 40

<210> 15  
 <211> 164  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(164)

<400> 15  
 act agt ggc acc atc acc ctg ccc tgc cgc atc aag cag atc atc aac 48  
 Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln Ile Ile Asn  
 1 5 10 15  
 atg tgg cag gag gtg ggc aag gcc atg tac gcc ccc ccc atc ggc ggc 96  
 Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro Ile Gly Gly  
 20 25 30  
 cag atc aag tgc ctg agc aac atc acc ggc ctg ctg ctg acc cgc gac 144  
 Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu Thr Arg Asp  
 35 40 45  
 ggc ggc agc gac aac tcg ag 164  
 Gly Gly Ser Asp Asn Ser  
 50

<210> 16  
 <211> 54  
 <212> PRT  
 <213> Artificial Sequence

<400> 16  
 Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln Ile Ile Asn  
 1 5 10 15  
 Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro Ile Gly Gly  
 20 25 30  
 Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu Thr Arg Asp  
 35 40 45  
 Gly Gly Ser Asp Asn Ser  
 50

<210> 17  
 <211> 200  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(200)

<400> 17

```

ctc gag cag cgg caa gga gat ttt ccg ccc cgg cgg cgg cga cat gcg      48
Leu Glu Gln Arg Gln Gly Asp Phe Pro Pro Arg Arg Arg Arg His Ala
 1           5           10           15

cga caa ctg gcg cag cga gct gta caa gta caa ggt ggt gaa gat cga      96
Arg Gln Leu Ala Gln Arg Ala Val Gln Val Gln Gly Gly Glu Asp Arg
          20           25           30

gcc cct ggg cat cgc ccc cac caa ggc caa gcg ccg cgt ggt gca gcg      144
Ala Pro Gly His Arg Pro His Gln Gly Gln Ala Pro Arg Gly Ala Ala
          35           40           45

cga gaa gcg cgc cgt ggg cat cgg cgc-tat gtt cct cgg ctt cct ggg      192
Arg Glu Ala Arg Arg Gly His Arg Arg Tyr Val Pro Arg Leu Pro Gly
 50           55           60

cgc tgc ag      200
Arg Cys
65

```

<210> 18  
 <211> 66  
 <212> PRT  
 <213> Artificial Sequence

```

<400> 18
Leu Glu Gln Arg Gln Gly Asp Phe Pro Pro Arg Arg Arg Arg His Ala
 1           5           10           15
Arg Gln Leu Ala Gln Arg Ala Val Gln Val Gln Gly Gly Glu Asp Arg
          20           25           30
Ala Pro Gly His Arg Pro His Gln Gly Gln Ala Pro Arg Gly Ala Ala
          35           40           45
Arg Glu Ala Arg Arg Gly His Arg Arg Tyr Val Pro Arg Leu Pro Gly
 50           55           60
Arg Cys
65

```

<210> 19  
 <211> 212  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(212)

```

<400> 19
ctc gag cag cgg caa gga gat ttt ccg ccc cgg cgg cgg cga cat gcg      48
Leu Glu Gln Arg Gln Gly Asp Phe Pro Pro Arg Arg Arg Arg His Ala
 1           5           10           15

cga caa ctg gcg cag cga gct gta caa gta caa ggt ggt gaa gat cga      96
Arg Gln Leu Ala Gln Arg Ala Val Gln Val Gln Gly Gly Glu Asp Arg
          20           25           30

gcc cct ggg cat cgc ccc cac caa ggc caa gcg ccg cgt ggt gca gcg      144
Ala Pro Gly His Arg Pro His Gln Gly Gln Ala Pro Arg Gly Ala Ala
          35           40           45

```

cga gaa gcg cgc cta ggg cat cgg cgc tat gtt cct cgg ctt cct ggg 192  
 Arg Glu Ala Arg Leu Gly His Arg Arg Tyr Val Pro Arg Leu Pro Gly  
           50                          55                          60

cgc tgc agc ccg ggg gat cc 212  
 Arg Cys Ser Pro Gly Asp  
           65                          70

<210> 20  
 <211> 70  
 <212> PRT  
 <213> Artificial Sequence

<400> 20  
 Leu Glu Gln Arg Gln Gly Asp Phe Pro Pro Arg Arg Arg Arg His Ala  
   1                          5                          10                          15  
 Arg Gln Leu Ala Gln Arg Ala Val Gln Val Gln Gly Gly Glu Asp Arg  
           20                          25                          30  
 Ala Pro Gly His Arg Pro His Gln Gly Gln Ala Pro Arg Gly Ala Ala  
           35                          40                          45  
 Arg Glu Ala Arg Leu Gly His Arg Arg Tyr Val Pro Arg Leu Pro Gly  
           50                          55                          60  
 Arg Cys Ser Pro Gly Asp  
           65                          70

<210> 21  
 <211> 200  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(200)

<400> 21  
 ctc gag cag cgg caa gga gat ttt ccg ccc cgg cgg cgg cga cat gcg 48  
 Leu Glu Gln Arg Gln Gly Asp Phe Pro Pro Arg Arg Arg Arg His Ala  
   1                          5                          10                          15

cga caa ctg gcg cag cga gct gta caa gta caa ggt ggt gaa gat cga 96  
 Arg Gln Leu Ala Gln Arg Ala Val Gln Val Gln Gly Gly Glu Asp Arg  
           20                          25                          30

gcc cct ggg cat cgc ccc cac caa ggc caa gcg ccg cgt ggt gca gcg 144  
 Ala Pro Gly His Arg Pro His Gln Gly Gln Ala Pro Arg Gly Ala Ala  
           35                          40                          45

cga gaa gag cgc cgt ggg cat cgg cgc tat gtt cct cgg ctt cct ggg 192  
 Arg Glu Glu Arg Arg Gly His Arg Arg Tyr Val Pro Arg Leu Pro Gly  
           50                          55                          60

cgc tgc ag 200  
 Arg Cys  
           65

<210> 22

<211> 66  
 <212> PRT  
 <213> Artificial Sequence

<400> 22  
 Leu Glu Gln Arg Gln Gly Asp Phe Pro Pro Arg Arg Arg Arg His Ala  
 1 5 10 15  
 Arg Gln Leu Ala Gln Arg Ala Val Gln Val Gln Gly Gly Glu Asp Arg  
 20 25 30  
 Ala Pro Gly His Arg Pro His Gln Gly Gln Ala Pro Arg Gly Ala Ala  
 35 40 45  
 Arg Glu Glu Arg Arg Gly His Arg Arg Tyr Val Pro Arg Leu Pro Gly  
 50 55 60  
 Arg Cys  
 65

<210> 23  
 <211> 178  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(178)

<400> 23  
 ctg cag gca gca cca tgg gcg ccg cca gcc tga ccc tga ccg tgc agg 48  
 Leu Gln Ala Ala Pro Trp Ala Pro Pro Ala \* Pro \* Pro Cys Arg  
 1 5 10  
 ccc gcc agc tgc tga gcg gca tcg tgc agc agc aga aca acc tgc tgc 96  
 Pro Ala Ser Cys \* Ala Ala Ser Cys Ser Ser Arg Thr Thr Cys Cys  
 15 20 25  
 gcg cca tcg agg ccc agc agc acc tgc tcc agc tga ccg tgt ggg gca 144  
 Ala Pro Ser Arg Pro Ser Ser Thr Cys Ser Ser \* Pro Cys Gly Ala  
 30 35 40  
 tca agc agc tcc agg ccc gcg tgc tgg ctc tag a 178  
 Ser Ser Ser Ser Arg Pro Ala Cys Trp Leu \*  
 45 50

<210> 24  
 <211> 54  
 <212> PRT  
 <213> Artificial Sequence

<400> 24  
 Leu Gln Ala Ala Pro Trp Ala Pro Pro Ala Pro Pro Cys Arg Pro Ala  
 1 5 10 15  
 Ser Cys Ala Ala Ser Cys Ser Ser Arg Thr Thr Cys Cys Ala Pro Ser  
 20 25 30  
 Arg Pro Ser Ser Thr Cys Ser Ser Pro Cys Gly Ala Ser Ser Ser Ser  
 35 40 45  
 Arg Pro Ala Cys Trp Leu  
 50

<210> 25

<211> 178  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(178)

<400> 25  
 ctg cag gca gca cca tgg gcg ccg cca gcc tga ccc tga ccg tgc agg 48  
 Leu Gln Ala Ala Pro Trp Ala Pro Pro Ala \* Pro \* Pro Cys Arg  
 1 5 10  
 ccc gcc agc tgc tga gcg gca tcg tgc agc agc aga aca acc tgc tgc 96  
 Pro Ala Ser Cys \* Ala Ala Ser Cys Ser Ser Arg Thr Thr Cys Cys  
 15 20 25  
 gcg cca tcg agg ccc agc agc acc tgc tcc agc tga ccg tgt ggg gca 144  
 Ala Pro Ser Arg Pro Ser Ser Thr Cys Ser Ser \* Pro Cys Gly Ala  
 30 35 40  
 tca agc agt gct gcg gcc gcg tgc tgg ctc tag a 178  
 Ser Ser Ser Ala Ala Ala Ala Cys Trp Leu \*  
 45 50

<210> 26  
 <211> 54  
 <212> PRT  
 <213> Artificial Sequence

<400> 26  
 Leu Gln Ala Ala Pro Trp Ala Pro Pro Ala Pro Pro Cys Arg Pro Ala  
 1 5 10 15  
 Ser Cys Ala Ala Ser Cys Ser Ser Arg Thr Thr Cys Cys Ala Pro Ser  
 20 25 30  
 Arg Pro Ser Ser Thr Cys Ser Ser Pro Cys Gly Ala Ser Ser Ser Ala  
 35 40 45  
 Ala Ala Ala Cys Trp Leu  
 50

<210> 27  
 <211> 77  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(77)

<400> 27  
 tct aga gcg cta cct cca gga cca gcg ctt cct ggg cat gtg ggg ctg 48  
 Ser Arg Ala Leu Pro Pro Gly Pro Ala Leu Pro Gly His Val Gly Leu  
 1 5 10 15  
 ctc cgg caa gct gat ctg cac cac gcc cg 77  
 Leu Arg Gln Ala Asp Leu His His Gly  
 20 25

<210> 28  
 <211> 25  
 <212> PRT  
 <213> Artificial Sequence

<400> 28  
 Ser Arg Ala Leu Pro Pro Gly Pro Ala Leu Pro Gly His Val Gly Leu  
 1 5 10 15  
 Leu Arg Gln Ala Asp Leu His His Gly  
 20 25

<210> 29  
 <211> 190  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(190)

<400> 29  
 cgg ccg tgc cct gga acg cca gct gga gca aca aga acc tga gcc aga 48  
 Arg Pro Cys Pro Gly Thr Pro Ala Gly Ala Thr Arg Thr \* Ala Arg  
 1 5 10 15  
 ttt ggg aca aca tga cct gga tgg agt ggg agc gcg aga tca gca act 96  
 Phe Gly Thr Thr \* Pro Gly Trp Ser Gly Ser Ala Arg Ser Ala Thr  
 20 25 30  
 aca ccg aga tca tct aca gcc tga tcg agg aga gcc aga acc agc agg 144  
 Thr Pro Arg Ser Ser Thr Ala \* Ser Arg Arg Ala Arg Thr Ser Arg  
 35 40 45  
 aga aga acg agc tgg acc tgc tcc agc tgg aca agt ggg caa gct t 190  
 Arg Arg Thr Ser Trp Thr Cys Ser Ser Trp Thr Ser Gly Gln Ala  
 50 55 60

<210> 30  
 <211> 60  
 <212> PRT  
 <213> Artificial Sequence

<400> 30  
 Arg Pro Cys Pro Gly Thr Pro Ala Gly Ala Thr Arg Thr Ala Arg Phe  
 1 5 10 15  
 Gly Thr Thr Pro Gly Trp Ser Gly Ser Ala Arg Ser Ala Thr Pro  
 20 25 30  
 Arg Ser Ser Thr Ala Ser Arg Arg Ala Arg Thr Ser Arg Arg Arg Thr  
 35 40 45  
 Ser Trp Thr Cys Ser Ser Trp Thr Ser Gly Gln Ala  
 50 55 60

<210> 31  
 <211> 177  
 <212> DNA  
 <213> Artificial Sequence



<220>  
 <221> CDS  
 <222> (1)...(177)

<400> 31  
 aag ctt gtg gaa ctg gtt caa cat cac caa ctg gct gtg gta cat caa 48  
 Lys Leu Val Glu Leu Val Gln His His Gln Leu Ala Val Val His Gln  
 1 5 10 15  
 gat ttt cat cat gat cgt ggg cgg cct gat cgg cct gcg cat cgt gtt 96  
 Asp Phe His His Asp Arg Gly Arg Pro Asp Arg Pro Ala His Arg Val  
 20 25 30  
 cac cgt gct gag cat cgt gaa ccg cgt gcg cca ggg cta cag ccc cct 144  
 His Arg Ala Glu His Arg Glu Pro Arg Ala Pro Gly Leu Gln Pro Pro  
 35 40 45  
 gag ctt cca gac ccg cct gcc cgt gcc ccg cgg 177  
 Glu Leu Pro Asp Pro Pro Ala Arg Ala Pro Arg  
 50 55

<210> 32  
 <211> 59  
 <212> PRT  
 <213> Artificial Sequence

<400> 32  
 Lys Leu Val Glu Leu Val Gln His His Gln Leu Ala Val Val His Gln  
 1 5 10 15  
 Asp Phe His His Asp Arg Gly Arg Pro Asp Arg Pro Ala His Arg Val  
 20 25 30  
 His Arg Ala Glu His Arg Glu Pro Arg Ala Pro Gly Leu Gln Pro Pro  
 35 40 45  
 Glu Leu Pro Asp Pro Pro Ala Arg Ala Pro Arg  
 50 55

<210> 33  
 <211> 140  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(140)

<400> 33  
 ccg cgg ccc cga ccg ccc cga ggg cat cga gga gga ggg cgg cga gcg 48  
 Pro Arg Pro Arg Pro Pro Arg Gly His Arg Gly Gly Gly Arg Arg Ala  
 1 5 10 15  
 cga ccg cga ccg cag cac ccg cct ggt gac cgg ctt cct gcc cct gat 96  
 Arg Pro Arg Pro Gln His Pro Pro Gly Asp Arg Leu Pro Ala Pro Asp  
 20 25 30  
 ctg gga cga cct gcg cag cct gtt cct gtt cag cta cca tcg at 140  
 Leu Gly Arg Pro Ala Gln Pro Val Pro Val Gln Leu Pro Ser  
 35 40 45

<210> 34  
 <211> 46  
 <212> PRT  
 <213> Artificial Sequence

<400> 34  
 Pro Arg Pro Arg Pro Pro Arg Gly His Arg Gly Gly Gly Arg Arg Ala  
 1 5 10 15  
 Arg Pro Arg Pro Gln His Pro Pro Gly Asp Arg Leu Pro Ala Pro Asp  
 20 25 30  
 Leu Gly Arg Pro Ala Gln Pro Val Pro Val Gln Leu Pro Ser  
 35 40 45

<210> 35  
 <211> 129  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(129)

<400> 35  
 atc gat tgc gcg acc tgc tgc tga tcg tgg ccc gca tcg tgg agc tgc 48  
 Ile Asp Cys Ala Thr Cys Cys \* Ser Trp Pro Ala Ser Trp Ser Cys  
 1 5 10 15  
 tgg gcc ggc gcg gct ggg aga tcc tga agt act ggt gga acc tgc tcc 96  
 Trp Ala Gly Ala Ala Gly Arg Ser \* Ser Thr Gly Gly Thr Cys Ser  
 20 25 30  
 agt act gga gcc agg agc tga aga act ctg cag 129  
 Ser Thr Gly Ala Arg Ser \* Arg Thr Leu Gln  
 35 40

<210> 36  
 <211> 40  
 <212> PRT  
 <213> Artificial Sequence

<400> 36  
 Ile Asp Cys Ala Thr Cys Cys Ser Trp Pro Ala Ser Trp Ser Cys Trp  
 1 5 10 15  
 Ala Gly Ala Ala Gly Arg Ser Ser Thr Gly Gly Thr Cys Ser Ser Thr  
 20 25 30  
 Gly Ala Arg Ser Arg Thr Leu Gln  
 35 40

<210> 37  
 <211> 114  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(114)

<400> 37  
 ctg cag tga gcc tgc tga acg cca ccg cca tgc ccg tgg ccg agg gca 48  
 Leu Gln \* Ala Cys \* Thr Pro Pro Pro Ser Pro Trp Pro Arg Ala  
 1 5 10  
 ccg acc gcg tga tgc agg tgg tgc agc gca tct ggc gcg gca tcc tgc 96  
 Pro Thr Ala \* Ser Arg Trp Cys Ser Ala Ser Gly Ala Ala Ser Cys  
 15 20 25  
 aca tcc cca ccc gaa ttc 114  
 Thr Ser Pro Pro Glu Phe  
 30 35

<210> 38  
 <211> 35  
 <212> PRT  
 <213> Artificial Sequence

<400> 38  
 Leu Gln Ala Cys Thr Pro Pro Pro Ser Pro Trp Pro Arg Ala Pro Thr  
 1 5 10 15  
 Ala Ser Arg Trp Cys Ser Ala Ser Gly Ala Ala Ser Cys Thr Ser Pro  
 20 25 30  
 Pro Glu Phe  
 35

<210> 39  
 <211> 41  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(41)

<400> 39  
 gaa ttc gcc agg gct tgc agc gcg ccc tgc tgt aag gat cc 41  
 Glu Phe Ala Arg Ala Ser Ser Ala Pro Cys Cys Lys Asp  
 1 5 10

<210> 40  
 <211> 13  
 <212> PRT  
 <213> Artificial Sequence

<400> 40  
 Glu Phe Ala Arg Ala Ser Ser Ala Pro Cys Cys Lys Asp  
 1 5 10

<210> 41  
 <211> 506  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(506)

<400> 41  
gct agc gcg gcc gac cgc ctg tgg gtg acc gtg tac tac ggc gtg ccc 48  
Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
1 5 10 15

gtg tgg aag gac gcc acc acc acc ctg ttc tgc gcc agc gac gcc aag 96  
Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
20 25 30

gcc tac gac acc gag gtg cac aac gtg tgg gcc acc cac gcg tgc gtg 144  
Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
35 40 45

ccc acc gac ccc aac ccc cag gag gtg gtg ctg ggc aac gtg acc gag 192  
Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
50 55 60

aac ttc aac atg ggc aag aac aac atg gtg gag cag atg cac gag gat 240  
Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp  
65 70 75 80

atc atc agc ctg tgg gac cag agc ctg aag ccc tgc gtg aag ctg acc 288  
Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr  
85 90 95

ccc ctg tgc gtg acc ctg aac tgc acc aag ctg aag aac agc acc gac 336  
Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr Asp  
100 105 110

acc aac aac acc cgc tgg ggc acc cag gag atg aag aac tgc agc ttc 384  
Thr Asn Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys Ser Phe  
115 120 125

aac atc agc acc agc gtg cgc aac aag atg aag cgc gag tac gcc ctg 432  
Asn Ile Ser Thr Ser Val Arg Asn Lys Met Lys Arg Glu Tyr Ala Leu  
130 135 140

ttc tac agc ctg gac atc gtg ccc atc gac aac gac aac acc agc tac 480  
Phe Tyr Ser Leu Asp Ile Val Pro Ile Asp Asn Asp Asn Thr Ser Tyr  
145 150 155 160

cgc ctg cgc agc tgc aac aca tcg at 506  
Arg Leu Arg Ser Cys Asn Thr Ser  
165

&lt;210&gt; 42

&lt;211&gt; 168

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

<400> 42  
Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
1 5 10 15  
Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
20 25 30  
Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
35 40 45

Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
 50 55 60  
 Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp  
 65 70 75 80  
 Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr  
 85 90 95  
 Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr Asp  
 100 105 110  
 Thr Asn Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys Ser Phe  
 115 120 125  
 Asn Ile Ser Thr Ser Val Arg Asn Lys Met Lys Arg Glu Tyr Ala Leu  
 130 135 140  
 Phe Tyr Ser Leu Asp Ile Val Pro Ile Asp Asn Asp Asn Thr Ser Tyr  
 145 150 155 160  
 Arg Leu Arg Ser Cys Asn Thr Ser  
 165

<210> 43  
 <211> 374  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(374)

<400> 43  
 tct aga acc aac tgg acc aac acc ctg aag cgc gtg gcc gag aag ctg 48  
 Ser Arg Thr Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu  
 1 5 10 15  
 cgc gag aag ttc aac aac acc acc atc gtg ttc aac cag agc tcc gcc 96  
 Arg Glu Lys Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser Ser Gly  
 20 25 30  
 gcc gac ccc gag atc gtg atg cac agc ttc aac tgc gcc gcc gag ttc 144  
 Gly Asp Pro Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe  
 35 40 45  
 ttc tac tgc aac acc acc cag ctg ttc aac agc acc tgg aac gag acc 192  
 Phe Tyr Cys Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr  
 50 55 60  
 aac agc gag gcc aac atc act agt gcc acc atc acc ctg ccc tgc cgc 240  
 Asn Ser Glu Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg  
 65 70 75 80  
 atc aag cag atc atc aac atg tgg cag gag gtg gcc aag gcc atg tac 288  
 Ile Lys Gln Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr  
 85 90 95  
 gcc ccc ccc atc gcc gcc cag atc aag tgc ctg agc aac atc acc gcc 336  
 Ala Pro Pro Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly  
 100 105 110  
 ctg ctg ctg acc cgc gac gcc gcc agc gac aac tgc ag 374  
 Leu Leu Leu Thr Arg Asp Gly Gly Ser Asp Asn Ser  
 115 120

<210> 44  
 <211> 124  
 <212> PRT  
 <213> Artificial Sequence

<400> 44  
 Ser Arg Thr Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu  
 1 5 10 15  
 Arg Glu Lys Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser Ser Gly  
 20 25 30  
 Gly Asp Pro Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe  
 35 40 45  
 Phe Tyr Cys Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr  
 50 55 60  
 Asn Ser Glu Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg  
 65 70 75 80  
 Ile Lys Gln Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr  
 85 90 95  
 Ala Pro Pro Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly  
 100 105 110  
 Leu Leu Leu Thr Arg Asp Gly Gly Ser Asp Asn Ser  
 115 120

<210> 45  
 <211> 1277  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(1277)

<400> 45  
 gct agc gcg gcc gac cgc ctg tgg gtg acc gtg tac tac ggc gtg ccc 48  
 Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
 1 5 10 15  
 gtg tgg aag gac gcc acc acc acc ctg ttc tgc gcc agc gac gcc aag 96  
 Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
 20 25 30  
 gcc tac gac acc gag gtg cac aac gtg tgg gcc acc cac gcg tgc gtg 144  
 Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
 35 40 45  
 ccc acc gac ccc aac ccc cag gag gtg gtg ctg ggc aac gtg acc gag 192  
 Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
 50 55 60  
 aac ttc aac atg ggc aag aac aac atg gtg gag cag atg cac gag gat 240  
 Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp  
 65 70 75 80  
 atc atc agc ctg tgg gac cag agc ctg aag ccc tgc gtg aag ctg acc 288  
 Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr  
 85 90 95  
 ccc ctg tgc gtg acc ctg aac tgc acc aag ctg aag aac agc acc gac 336

Pro	Leu	Cys	Val	Thr	Leu	Asn	Cys	Thr	Lys	Leu	Lys	Asn	Ser	Thr	Asp	
			100					105					110			
acc	aac	aac	acc	cgc	tgg	ggc	acc	cag	gag	atg	aag	aac	tgc	agc	ttc	384
Thr	Asn	Asn	Thr	Arg	Trp	Gly	Thr	Gln	Glu	Met	Lys	Asn	Cys	Ser	Phe	
		115					120					125				
aac	atc	agc	acc	agc	gtg	cgc	aac	aag	atg	aag	cgc	gag	tac	gcc	ctg	432
Asn	Ile	Ser	Thr	Ser	Val	Arg	Asn	Lys	Met	Lys	Arg	Glu	Tyr	Ala	Leu	
		130				135					140					
ttc	tac	agc	ctg	gac	atc	gtg	ccc	atc	gac	aac	gac	aac	acc	agc	tac	480
Phe	Tyr	Ser	Leu	Asp	Ile	Val	Pro	Ile	Asp	Asn	Asp	Asn	Thr	Ser	Tyr	
145					150					155					160	
cgc	ctg	cgc	agc	tgc	aac	aca	tcg	atc	atc	acc	cag	gcc	tgc	ccc	aag	528
Arg	Leu	Arg	Ser	Cys	Asn	Thr	Ser	Ile	Ile	Thr	Gln	Ala	Cys	Pro	Lys	
				165					170					175		
gtg	agc	ttc	gag	ccc	atc	ccc	atc	cac	ttc	tgc	gcc	ccc	gcc	ggc	ttc	576
Val	Ser	Phe	Glu	Pro	Ile	Pro	Ile	His	Phe	Cys	Ala	Pro	Ala	Gly	Phe	
			180					185					190			
gcc	atc	ctg	aag	tgc	aac	aac	aag	acc	ttc	aac	ggc	acc	ggc	ccc	tgc	624
Ala	Ile	Leu	Lys	Cys	Asn	Asn	Lys	Thr	Phe	Asn	Gly	Thr	Gly	Pro	Cys	
		195					200					205				
acc	aac	gtg	agc	acc	gtg	cag	tgc	acc	cac	gga	att	cgc	ccc	gtg	gtg	672
Thr	Asn	Val	Ser	Thr	Val	Gln	Cys	Thr	His	Gly	Ile	Arg	Pro	Val	Val	
		210				215					220					
agc	acc	cag	ctg	ctg	ctg	aac	ggc	agc	ctg	gcc	gag	gag	gag	gtg	gtg	720
Ser	Thr	Gln	Leu	Leu	Leu	Asn	Gly	Ser	Leu	Ala	Glu	Glu	Glu	Val	Val	
225					230					235					240	
atc	aga	tct	gag	aac	ttc	acc	aac	aac	gcc	aag	acc	atc	atc	gtg	cag	768
Ile	Arg	Ser	Glu	Asn	Phe	Thr	Asn	Asn	Ala	Lys	Thr	Ile	Ile	Val	Gln	
				245					250					255		
ctg	aac	gag	agc	gtg	gag	atc	aac	tgc	acc	cgc	ccc	aac	aac	aac	acc	816
Leu	Asn	Glu	Ser	Val	Glu	Ile	Asn	Cys	Thr	Arg	Pro	Asn	Asn	Asn	Thr	
			260					265				270				
cgc	aag	agc	atc	cac	atc	ggc	cct	ggc	cgc	gcc	ttc	tac	acc	acc	ggc	864
Arg	Lys	Ser	Ile	His	Ile	Gly	Pro	Gly	Arg	Ala	Phe	Tyr	Thr	Thr	Gly	
		275				280						285				
gac	atc	atc	ggc	gac	atc	cgc	cag	gcc	cac	tgc	aac	atc	tct	aga	acc	912
Asp	Ile	Ile	Gly	Asp	Ile	Arg	Gln	Ala	His	Cys	Asn	Ile	Ser	Arg	Thr	
		290				295					300					
aac	tgg	acc	aac	acc	ctg	aag	cgc	gtg	gcc	gag	aag	ctg	cgc	gag	aag	960
Asn	Trp	Thr	Asn	Thr	Leu	Lys	Arg	Val	Ala	Glu	Lys	Leu	Arg	Glu	Lys	
305					310					315					320	
ttc	aac	aac	acc	acc	atc	gtg	ttc	aac	cag	agc	tcc	ggc	ggc	gac	ccc	1008
Phe	Asn	Asn	Thr	Thr	Ile	Val	Phe	Asn	Gln	Ser	Ser	Gly	Gly	Asp	Pro	
				325					330					335		

gag atc gtg atg cac agc ttc aac tgc ggc ggc gag ttc ttc tac tgc 1056  
 Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe Phe Tyr Cys  
                   340                  345                  350  
  
 aac acc acc cag ctg ttc aac agc acc tgg aac gag acc aac agc gag 1104  
 Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr Asn Ser Glu  
                   355                  360                  365  
  
 ggc aac atc act agt ggc acc atc acc ctg ccc tgc cgc atc aag cag 1152  
 Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln  
                   370                  375                  380  
  
 atc atc aac atg tgg cag gag gtg ggc aag gcc atg tac gcc ccc ccc 1200  
 Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro  
                   385                  390                  395                  400  
  
 atc ggc ggc cag atc aag tgc ctg agc aac atc acc ggc ctg ctg ctg 1248  
 Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu  
                   405                  410                  415  
  
 acc cgc gac ggc ggc agc gac aac tcg ag 1277  
 Thr Arg Asp Gly Gly Ser Asp Asn Ser  
                   420                  425

<210> 46  
 <211> 425  
 <212> PRT  
 <213> Artificial Sequence

<400> 46  
 Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
 1                  5                  10                  15  
 Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
                   20                  25                  30  
 Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
                   35                  40                  45  
 Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
                   50                  55                  60  
 Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp  
                   65                  70                  75                  80  
 Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr  
                   85                  90                  95  
 Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr Asp  
                   100                  105                  110  
 Thr Asn Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys Ser Phe  
                   115                  120                  125  
 Asn Ile Ser Thr Ser Val Arg Asn Lys Met Lys Arg Glu Tyr Ala Leu  
                   130                  135                  140  
 Phe Tyr Ser Leu Asp Ile Val Pro Ile Asp Asn Asp Asn Thr Ser Tyr  
                   145                  150                  155                  160  
 Arg Leu Arg Ser Cys Asn Thr Ser Ile Ile Thr Gln Ala Cys Pro Lys  
                   165                  170                  175  
 Val Ser Phe Glu Pro Ile Pro Ile His Phe Cys Ala Pro Ala Gly Phe  
                   180                  185                  190  
 Ala Ile Leu Lys Cys Asn Asn Lys Thr Phe Asn Gly Thr Gly Pro Cys  
                   195                  200                  205  
 Thr Asn Val Ser Thr Val Gln Cys Thr His Gly Ile Arg Pro Val Val  
                   210                  215                  220



Ser Thr Gln Leu Leu Leu Asn Gly Ser Leu Ala Glu Glu Glu Val Val  
 225 230 235 240  
 Ile Arg Ser Glu Asn Phe Thr Asn Asn Ala Lys Thr Ile Ile Val Gln  
 245 250 255  
 Leu Asn Glu Ser Val Glu Ile Asn Cys Thr Arg Pro Asn Asn Asn Thr  
 260 265 270  
 Arg Lys Ser Ile His Ile Gly Pro Gly Arg Ala Phe Tyr Thr Thr Gly  
 275 280 285  
 Asp Ile Ile Gly Asp Ile Arg Gln Ala His Cys Asn Ile Ser Arg Thr  
 290 295 300  
 Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu Arg Glu Lys  
 305 310 315 320  
 Phe Asn Asn Thr Thr Ile Val Phe Asn-Gln Ser Ser Gly Gly Asp Pro  
 325 330 335  
 Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe Phe Tyr Cys  
 340 345 350  
 Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr Asn Ser Glu  
 355 360 365  
 Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln  
 370 375 380  
 Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro  
 385 390 395 400  
 Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu  
 405 410 415  
 Thr Arg Asp Gly Gly Ser Asp Asn Ser  
 420 425

<210> 47  
 <211> 1277  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(1277)

<400> 47

gct agc gcg gcc gac cgc ctg tgg gtg acc gtg tac tac ggc gtg ccc	48
Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro	
1 5 10 15	
gtg tgg aag gac gcc acc acc acc ctg ttc tgc gcc agc gac gcc aag	96
Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys	
20 25 30	
gcc tac gac acc gag gtg cac aac gtg tgg gcc acc cac gcg tgc gtg	144
Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val	
35 40 45	
ccc acc gac ccc aac ccc cag gag gtg gtg ctg ggc aac gtg acc gag	192
Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu	
50 55 60	
aac ttc aac atg ggc aag aac aac atg gtg gag cag atg cac gag gat	240
Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp	
65 70 75 80	
atc atc agc ctg tgg gac cag agc ctg aag ccc tgc gtg aag ctg acc	288
Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr	

85															90					95					
ccc	ctg	tgc	gtg	acc	ctg	caa	tgc	acc	aag	ctg	aag	cag	agc	acc	gac	336									
Pro	Leu	Cys	Val	Thr	Leu	Gln	Cys	Thr	Lys	Leu	Lys	Gln	Ser	Thr	Asp										
			100					105					110												
acc	cag	aac	acc	cgc	tgg	ggc	acc	cag	gag	atg	aag	aac	tgc	agc	ttc	384									
Thr	Gln	Asn	Thr	Arg	Trp	Gly	Thr	Gln	Glu	Met	Lys	Asn	Cys	Ser	Phe										
		115					120					125													
aac	atc	agc	acc	agc	gtg	cgc	aac	aag	atg	aag	cgc	gag	tac	gcc	ctg	432									
Asn	Ile	Ser	Thr	Ser	Val	Arg	Asn	Lys	Met	Lys	Arg	Glu	Tyr	Ala	Leu										
	130					135					140														
ttc	tac	agc	ctg	gac	atc	gtg	ccc	atc	gac	aac	gac	aac	acc	agc	tac	480									
Phe	Tyr	Ser	Leu	Asp	Ile	Val	Pro	Ile	Asp	Asn	Asp	Asn	Thr	Ser	Tyr										
145					150				155						160										
cgc	ctg	cgc	agc	tgc	aac	aca	tcg	atc	atc	acc	cag	gcc	tgc	ccc	aag	528									
Arg	Leu	Arg	Ser	Cys	Asn	Thr	Ser	Ile	Ile	Thr	Gln	Ala	Cys	Pro	Lys										
				165					170					175											
gtg	agc	ttc	gag	ccc	atc	ccc	atc	cac	ttc	tgc	gcc	ccc	gcc	ggc	ttc	576									
Val	Ser	Phe	Glu	Pro	Ile	Pro	Ile	His	Phe	Cys	Ala	Pro	Ala	Gly	Phe										
			180					185					190												
gcc	atc	ctg	aag	tgc	aac	aac	aag	acc	ttc	aac	ggc	acc	ggc	ccc	tgc	624									
Ala	Ile	Leu	Lys	Cys	Asn	Asn	Lys	Thr	Phe	Asn	Gly	Thr	Gly	Pro	Cys										
		195					200					205													
acc	aac	gtg	agc	acc	gtg	cag	tgc	acc	cac	gga	att	cgc	ccc	gtg	gtg	672									
Thr	Asn	Val	Ser	Thr	Val	Gln	Cys	Thr	His	Gly	Ile	Arg	Pro	Val	Val										
	210					215					220														
agc	acc	cag	ctg	ctg	ctg	aac	ggc	agc	ctg	gcc	gag	gag	gag	gtg	gtg	720									
Ser	Thr	Gln	Leu	Leu	Leu	Asn	Gly	Ser	Leu	Ala	Glu	Glu	Glu	Val	Val										
225					230					235				240											
atc	aga	tct	gag	aac	ttc	acc	aac	aac	gcc	aag	acc	atc	atc	gtg	cag	768									
Ile	Arg	Ser	Glu	Asn	Phe	Thr	Asn	Asn	Ala	Lys	Thr	Ile	Ile	Val	Gln										
			245						250					255											
ctg	aac	gag	agc	gtg	gag	atc	aac	tgc	acc	cgc	ccc	aac	aac	aac	acc	816									
Leu	Asn	Glu	Ser	Val	Glu	Ile	Asn	Cys	Thr	Arg	Pro	Asn	Asn	Asn	Thr										
			260					265						270											
cgc	aag	agc	atc	cac	atc	ggc	cct	ggc	cgc	gcc	ttc	tac	acc	acc	ggc	864									
Arg	Lys	Ser	Ile	His	Ile	Gly	Pro	Gly	Arg	Ala	Phe	Tyr	Thr	Thr	Gly										
		275					280					285													
gac	atc	atc	ggc	gac	atc	cgc	cag	gcc	cac	tgc	aac	atc	tct	aga	acc	912									
Asp	Ile	Ile	Gly	Asp	Ile	Arg	Gln	Ala	His	Cys	Asn	Ile	Ser	Arg	Thr										
	290					295					300														
aac	tgg	acc	aac	acc	ctg	aag	cgc	gtg	gcc	gag	aag	ctg	cgc	gag	aag	960									
Asn	Trp	Thr	Asn	Thr	Leu	Lys	Arg	Val	Ala	Glu	Lys	Leu	Arg	Glu	Lys										
305					310				315					320											
ttc	aac	aac	acc	acc	atc	gtg	ttc	aac	cag	agc	tcc	ggc	ggc	gac	ccc	1008									

Phe	Asn	Asn	Thr	Thr	Ile	Val	Phe	Asn	Gln	Ser	Ser	Gly	Gly	Asp	Pro		
				325					330					335			
gag	atc	gtg	atg	cac	agc	ttc	aac	tgc	ggc	ggc	gag	ttc	ttc	tac	tgc		1056
Glu	Ile	Val	Met	His	Ser	Phe	Asn	Cys	Gly	Gly	Glu	Phe	Phe	Tyr	Cys		
			340					345					350				
aac	acc	acc	cag	ctg	ttc	aac	agc	acc	tgg	aac	gag	acc	aac	agc	gag		1104
Asn	Thr	Thr	Gln	Leu	Phe	Asn	Ser	Thr	Trp	Asn	Glu	Thr	Asn	Ser	Glu		
			355				360					365					
ggc	aac	atc	act	agt	ggc	acc	atc	acc	ctg	ccc	tgc	cgc	atc	aag	cag		1152
Gly	Asn	Ile	Thr	Ser	Gly	Thr	Ile	Thr	Leu	Pro	Cys	Arg	Ile	Lys	Gln		
	370					375					380						
atc	atc	aac	atg	tgg	cag	gag	gtg	ggc	aag	gcc	atg	tac	gcc	ccc	ccc		1200
Ile	Ile	Asn	Met	Trp	Gln	Glu	Val	Gly	Lys	Ala	Met	Tyr	Ala	Pro	Pro		
	385				390				395					400			
atc	ggc	ggc	cag	atc	aag	tgc	ctg	agc	aac	atc	acc	ggc	ctg	ctg	ctg		1248
Ile	Gly	Gly	Gln	Ile	Lys	Cys	Leu	Ser	Asn	Ile	Thr	Gly	Leu	Leu	Leu		
			405					410					415				
acc	cgc	gac	ggc	ggc	agc	gac	aac	tcg	ag								1277
Thr	Arg	Asp	Gly	Gly	Ser	Asp	Asn	Ser									
			420				425										

<210> 48  
 <211> 425  
 <212> PRT  
 <213> Artificial Sequence

<400> 48

Ala	Ser	Ala	Ala	Asp	Arg	Leu	Trp	Val	Thr	Val	Tyr	Tyr	Gly	Val	Pro		
1				5					10					15			
Val	Trp	Lys	Asp	Ala	Thr	Thr	Thr	Leu	Phe	Cys	Ala	Ser	Asp	Ala	Lys		
		20						25					30				
Ala	Tyr	Asp	Thr	Glu	Val	His	Asn	Val	Trp	Ala	Thr	His	Ala	Cys	Val		
	35					40					45						
Pro	Thr	Asp	Pro	Asn	Pro	Gln	Glu	Val	Val	Leu	Gly	Asn	Val	Thr	Glu		
	50					55					60						
Asn	Phe	Asn	Met	Gly	Lys	Asn	Asn	Met	Val	Glu	Gln	Met	His	Glu	Asp		
	65				70					75				80			
Ile	Ile	Ser	Leu	Trp	Asp	Gln	Ser	Leu	Lys	Pro	Cys	Val	Lys	Leu	Thr		
			85					90					95				
Pro	Leu	Cys	Val	Thr	Leu	Gln	Cys	Thr	Lys	Leu	Lys	Gln	Ser	Thr	Asp		
		100						105					110				
Thr	Gln	Asn	Thr	Arg	Trp	Gly	Thr	Gln	Glu	Met	Lys	Asn	Cys	Ser	Phe		
	115					120						125					
Asn	Ile	Ser	Thr	Ser	Val	Arg	Asn	Lys	Met	Lys	Arg	Glu	Tyr	Ala	Leu		
	130				135						140						
Phe	Tyr	Ser	Leu	Asp	Ile	Val	Pro	Ile	Asp	Asn	Asp	Asn	Thr	Ser	Tyr		
	145				150				155					160			
Arg	Leu	Arg	Ser	Cys	Asn	Thr	Ser	Ile	Ile	Thr	Gln	Ala	Cys	Pro	Lys		
			165					170					175				
Val	Ser	Phe	Glu	Pro	Ile	Pro	Ile	His	Phe	Cys	Ala	Pro	Ala	Gly	Phe		
			180					185					190				
Ala	Ile	Leu	Lys	Cys	Asn	Asn	Lys	Thr	Phe	Asn	Gly	Thr	Gly	Pro	Cys		

```
<210> 49
<211> 144
<212> PRT
<213> Artificial Sequence
```

<400> 49																	
gct	agc	gcg	gcc	gac	cgc	ctg	tgg	gtg	acc	gtg	tac	tac	ggc	gtg	ccc	48	
gtg	tgg	aag	gac	gcc	acc	acc	acc	ctg	ttc	tgc	gcc	agc	gac	gcc	aag	96	
gcc	tac	gac	acc	gag	gtg	cac	aac	gtg	tgg	gcc	acc	cac	gcg	tgc	gtg	144	
ccc	acc	gac	ccc	aac	ccc	cag	gag	gtg	gtg	ctg	ggc	aac	gtg	acc	gag	192	
aac	ttc	aac	atg	ggc	aag	aac	aac	atg	gtg	gag	cag	atg	cac	gag	gat	240	
atc	atc	agc	ctg	tgg	gac	cag	agc	ctg	aag	ccc	tgc	gtg	aag	ctg	acc	288	
ccc	ctg	tgc	gtg	acc	ctg	caa	tgc	acc	aag	ctg	aag	cag	agc	acc	gac	336	
acc	cag	aac	acc	cgc	tgg	ggc	acc	cag	gag	atg	aag	aac	tgc	agc	ttc	384	
cag	atc	agc	acc	agc	gtg	cgc	aac	aag	atg	aag	cgc	gag	tac	gcc	ctg	432	
ttc	tac	agc	ctg	gac	atc	gtg	ccc	atc	gac	aac	gac	cag	acc	agc	tac	480	
cgc	ctg	cgc	agc	tgc	aac	aca	tgc	atc	atc	acc	cag	gcc	tgc	ccc	aag	528	
gtg	agc	ttc	gag	ccc	atc	ccc	atc	cac	ttc	tgc	gcc	ccc	gcc	ggc	ttc	576	
gcc	atc	ctg	aag	tgc	aac	aac	aag	acc	ttc	aac	ggc	acc	ggc	ccc	tgc	624	
acc	aac	gtg	agc	acc	gtg	cag	tgc	acc	cac	gga	att	cgc	ccc	gtg	gtg	672	
agc	acc	cag	ctg	ctg	ctg	aac	ggc	agc	ctg	gcc	gag	gag	gag	gtg	gtg	720	
atc	aga	tct	gag	aac	ttc	acc	aac	aac	gcc	aag	acc	atc	atc	gtg	cag	768	
ctg	aac	gac	agc	gtg	gag	atc	aac	tgc	acc	cgc	ccc	aac	aac	aac	acc	816	
cgc	aag	agc	atc	cac	atc	ggc	cct	ggc	cgc	gcc	ttc	tac	acc	acc	ggc	864	
gac	atc	atc	ggc	gac	atc	cgc	cag	gcc	cac	tgc	aac	atc	tct	aga	acc	912	
aac	tgg	acc	aac	acc	ctg	aag	cgc	gtg	gcc	gag	aag	ctg	cgc	gag	aag	960	
ttc	aac	aac	acc	acc	atc	gtg	ttc	aac	cag	agc	tcc	ggc	ggc	gac	ccc	1008	
gag	atc	gtg	atg	cac	gtc	ttc	aac	tgc	ggc	ggc	gag	ttc	ttc	tac	tgc	1056	
aac	acc	acc	cag	ctg	ttc	aac	agc	acc	tgg	aac	gag	acc	aac	agc	gag	1104	

```

ggc aac atc act agt ggc acc atc acc ctg ccc tgc cgc atc aag cag 1152
atc atc aac atg tgg cag gag gtg ggc aag gcc atg tac gcc ccc ccc 1200
atc ggc ggc cag atc aag tgc ctg agc aac atc acc ggc ctg ctg ctg 1248
acc cgc gac ggc ggc agc gac aac tcg ag 1277

```

```

<210> 50
<211> 425
<212> PRT
<213> Artificial Sequence

```

```

<400> 50
Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro
1 5 10 15
Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys
20 25 30
Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val
35 40 45
Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu
50 55 60
Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp
65 70 75 80
Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr
85 90 95
Pro Leu Cys Val Thr Leu Gln Cys Thr Lys Leu Lys Gln Ser Thr Asp
100 105 110
Thr Gln Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys Ser Phe
115 120 125
Gln Ile Ser Thr Ser Val Arg Asn Lys Met Lys Arg Glu Tyr Ala Leu
130 135 140
Phe Tyr Ser Leu Asp Ile Val Pro Ile Asp Asn Asp Gln Thr Ser Tyr
145 150 155 160
Arg Leu Arg Ser Cys Asn Thr Ser Ile Ile Thr Gln Ala Cys Pro Lys
165 170 175
Val Ser Phe Glu Pro Ile Pro Ile His Phe Cys Ala Pro Ala Gly Phe
180 185 190
Ala Ile Leu Lys Cys Asn Asn Lys Thr Phe Asn Gly Thr Gly Pro Cys
195 200 205
Thr Asn Val Ser Thr Val Gln Cys Thr His Gly Ile Arg Pro Val Val
210 215 220
Ser Thr Gln Leu Leu Leu Asn Gly Ser Leu Ala Glu Glu Glu Val Val
225 230 235 240
Ile Arg Ser Glu Asn Phe Thr Asn Asn Ala Lys Thr Ile Ile Val Gln
245 250 255
Leu Asn Glu Ser Val Glu Ile Asn Cys Thr Arg Pro Asn Asn Asn Thr
260 265 270
Arg Lys Ser Ile His Ile Gly Pro Gly Arg Ala Phe Tyr Thr Thr Gly
275 280 285
Asp Ile Ile Gly Asp Ile Arg Gln Ala His Cys Asn Ile Ser Arg Thr
290 295 300
Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu Arg Glu Lys
305 310 315 320
Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser Ser Gly Gly Asp Pro
325 330 335
Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe Phe Tyr Cys
340 345 350
Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr Asn Ser Glu
355 360 365
Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln
370 375 380

```

Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro  
 385 390 395 400  
 Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu  
 405 410 415  
 Thr Arg Asp Gly Gly Ser Asp Asn Ser  
 420 425

<210> 51  
 <211> 144  
 <212> PRT  
 <213> Artificial Sequence

<400> 51  
 gct agc gcg gcc gac cgc ctg tgg gtg acc gtg tac tac ggc gtg ccc 48  
 gtg tgg aag gac gcc acc acc acc ctg ttc tgc gcc agc gac gcc aag 96  
 gcc tac gac acc gag gtg cac aac gtg tgg gcc acc cac gcg tgc gtg 144  
 ccc acc gac ccc aac ccc cag gag gtg gtg ctg ggc aac gtg acc gag 192  
 aac ttc aac atg ggc aag aac aac atg gtg gag cag atg cac gag gat 240  
 atc atc agc ctg tgg gac cag agc ctg aag ccc tgc gtg aag ctg acc 288  
 ccc ctg tgc gtg acc ctg aac tgc acc aag ctg aag aac agc acc gac 336  
 acc aac aac acc cgc tgg ggc acc cag gag atg aag aac tgc agc ttc 384  
 cag atc agc acc agc gtg cgc aac aag atg aag cgc gag tac gcc ctg 432  
 ttc tac agc ctg gac atc gtg ccc atc gac aac gac cag acc agc tac 480  
 cgc ctg cgc agc tgc aac aca tgc atc atc acc cag gcc tgc ccc aag 528  
 gtg agc ttc gag ccc atc ccc atc cac ttc tgc gcc ccc gcc ggc ttc 576  
 gcc atc ctg aag tgc aac aac aag acc ttc aac ggc acc ggc ccc tgc 624  
 acc aac gtg agc acc gtg cag tgc acc cac gga att cgc ccc gtg gtg 672  
 agc acc cag ctg ctg ctg aac ggc agc ctg gcc gag gag gag gtg gtg 720  
 atc aga tct gag aac ttc acc aac aac gcc aag acc atc atc gtg cag 768  
 ctg aac gag agc gtg gag atc aac tgc acc cgc ccc aac aac aac acc 816  
 cgc aag agc atc cac atc ggc cct ggc cgc gcc ttc tac acc acc ggc 864  
 gac atc atc ggc gac atc cgc cag gcc cac tgc aac atc tct aga acc 912  
 aac tgg acc aac acc ctg aag cgc gtg gcc gag aag ctg cgc gag aag 960  
 ttc aac aac acc acc atc gtg ttc aac cag agc tcc ggc ggc gac ccc 1008  
 gag atc gtg atg cac agc ttc aac tgc ggc ggc gag ttc ttc tac tgc 1056  
 aac acc acc cag ctg ttc aac agc acc tgg aac gag acc aac agc gag 1104  
 ggc aac atc act agt ggc acc atc acc ctg ccc tgc cgc atc aag cag 1152  
 atc atc aac atg tgg cag gag gtg ggc aag gcc atg tac gcc ccc ccc 1200  
 atc ggc ggc cag atc aag tgc ctg agc aac atc acc ggc ctg ctg ctg 1248  
 acc cgc gac ggc ggc agc gac aac tgc ag 1277

<210> 52  
 <211> 425  
 <212> PRT  
 <213> Artificial Sequence

<400> 52  
 Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
 1 5 10 15  
 Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
 20 25 30  
 Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
 35 40 45  
 Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
 50 55 60  
 Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp  
 65 70 75 80  
 Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr  
 85 90 95

Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr Asp  
 100 105 110  
 Thr Asn Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys Ser Phe  
 115 120 125  
 Gln Ile Ser Thr Ser Val Arg Asn Lys Met Lys Arg Glu Tyr Ala Leu  
 130 135 140  
 Phe Tyr Ser Leu Asp Ile Val Pro Ile Asp Asn Asp Gln Thr Ser Tyr  
 145 150 155 160  
 Arg Leu Arg Ser Cys Asn Thr Ser Ile Ile Thr Gln Ala Cys Pro Lys  
 165 170 175  
 Val Ser Phe Glu Pro Ile Pro Ile His Phe Cys Ala Pro Ala Gly Phe  
 180 185 190  
 Ala Ile Leu Lys Cys Asn Asn Lys Thr Phe Asn Gly Thr Gly Pro Cys  
 195 200 205  
 Thr Asn Val Ser Thr Val Gln Cys Thr His Gly Ile Arg Pro Val Val  
 210 215 220  
 Ser Thr Gln Leu Leu Leu Asn Gly Ser Leu Ala Glu Glu Glu Val Val  
 225 230 235 240  
 Ile Arg Ser Glu Asn Phe Thr Asn Asn Ala Lys Thr Ile Ile Val Gln  
 245 250 255  
 Leu Asn Glu Ser Val Glu Ile Asn Cys Thr Arg Pro Asn Asn Asn Thr  
 260 265 270  
 Arg Lys Ser Ile His Ile Gly Pro Gly Arg Ala Phe Tyr Thr Thr Gly  
 275 280 285  
 Asp Ile Ile Gly Asp Ile Arg Gln Ala His Cys Asn Ile Ser Arg Thr  
 290 295 300  
 Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu Arg Glu Lys  
 305 310 315 320  
 Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser Ser Gly Gly Asp Pro  
 325 330 335  
 Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe Phe Tyr Cys  
 340 345 350  
 Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr Asn Ser Glu  
 355 360 365  
 Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln  
 370 375 380  
 Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro  
 385 390 395 400  
 Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu  
 405 410 415  
 Thr Arg Asp Gly Gly Ser Asp Asn Ser  
 420 425

<210> 53  
 <211> 432  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(432)

<400> 53  
 tct aga gcg cta cct cca gga cca gcg ctt cct ggg cat gtg ggg ctg  
 Ser Arg Ala Leu Pro Pro Gly Pro Ala Leu Pro Gly His Val Gly Leu  
 1 5 10 15

48

ctc cgg caa gct gat ctg cac cac ggc cgt gcc ctg gaa cgc cag ctg  
 Leu Arg Gln Ala Asp Leu His His Gly Arg Ala Leu Glu Arg Gln Leu

96

27

20	25	30	
gag caa caa gaa cct gag cca gat ttg gga caa cat gac ctg gat gga Glu Gln Gln Glu Pro Glu Pro Asp Leu Gly Gln His Asp Leu Asp Gly 35 40 45			144
gtg gga gcg cga gat cag caa cta cac cga gat cat cta cag cct gat Val Gly Ala Arg Asp Gln Gln Leu His Arg Asp His Leu Gln Pro Asp 50 55 60			192
cga gga gag cca gaa cca gca gga gaa gaa cga gct gga cct gct cca Arg Gly Glu Pro Glu Pro Ala Gly Glu Glu Arg Ala Gly Pro Ala Pro 65 70 75 80			240
gct gga caa gtg ggc aag ctt gtg gaa ctg gtt caa cat cac caa ctg Ala Gly Gln Val Gly Lys Leu Val Glu Leu Val Gln His His Gln Leu 85 90 95			288
gct gtg gta cat caa gat ttt cat cat gat cgt ggg cgg cct gat cgg Ala Val Val His Gln Asp Phe His His Asp Arg Gly Arg Pro Asp Arg 100 105 110			336
cct gcg cat cgt gtt cac cgt gct gag cat cgt gaa ccg cgt gcg cca Pro Ala His Arg Val His Arg Ala Glu His Arg Glu Pro Arg Ala Pro 115 120 125			384
ggg cta cag ccc cct gag ctt cca gac ccg cct gcc cgt gcc ccg cgg Gly Leu Gln Pro Pro Glu Leu Pro Asp Pro Pro Ala Arg Ala Pro Arg 130 135 140			432

<210> 54  
 <211> 144  
 <212> PRT  
 <213> Artificial Sequence

<400> 54  
 Ser Arg Ala Leu Pro Pro Gly Pro Ala Leu Pro Gly His Val Gly Leu  
 1 5 10 15  
 Leu Arg Gln Ala Asp Leu His His Gly Arg Ala Leu Glu Arg Gln Leu  
 20 25 30  
 Glu Gln Gln Glu Pro Glu Pro Asp Leu Gly Gln His Asp Leu Asp Gly  
 35 40 45  
 Val Gly Ala Arg Asp Gln Gln Leu His Arg Asp His Leu Gln Pro Asp  
 50 55 60  
 Arg Gly Glu Pro Glu Pro Ala Gly Glu Glu Arg Ala Gly Pro Ala Pro  
 65 70 75 80  
 Ala Gly Gln Val Gly Lys Leu Val Glu Leu Val Gln His His Gln Leu  
 85 90 95  
 Ala Val Val His Gln Asp Phe His His Asp Arg Gly Arg Pro Asp Arg  
 100 105 110  
 Pro Ala His Arg Val His Arg Ala Glu His Arg Glu Pro Arg Ala Pro  
 115 120 125  
 Gly Met Gln Pro Pro Glu Leu Pro Asp Pro Pro Ala Arg Val Thr Asp  
 130 135 140

<210> 55  
 <211> 434  
 <212> DNA



## &lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)...(434)

&lt;400&gt; 55

tct	aga	gcg	cta	cct	cca	gga	cca	gcg	ctt	cct	ggg	cat	gtg	ggg	ctg	48
Ser	Arg	Ala	Leu	Pro	Pro	Gly	Pro	Ala	Leu	Pro	Gly	His	Val	Gly	Leu	
1				5				10					15			
ctc	cgg	caa	gct	gat	ctg	cac	cac	ggc	cgt	gcc	ctg	gaa	cgc	cag	ctg	96
Leu	Arg	Gln	Ala	Asp	Leu	His	His	Gly	Arg	Ala	Leu	Glu	Arg	Gln	Leu	
			20					25				30				
gag	caa	caa	gaa	cct	gag	cca	gat	ttg	gga	caa	cat	gac	ctg	gat	gga	144
Glu	Gln	Gln	Glu	Pro	Glu	Pro	Asp	Leu	Gly	Gln	His	Asp	Leu	Asp	Gly	
			35				40					45				
gtg	gga	gcg	cga	gat	cag	caa	cta	cac	cga	gat	cat	cta	cag	cct	gat	192
Val	Gly	Ala	Arg	Asp	Gln	Gln	Leu	His	Arg	Asp	His	Leu	Gln	Pro	Asp	
	50					55					60					
cga	gga	gag	cca	gaa	cca	gca	gga	gaa	gaa	cga	gct	gga	cct	gct	cca	240
Arg	Gly	Glu	Pro	Glu	Pro	Ala	Gly	Glu	Glu	Arg	Ala	Gly	Pro	Ala	Pro	
65				70				75						80		
gct	gga	caa	gtg	ggc	aag	ctt	gtg	gaa	ctg	gtt	caa	cat	cac	caa	ctg	288
Ala	Gly	Gln	Val	Gly	Lys	Leu	Val	Glu	Val	Gln	His	His	Gln	Leu		
			85					90					95			
gct	gtg	gta	cat	caa	gat	ttt	cat	cat	gat	cgt	ggg	cgg	cct	gat	cgg	336
Ala	Val	Val	His	Gln	Asp	Phe	His	His	Asp	Arg	Gly	Arg	Pro	Asp	Arg	
			100					105					110			
cct	gcg	cat	cgt	gtt	cac	cgt	gct	gag	cat	cgt	gaa	ccg	cgt	gcg	cca	384
Pro	Ala	His	Arg	Val	His	Arg	Ala	Glu	His	Arg	Glu	Pro	Arg	Ala	Pro	
			115				120					125				
ggg	atg	cag	ccc	cct	gag	ctt	cca	gac	ccg	cct	gcc	cgt	gtg	acg	gat	432
Gly	Met	Gln	Pro	Pro	Glu	Leu	Pro	Asp	Pro	Pro	Ala	Arg	Val	Thr	Asp	
	130					135					140					
cc																434

&lt;210&gt; 56

&lt;211&gt; 144

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;400&gt; 56

Ser	Arg	Ala	Leu	Pro	Pro	Gly	Pro	Ala	Leu	Pro	Gly	His	Val	Gly	Leu
1				5				10					15		
Leu	Arg	Gln	Ala	Asp	Leu	His	His	Gly	Arg	Ala	Leu	Glu	Arg	Gln	Leu
			20					25				30			
Glu	Gln	Gln	Glu	Pro	Glu	Pro	Asp	Leu	Gly	Gln	His	Asp	Leu	Asp	Gly
			35				40					45			
Val	Gly	Ala	Arg	Asp	Gln	Gln	Leu	His	Arg	Asp	His	Leu	Gln	Pro	Asp
	50					55					60				

Arg Gly Glu Pro Glu Pro Ala Gly Glu Glu Arg Ala Gly Pro Ala Pro  
 65 70 75 80  
 Ala Gly Gln Val Gly Lys Leu Val Glu Leu Val Gln His His Gln Leu  
 85 90 95  
 Ala Val Val His Gln Asp Phe His His Asp Arg Gly Arg Pro Asp Arg  
 100 105 110  
 Pro Ala His Arg Val His Arg Ala Glu His Arg Glu Pro Arg Ala Pro  
 115 120 125  
 Gly Met Gln Pro Pro Glu Leu Pro Asp Pro Pro Ala Arg Val Thr Asp  
 130 135 140

<210> 57  
 <211> 281  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(281)

<400> 57  
 tct aga gcg cta cct cca gga cca gcg ctt cct ggg cat gtg ggg ctg 48  
 Ser Arg Ala Leu Pro Pro Gly Pro Ala Leu Pro Gly His Val Gly Leu  
 1 5 10 15  
 ctc cgg caa gct gat ctg cac cac ggc cgt gcc ctg gaa cgc cag ctg 96  
 Leu Arg Gln Ala Asp Leu His His Gly Arg Ala Leu Glu Arg Gln Leu  
 20 25 30  
 gag caa caa gaa cct gag cca gat ttg gga caa cat gac ctg gat gga 144  
 Glu Gln Gln Glu Pro Glu Pro Asp Leu Gly Gln His Asp Leu Asp Gly  
 35 40 45  
 gtg gga gcg cga gat cag caa cta cac cga gat cat cta cag cct gat 192  
 Val Gly Ala Arg Asp Gln Gln Leu His Arg Asp His Leu Gln Pro Asp  
 50 55 60  
 cga gga gag cca gaa cca gca gga gaa gaa cga gct gga cct gct cca 240  
 Arg Gly Glu Pro Glu Pro Ala Gly Glu Glu Arg Ala Gly Pro Ala Pro  
 65 70 75 80  
 gct gga caa gtg ggc aag ctt gtg tga ctg att gag gat cc 281  
 Ala Gly Gln Val Gly Lys Leu Val \* Leu Ile Glu Asp  
 85 90

<210> 58  
 <211> 92  
 <212> PRT  
 <213> Artificial Sequence

<400> 58  
 Ser Arg Ala Leu Pro Pro Gly Pro Ala Leu Pro Gly His Val Gly Leu  
 1 5 10 15  
 Leu Arg Gln Ala Asp Leu His His Gly Arg Ala Leu Glu Arg Gln Leu  
 20 25 30  
 Glu Gln Gln Glu Pro Glu Pro Asp Leu Gly Gln His Asp Leu Asp Gly  
 35 40 45  
 Val Gly Ala Arg Asp Gln Gln Leu His Arg Asp His Leu Gln Pro Asp

50                      55                      60  
 Arg Gly Glu Pro Glu Pro Ala Gly Glu Glu Arg Ala Gly Pro Ala Pro  
 65                      70                      75                      80  
 Ala Gly Gln Val Gly Lys Leu Val Leu Ile Glu Asp  
                     85                      90

<210> 59  
 <211> 272  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(272)

<400> 59  
 atc gat tgc gcg acc tgc tgc tga tgc tgg ccc gca tgc tgg agc tgc                      48  
 ile asp cys ala thr cys cys \* ser trp pro ala ser trp ser cys  
   1                      5                      10                      15  
 tgg gcc gcc gcg gct ggg aga tcc tga agt act ggt gga acc tgc tcc                      96  
 trp ala gly ala ala gly arg ser \* ser thr gly gly thr cys ser  
                     20                      25                      30  
 agt act gga gcc agg agc tga aga act ctg cag tga gcc tgc tga acg                      144  
 ser thr gly ala arg ser \* arg thr leu gln \* ala cys \* thr  
                     35                      40  
 cca ccg cca tgc ccg tgg ccg agg gca ccg acc gcg tga tgc agg tgg                      192  
 pro pro pro ser pro trp pro arg ala pro thr ala \* ser arg trp  
   45                      50                      55  
 tgc agc gca tct ggc gcg gca tcc tgc aca tcc cca ccc gaa ttc gcc                      240  
 cys ser ala ser gly ala ala ser cys thr ser pro pro glu phe ala  
   60                      65                      70  
 agg gct tgc agc gcg ccc tgc tgt aag gat cc                      272  
 arg ala ser ser ala pro cys cys lys asp  
   75                      80

<210> 60  
 <211> 84  
 <212> PRT  
 <213> Artificial Sequence

<400> 60  
 ile asp cys ala thr cys cys ser trp pro ala ser trp ser cys trp  
   1                      5                      10                      15  
 ala gly ala ala gly arg ser ser thr gly gly thr cys ser ser thr  
                     20                      25                      30  
 gly ala arg ser arg thr leu gln ala cys thr pro pro pro ser pro  
   35                      40                      45  
 trp pro arg ala pro thr ala ser arg trp cys ser ala ser gly ala  
   50                      55                      60  
 ala ser cys thr ser pro pro glu phe ala arg ala ser ser ala pro  
   65                      70                      75                      80  
 cys cys lys asp

<210> 61  
 <211> 798  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(798)

<400> 61  
 ctc gag cag cgg caa gga gat ttt ccg ccc cgg cgg cgg cga cat gcg 48  
 Leu Glu Gln Arg Gln Gly Asp Phe Pro Pro Arg Arg Arg Arg His Ala  
 1 5 10 15  
 cga caa ctg gcg cag cga gct gta caa gta caa ggt ggt gaa gat cga 96  
 Arg Gln Leu Ala Gln Arg Ala Val Gln Val Gln Gly Gly Glu Asp Arg  
 20 25 30  
 gcc cct ggg cat cgc ccc cac caa ggc caa gcg ccg cgt ggt gca gcg 144  
 Ala Pro Gly His Arg Pro His Gln Gly Gln Ala Pro Arg Gly Ala Ala  
 35 40 45  
 cga gaa gcg cgc cgt ggg cat cgg cgc tat gtt cct cgg ctt cct ggg 192  
 Arg Glu Ala Arg Arg Gly His Arg Arg Tyr Val Pro Arg Leu Pro Gly  
 50 55 60  
 cgc tgc agg cag cac cat ggg cgc cgc cag cct gac cct gac cgt gca 240  
 Arg Cys Arg Gln His His Gly Arg Arg Gln Pro Asp Pro Asp Arg Ala  
 65 70 75 80  
 ggc ccg cca gct gct gag cgg cat cgt gca gca gca gaa caa cct gct 288  
 Gly Pro Pro Ala Ala Glu Arg His Arg Ala Ala Ala Glu Gln Pro Ala  
 85 90 95  
 gcg cgc cat cga ggc cca gca gca cct gct cca gct gac cgt gtg ggg 336  
 Ala Arg His Arg Gly Pro Ala Ala Pro Ala Pro Ala Asp Arg Val Gly  
 100 105 110  
 cat caa gca gct cca ggc ccg cgt gct ggc tct aga gcg cta cct cca 384  
 His Gln Ala Ala Pro Gly Pro Arg Ala Gly Ser Arg Ala Leu Pro Pro  
 115 120 125  
 gga cca gcg ctt cct ggg cat gtg ggg ctg ctc cgg caa gct gat ctg 432  
 Gly Pro Ala Leu Pro Gly His Val Gly Leu Leu Arg Gln Ala Asp Leu  
 130 135 140  
 cac cac ggc cgt gcc ctg gaa cgc cag ctg gag caa caa gaa cct gag 480  
 His His Gly Arg Ala Leu Glu Arg Gln Leu Glu Gln Gln Glu Pro Glu  
 145 150 155 160  
 cca gat ttg gga caa cat gac ctg gat gga gtg gga gcg cga gat cag 528  
 Pro Asp Leu Gly Gln His Asp Leu Asp Gly Val Gly Ala Arg Asp Gln  
 165 170 175  
 caa cta cac cga gat cat cta cag cct gat cga gga gag cca gaa cca 576  
 Gln Leu His Arg Asp His Leu Gln Pro Asp Arg Gly Glu Pro Glu Pro  
 180 185 190

gca gga gaa gaa cga gct gga cct gct cca gct gga caa gtg ggc aag 624  
 Ala Gly Glu Glu Arg Ala Gly Pro Ala Pro Ala Gly Gln Val Gly Lys  
 195 200 205  
 ctt gtg gaa ctg gtt caa cat cac caa ctg gct gtg gta cat caa gat 672  
 Leu Val Glu Leu Val Gln His His Gln Leu Ala Val Val His Gln Asp  
 210 215 220  
 ttt cat cat gat cgt ggg cgg cct gat cgg cct gcg cat cgt gtt cac 720  
 Phe His His Asp Arg Gly Arg Pro Asp Arg Pro Ala His Arg Val His  
 225 230 235 240  
 cgt gct gag cat cgt gaa ccg cgt gcg cca ggg cta cag ccc cct gag 768  
 Arg Ala Glu His Arg Glu Pro Arg Ala Pro Gly Leu Gln Pro Pro Glu  
 245 250 255  
 ctt cca gac ccg cct gcc cgt gcc ccg cgg 798  
 Leu Pro Asp Pro Pro Ala Arg Ala Pro Arg  
 260 265

<210> 62  
 <211> 266  
 <212> PRT  
 <213> Artificial Sequence

<400> 62  
 Leu Glu Gln Arg Gln Gly Asp Phe Pro Pro Arg Arg Arg Arg His Ala  
 1 5 10 15  
 Arg Gln Leu Ala Gln Arg Ala Val Gln Val Gln Gly Gly Glu Asp Arg  
 20 25 30  
 Ala Pro Gly His Arg Pro His Gln Gly Gln Ala Pro Arg Gly Ala Ala  
 35 40 45  
 Arg Glu Ala Arg Arg Gly His Arg Arg Tyr Val Pro Arg Leu Pro Gly  
 50 55 60  
 Arg Cys Arg Gln His His Gly Arg Arg Gln Pro Asp Pro Asp Arg Ala  
 65 70 75 80  
 Gly Pro Pro Ala Ala Glu Arg His Arg Ala Ala Glu Gln Pro Ala  
 85 90 95  
 Ala Arg His Arg Gly Pro Ala Ala Pro Ala Pro Ala Asp Arg Val Gly  
 100 105 110  
 His Gln Ala Ala Pro Gly Pro Arg Ala Gly Ser Arg Ala Leu Pro Pro  
 115 120 125  
 Gly Pro Ala Leu Pro Gly His Val Gly Leu Leu Arg Gln Ala Asp Leu  
 130 135 140  
 His His Gly Arg Ala Leu Glu Arg Gln Leu Glu Gln Gln Glu Pro Glu  
 145 150 155 160  
 Pro Asp Leu Gly Gln His Asp Leu Asp Gly Val Gly Ala Arg Asp Gln  
 165 170 175  
 Gln Leu His Arg Asp His Leu Gln Pro Asp Arg Gly Glu Pro Glu Pro  
 180 185 190  
 Ala Gly Glu Glu Arg Ala Gly Pro Ala Pro Ala Gly Gln Val Gly Lys  
 195 200 205  
 Leu Val Glu Leu Val Gln His His Gln Leu Ala Val Val His Gln Asp  
 210 215 220  
 Phe His His Asp Arg Gly Arg Pro Asp Arg Pro Ala His Arg Val His  
 225 230 235 240  
 Arg Ala Glu His Arg Glu Pro Arg Ala Pro Gly Leu Gln Pro Pro Glu  
 245 250 255

Leu Pro Asp Pro Pro Ala Arg Ala Pro Arg  
260 265

<210> 63  
<211> 800  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> CDS  
<222> (1)...(800)

<400> 63  
ctc gag cag cgg caa gga gat ttt ccg ccc cgg cgg cgg cga cat gcg 48  
Leu Glu Gln Arg Gln Gly Asp Phe Pro Pro Arg Arg Arg Arg His Ala  
1 5 10 15  
cga caa ctg gcg cag cga gct gta caa gta caa ggt ggt gaa gat cga 96  
Arg Gln Leu Ala Gln Arg Ala Val Gln Val Gln Gly Gly Glu Asp Arg  
20 25 30  
gcc cct ggg cat cgc ccc cac caa ggc caa gcg ccg cgt ggt gca gcg 144  
Ala Pro Gly His Arg Pro His Gln Gly Gln Ala Pro Arg Gly Ala Ala  
35 40 45  
cga gaa gcg cgc cgt ggg cat ccg cgc tat gtt cct ccg ctt cct ggg 192  
Arg Glu Ala Arg Arg Gly His Arg Arg Tyr Val Pro Arg Leu Pro Gly  
50 55 60  
cgc tgc agg cag cac cat ggg cgc cgc cag cct gac cct gac cgt gca 240  
Arg Cys Arg Gln His His Gly Arg Arg Gln Pro Asp Pro Asp Arg Ala  
65 70 75 80  
ggc ccg cca gct gct gag cgg cat cgt gca gca gca gaa caa cct gct 288  
Gly Pro Pro Ala Ala Glu Arg His Arg Ala Ala Ala Glu Gln Pro Ala  
85 90 95  
gcg cgc cat cga ggc cca gca gca cct gct cca gct gac cgt gtg ggg 336  
Ala Arg His Arg Gly Pro Ala Ala Pro Ala Pro Ala Asp Arg Val Gly  
100 105 110  
cat caa gca gct cca ggc ccg cgt gct ggc tct aga gcg cta cct cca 384  
His Gln Ala Ala Pro Gly Pro Arg Ala Gly Ser Arg Ala Leu Pro Pro  
115 120 125  
gga cca gcg ctt cct ggg cat gtg ggg ctg ctg ccg caa gct gat ctg 432  
Gly Pro Ala Leu Pro Gly His Val Gly Leu Leu Arg Gln Ala Asp Leu  
130 135 140  
cac cac ggc cgt gcc ctg gaa cgc cag ctg gag caa caa gaa cct gag 480  
His His Gly Arg Ala Leu Glu Arg Gln Leu Glu Gln Gln Glu Pro Glu  
145 150 155 160  
cca gat ttg gga caa cat gac ctg gat gga gtg gga gcg cga gat cag 528  
Pro Asp Leu Gly Gln His Asp Leu Asp Gly Val Gly Ala Arg Asp Gln  
165 170 175  
caa cta cac cga gat cat cta cag cct gat cga gga gag cca gaa cca 576  
Gln Leu His Arg Asp His Leu Gln Pro Asp Arg Gly Glu Pro Glu Pro

180	185	190	
gca gga gaa gaa cga gct gga cct gct cca gct gga caa gtg ggc aag			624
Ala Gly Glu Glu Arg Ala Gly Pro Ala Pro Ala Gly Gln Val Gly Lys			
195	200	205	
ctt gtg gaa ctg gtt caa cat cac caa ctg gct gtg gta cat caa gat			672
Leu Val Glu Leu Val Gln His His Gln Leu Ala Val Val His Gln Asp			
210	215	220	
ttt cat cat gat cgt ggg cgg cct gat cgg cct gcg cat cgt gtt cac			720
Phe His His Asp Arg Gly Arg Pro Asp Arg Pro Ala His Arg Val His			
225	230	235	240
cgt gct gag cat cgt gaa ccg cgt gcg cca ggg atg cag ccc cct gag			768
Arg Ala Glu His Arg Glu Pro Arg Ala Pro Gly Met Gln Pro Pro Glu			
245	250	255	
ctt cca gac ccg cct gcc cgt gtg acg gat cc			800
Leu Pro Asp Pro Pro Ala Arg Val Thr Asp			
260	265		

<210> 64  
 <211> 266  
 <212> PRT  
 <213> Artificial Sequence

<400> 64

Leu	Glu	Gln	Arg	Gln	Gly	Asp	Phe	Pro	Pro	Arg	Arg	Arg	Arg	His	Ala
1				5					10					15	
Arg	Gln	Leu	Ala	Gln	Arg	Ala	Val	Gln	Val	Gln	Gly	Gly	Glu	Asp	Arg
		20						25					30		
Ala	Pro	Gly	His	Arg	Pro	His	Gln	Gly	Gln	Ala	Pro	Arg	Gly	Ala	Ala
		35				40						45			
Arg	Glu	Ala	Arg	Arg	Gly	His	Arg	Arg	Tyr	Val	Pro	Arg	Leu	Pro	Gly
	50					55				60					
Arg	Cys	Arg	Gln	His	His	Gly	Arg	Arg	Gln	Pro	Asp	Pro	Asp	Arg	Ala
65				70						75				80	
Gly	Pro	Pro	Ala	Ala	Glu	Arg	His	Arg	Ala	Ala	Ala	Glu	Gln	Pro	Ala
			85						90				95		
Ala	Arg	His	Arg	Gly	Pro	Ala	Ala	Pro	Ala	Pro	Ala	Asp	Arg	Val	Gly
			100					105					110		
His	Gln	Ala	Ala	Pro	Gly	Pro	Arg	Ala	Gly	Ser	Arg	Ala	Leu	Pro	Pro
		115					120					125			
Gly	Pro	Ala	Leu	Pro	Gly	His	Val	Gly	Leu	Leu	Arg	Gln	Ala	Asp	Leu
	130					135					140				
His	His	Gly	Arg	Ala	Leu	Glu	Arg	Gln	Leu	Glu	Gln	Gln	Glu	Pro	Glu
145				150						155				160	
Pro	Asp	Leu	Gly	Gln	His	Asp	Leu	Asp	Gly	Val	Gly	Ala	Arg	Asp	Gln
			165						170					175	
Gln	Leu	His	Arg	Asp	His	Leu	Gln	Pro	Asp	Arg	Gly	Glu	Pro	Glu	Pro
			180					185					190		
Ala	Gly	Glu	Arg	Ala	Gly	Pro	Ala	Pro	Ala	Gly	Gln	Val	Gly	Lys	
		195				200					205				
Leu	Val	Glu	Leu	Val	Gln	His	His	Gln	Leu	Ala	Val	Val	His	Gln	Asp
	210					215					220				
Phe	His	His	Asp	Arg	Gly	Arg	Pro	Asp	Arg	Pro	Ala	His	Arg	Val	His
225					230					235					240

Arg Ala Glu His Arg Glu Pro Arg Ala Pro Gly Met Gln Pro Pro Glu  
245 255  
Leu Pro Asp Pro Pro Ala Arg Val Thr Asp  
260 265

```
<210> 65
<211> 647
<212> DNA
<213> Artificial Sequence
```

```
<220>  
<221> CDS  
<222> (1)...(647)
```

<400> 65																	48
ctc	gag	cag	cgg	caa	gga	gat	ttt	ccg	ccc	cgg	cgg	cgg	cga	cat	gcg		
Leu	Glu	Gln	Arg	Gln	Gly	Asp	Phe	Pro	Pro	Arg	Arg	Arg	Arg	His	Ala		
1				5					10					15			
cga caa ctg gcg cag cga gct gta caa gta caa ggt ggt gaa gat cga																	96
Arg	Gln	Leu	Ala	Gln	Arg	Ala	Val	Gln	Val	Gln	Gly	Gly	Glu	Asp	Arg		
			20					25					30				
gcc cct ggg cat cgc ccc cac caa ggc caa gcg ccg cgt ggt gca gcg																	144
Ala	Pro	Gly	His	Arg	Pro	His	Gln	Gly	Gln	Ala	Pro	Arg	Gly	Ala	Ala		
		35					40					45					
cga gaa gcg cgc cgt ggg cat cgg cgc tat gtt cct cgg ctt cct ggg																	192
Arg	Glu	Ala	Arg	Arg	Gly	His	Arg	Arg	Tyr	Val	Pro	Arg	Leu	Pro	Gly		
	50					55					60						
cgc tgc agg cag cac cat ggg cgc cgc cag cct gac cct gac cgt gca																	240
Arg	Cys	Arg	Gln	His	His	Gly	Arg	Arg	Gln	Pro	Asp	Pro	Asp	Arg	Ala		
65					70					75					80		
ggc ccg cca gct gct gag cgg cat cgt gca gca gca gaa caa cct gct																	288
Gly	Pro	Pro	Ala	Ala	Glu	Arg	His	Arg	Ala	Ala	Ala	Glu	Gln	Pro	Ala		
				85					90					95			
gcg cgc cat cga ggc cca gca gca cct gct cca gct gac cgt gtg ggg																	336
Ala	Arg	His	Arg	Gly	Pro	Ala	Ala	Pro	Ala	Pro	Ala	Asp	Arg	Val	Gly		
			100					105					110				
cat caa gca gct cca ggc ccg cgt gct ggc tct aga gcg cta cct cca																	384
His	Gln	Ala	Ala	Pro	Gly	Pro	Arg	Ala	Gly	Ser	Arg	Ala	Leu	Pro	Pro		
		115					120					125					
gga cca gcg ctt cct ggg cat gtg ggg ctg ctc ccg caa gct gat ctg																	432
Gly	Pro	Ala	Leu	Pro	Gly	His	Val	Gly	Leu	Leu	Arg	Gln	Ala	Asp	Leu		
	130					135					140						
cac cac ggc cgt gcc ctg gaa cgc cag ctg gag caa caa gaa cct gag																	480
His	His	Gly	Arg	Ala	Leu	Glu	Arg	Gln	Leu	Glu	Gln	Gln	Glu	Pro	Glu		
145					150					155					160		
cca gat ttg gga caa cat gac ctg gat gga gtg gga gcg cga gat cag																	528
Pro	Asp	Leu	Gly	Gln	His	Asp	Leu	Asp	Gly	Val	Gly	Ala	Arg	Asp	Gln		
				165					170					175			



caa cta cac cga gat cat cta cag cct gat cga gga gag cca gaa cca 576  
 Gln Leu His Arg Asp His Leu Gln Pro Asp Arg Gly Glu Pro Glu Pro  
 180 185 190  
 gca gga gaa gaa cga gct gga cct gct cca gct gga caa gtg ggc aag 624  
 Ala Gly Glu Glu Arg Ala Gly Pro Ala Pro Ala Gly Gln Val Gly Lys  
 195 200 205  
 ctt gtg tga ctg att gag gat cc 647  
 Leu Val \* Leu Ile Glu Asp  
 210

<210> 66  
 <211> 214  
 <212> PRT  
 <213> Artificial Sequence

<400> 66  
 Leu Glu Gln Arg Gln Gly Asp Phe Pro Pro Arg Arg Arg Arg His Ala  
 1 5 10 15  
 Arg Gln Leu Ala Gln Arg Ala Val Gln Val Gln Gly Gly Glu Asp Arg  
 20 25 30  
 Ala Pro Gly His Arg Pro His Gln Gly Gln Ala Pro Arg Gly Ala Ala  
 35 40 45  
 Arg Glu Ala Arg Arg Gly His Arg Arg Tyr Val Pro Arg Leu Pro Gly  
 50 55 60  
 Arg Cys Arg Gln His His Gly Arg Arg Gln Pro Asp Pro Asp Arg Ala  
 65 70 75 80  
 Gly Pro Pro Ala Ala Glu Arg His Arg Ala Ala Ala Glu Gln Pro Ala  
 85 90 95  
 Ala Arg His Arg Gly Pro Ala Ala Pro Ala Pro Ala Asp Arg Val Gly  
 100 105 110  
 His Gln Ala Ala Pro Gly Pro Arg Ala Gly Ser Arg Ala Leu Pro Pro  
 115 120 125  
 Gly Pro Ala Leu Pro Gly His Val Gly Leu Leu Arg Gln Ala Asp Leu  
 130 135 140  
 His His Gly Arg Ala Leu Glu Arg Gln Leu Glu Gln Gln Glu Pro Glu  
 145 150 155 160  
 Pro Asp Leu Gly Gln His Asp Leu Asp Gly Val Gly Ala Arg Asp Gln  
 165 170 175  
 Gln Leu His Arg Asp His Leu Gln Pro Asp Arg Gly Glu Pro Glu Pro  
 180 185 190  
 Ala Gly Glu Glu Arg Ala Gly Pro Ala Pro Ala Gly Gln Val Gly Lys  
 195 200 205  
 Leu Val Leu Ile Glu Asp  
 210

<210> 67  
 <211> 1918  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(1918)

<400> 67  
 gct agc gcg gcc gac cgc ctg tgg gtg acc gtg tac tac ggc gtg ccc 48

Ala	Ser	Ala	Ala	Asp	Arg	Leu	Trp	Val	Thr	Val	Tyr	Tyr	Gly	Val	Pro	
1				5					10					15		
gtg	tgg	aag	gac	gcc	acc	acc	acc	ctg	ttc	tgc	gcc	agc	gac	gcc	aag	96
Val	Trp	Lys	Asp	Ala	Thr	Thr	Thr	Leu	Phe	Cys	Ala	Ser	Asp	Ala	Lys	
			20					25					30			
gcc	tac	gac	acc	gag	gtg	cac	aac	gtg	tgg	gcc	acc	cac	gcg	tgc	gtg	144
Ala	Tyr	Asp	Thr	Glu	Val	His	Asn	Val	Trp	Ala	Thr	His	Ala	Cys	Val	
			35				40					45				
ccc	acc	gac	ccc	aac	ccc	cag	gag	gtg	gtg	ctg	ggc	aac	gtg	acc	gag	192
Pro	Thr	Asp	Pro	Asn	Pro	Gln	Glu	Val	Val	Leu	Gly	Asn	Val	Thr	Glu	
			50			55					60					
aac	ttc	aac	atg	ggc	aag	aac	aac	atg	gtg	gag	cag	atg	cac	gag	gat	240
Asn	Phe	Asn	Met	Gly	Lys	Asn	Asn	Met	Val	Glu	Gln	Met	His	Glu	Asp	
65					70				75						80	
atc	atc	agc	ctg	tgg	gac	cag	agc	ctg	aag	ccc	tgc	gtg	aag	ctg	acc	288
Ile	Ile	Ser	Leu	Trp	Asp	Gln	Ser	Leu	Lys	Pro	Cys	Val	Lys	Leu	Thr	
				85				90						95		
ccc	ctg	tgc	gtg	acc	ctg	aac	tgc	acc	aag	ctg	aag	aac	agc	acc	gac	336
Pro	Leu	Cys	Val	Thr	Leu	Asn	Cys	Thr	Lys	Leu	Lys	Asn	Ser	Thr	Asp	
			100					105					110			
acc	aac	aac	acc	cgc	tgg	ggc	acc	cag	gag	atg	aag	aac	tgc	agc	ttc	384
Thr	Asn	Asn	Thr	Arg	Trp	Gly	Thr	Gln	Glu	Met	Lys	Asn	Cys	Ser	Phe	
			115				120						125			
aac	atc	agc	acc	agc	gtg	cgc	aac	aag	atg	aag	cgc	gag	tac	gcc	ctg	432
Asn	Ile	Ser	Thr	Ser	Val	Arg	Asn	Lys	Met	Lys	Arg	Glu	Tyr	Ala	Leu	
	130					135					140					
ttc	tac	agc	ctg	gac	atc	gtg	ccc	atc	gac	aac	gac	aac	acc	agc	tac	480
Phe	Tyr	Ser	Leu	Asp	Ile	Val	Pro	Ile	Asp	Asn	Asp	Asn	Thr	Ser	Tyr	
145					150					155					160	
cgc	ctg	cgc	agc	tgc	aac	aca	tcg	atc	atc	acc	cag	gcc	tgc	ccc	aag	528
Arg	Leu	Arg	Ser	Cys	Asn	Thr	Ser	Ile	Ile	Thr	Gln	Ala	Cys	Pro	Lys	
				165				170						175		
gtg	agc	ttc	gag	ccc	atc	ccc	atc	cac	ttc	tgc	gcc	ccc	gcc	ggc	ttc	576
Val	Ser	Phe	Glu	Pro	Ile	Pro	Ile	His	Phe	Cys	Ala	Pro	Ala	Gly	Phe	
			180					185					190			
gcc	atc	ctg	aag	tgc	aac	aac	aag	acc	ttc	aac	ggc	acc	ggc	ccc	tgc	624
Ala	Ile	Leu	Lys	Cys	Asn	Asn	Lys	Thr	Phe	Asn	Gly	Thr	Gly	Pro	Cys	
			195				200					205				
acc	aac	gtg	agc	acc	gtg	cag	tgc	acc	cac	gga	att	cgc	ccc	gtg	gtg	672
Thr	Asn	Val	Ser	Thr	Val	Gln	Cys	Thr	His	Gly	Ile	Arg	Pro	Val	Val	
			210			215					220					
agc	acc	cag	ctg	ctg	ctg	aac	ggc	agc	ctg	gcc	gag	gag	gag	gtg	gtg	720
Ser	Thr	Gln	Leu	Leu	Leu	Asn	Gly	Ser	Leu	Ala	Glu	Glu	Glu	Val	Val	
225					230					235					240	

atc aga tct gag aac ttc acc aac aac gcc aag acc atc atc gtg cag Ile Arg Ser Glu Asn Phe Thr Asn Asn Ala Lys Thr Ile Ile Val Gln 245 250 255	768
ctg aac gag agc gtg gag atc aac tgc acc cgc ccc aac aac aac acc Leu Asn Glu Ser Val Glu Ile Asn Cys Thr Arg Pro Asn Asn Asn Thr 260 265 270	816
cgc aag agc atc cac atc ggc cct ggc cgc gcc ttc tac acc acc ggc Arg Lys Ser Ile His Ile Gly Pro Gly Arg Ala Phe Tyr Thr Thr Gly 275 280 285	864
gac atc atc ggc gac atc cgc cag gcc cat tgc aac atc tct aga acc Asp Ile Ile Gly Asp Ile Arg Gln Ala His Cys Asn Ile Ser Arg Thr 290 295 300	912
aac tgg acc aac acc ctg aag cgc gtg gcc gag aag ctg cgc gag aag Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu Arg Glu Lys 305 310 315 320	960
ttc aac aac acc acc atc gtg ttc aac cag agc tcc ggc ggc gac ccc Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser Ser Gly Gly Asp Pro 325 330 335	1008
gag atc gtg atg cac agc ttc aac tgc ggc ggc gag ttc ttc tac tgc Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe Phe Tyr Cys 340 345 350	1056
aac acc acc cag ctg ttc aac agc acc tgg aac gag acc aac agc gag Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr Asn Ser Glu 355 360 365	1104
ggc aac atc act agt ggc acc atc acc ctg ccc tgc cgc atc aag cag Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln 370 375 380	1152
atc atc aac atg tgg cag gag gtg ggc aag gcc atg tac gcc ccc ccc Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro 385 390 395 400	1200
atc ggc ggc cag atc aag tgc ctg agc aac atc acc ggc ctg ctg ctg Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu 405 410 415	1248
acc cgc gac ggc ggc agc gac aac tcg agc agc ggc aag gag att ttc Thr Arg Asp Gly Gly Ser Asp Asn Ser Ser Ser Gly Lys Glu Ile Phe 420 425 430	1296
cgc ccc ggc ggc ggc gac atg cgc gac aac tgg cgc agc gag ctg tac Arg Pro Gly Gly Gly Asp Met Arg Asp Asn Trp Arg Ser Glu Leu Tyr 435 440 445	1344
aag tac aag gtg gtg aag atc gag ccc ctg ggc atc gcc ccc acc aag Lys Tyr Lys Val Val Lys Ile Glu Pro Leu Gly Ile Ala Pro Thr Lys 450 455 460	1392
gcc aag cgc cgc gtg gtg cag cgc gag aag cgc gcc gtg ggc atc ggc Ala Lys Arg Arg Val Val Gln Arg Glu Lys Arg Ala Val Gly Ile Gly 465 470 475 480	1440

gct atg ttc ctc ggc ttc ctg ggc gct gca ggc agc acc atg ggc gcc 1488  
 Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly Ser Thr Met Gly Ala  
 485 490 495  
 gcc agc ctg acc ctg acc gtg cag gcc cgc cag ctg ctg agc ggc atc 1536  
 Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln Leu Leu Ser Gly Ile  
 500 505 510  
 gtg cag cag cag aac aac ctg ctg cgc gcc atc gag gcc cag cag cac 1584  
 Val Gln Gln Gln Asn Asn Leu Leu Arg Ala Ile Glu Ala Gln Gln His  
 515 520 525  
 ctg ctc cag ctg acc gtg tgg ggc atc aag cag ctc cag gcc cgc gtg 1632  
 Leu Leu Gln Leu Thr Val Trp Gly Ile Lys Gln Leu Gln Ala Arg Val  
 530 535 540  
 ctg gct cta gag cgc tac ctc cag gac cag cgc ttc ctg ggc atg tgg 1680  
 Leu Ala Leu Glu Arg Tyr Leu Gln Asp Gln Arg Phe Leu Gly Met Trp  
 545 550 555 560  
 ggc tgc tcc ggc aag ctg atc tgc acc acg gcc gtg ccc tgg aac gcc 1728  
 Gly Cys Ser Gly Lys Leu Ile Cys Thr Thr Ala Val Pro Trp Asn Ala  
 565 570 575  
 agc tgg agc aac aag aac ctg agc cag att tgg gac aac atg acc tgg 1776  
 Ser Trp Ser Asn Lys Asn Leu Ser Gln Ile Trp Asp Asn Met Thr Trp  
 580 585 590  
 atg gag tgg gag cgc gag atc agc aac tac acc gag atc atc tac agc 1824  
 Met Glu Trp Glu Arg Glu Ile Ser Asn Tyr Thr Glu Ile Ile Tyr Ser  
 595 600 605  
 ctg atc gag gag agc cag aac cag cag gag aag aac gag ctg gac ctg 1872  
 Leu Ile Glu Glu Ser Gln Asn Gln Gln Glu Lys Asn Glu Leu Asp Leu  
 610 615 620  
 ctc cag ctg gac aag tgg gca agc ttg tgt gac tga ttg agg atc c 1918  
 Leu Gln Leu Asp Lys Trp Ala Ser Leu Cys Asp \* Leu Arg Ile  
 625 630 635

<210> 68  
 <211> 638  
 <212> PRT  
 <213> Artificial Sequence

<400> 68  
 Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
 1 5 10 15  
 Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
 20 25 30  
 Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
 35 40 45  
 Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
 50 55 60  
 Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp  
 65 70 75 80  
 Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr

				85					90					95			
Pro	Leu	Cys	Val	Thr	Leu	Asn	Cys	Thr	Lys	Leu	Lys	Asn	Ser	Thr	Asp		
			100						105					110			
Thr	Asn	Asn	Thr	Arg	Trp	Gly	Thr	Gln	Glu	Met	Lys	Asn	Cys	Ser	Phe		
		115					120						125				
Asn	Ile	Ser	Thr	Ser	Val	Arg	Asn	Lys	Met	Lys	Arg	Glu	Tyr	Ala	Leu		
	130					135					140						
Phe	Tyr	Ser	Leu	Asp	Ile	Val	Pro	Ile	Asp	Asn	Asp	Asn	Thr	Ser	Tyr		
145				150					155					160			
Arg	Leu	Arg	Ser	Cys	Asn	Thr	Ser	Ile	Ile	Thr	Gln	Ala	Cys	Pro	Lys		
				165					170					175			
Val	Ser	Phe	Glu	Pro	Ile	Pro	Ile	His	Phe	Cys	Ala	Pro	Ala	Gly	Phe		
			180					185					190				
Ala	Ile	Leu	Lys	Cys	Asn	Asn	Lys	Thr	Phe	Asn	Gly	Thr	Gly	Pro	Cys		
		195					200					205					
Thr	Asn	Val	Ser	Thr	Val	Gln	Cys	Thr	His	Gly	Ile	Arg	Pro	Val	Val		
	210					215					220						
Ser	Thr	Gln	Leu	Leu	Leu	Asn	Gly	Ser	Leu	Ala	Glu	Glu	Glu	Val	Val		
225				230						235				240			
Ile	Arg	Ser	Glu	Asn	Phe	Thr	Asn	Asn	Ala	Lys	Thr	Ile	Ile	Val	Gln		
				245					250					255			
Leu	Asn	Glu	Ser	Val	Glu	Ile	Asn	Cys	Thr	Arg	Pro	Asn	Asn	Asn	Thr		
			260					265					270				
Arg	Lys	Ser	Ile	His	Ile	Gly	Pro	Gly	Arg	Ala	Phe	Tyr	Thr	Thr	Gly		
	275						280					285					
Asp	Ile	Ile	Gly	Asp	Ile	Arg	Gln	Ala	His	Cys	Asn	Ile	Ser	Arg	Thr		
	290					295				300							
Asn	Trp	Thr	Asn	Thr	Leu	Lys	Arg	Val	Ala	Glu	Lys	Leu	Arg	Glu	Lys		
305					310					315				320			
Phe	Asn	Asn	Thr	Thr	Ile	Val	Phe	Asn	Gln	Ser	Ser	Gly	Gly	Asp	Pro		
				325					330					335			
Glu	Ile	Val	Met	His	Ser	Phe	Asn	Cys	Gly	Gly	Glu	Phe	Phe	Tyr	Cys		
			340					345					350				
Asn	Thr	Thr	Gln	Leu	Phe	Asn	Ser	Thr	Trp	Asn	Glu	Thr	Asn	Ser	Glu		
		355					360					365					
Gly	Asn	Ile	Thr	Ser	Gly	Thr	Ile	Thr	Leu	Pro	Cys	Arg	Ile	Lys	Gln		
	370					375					380						
Ile	Ile	Asn	Met	Trp	Gln	Glu	Val	Gly	Lys	Ala	Met	Tyr	Ala	Pro	Pro		
385					390					395				400			
Ile	Gly	Gly	Gln	Ile	Lys	Cys	Leu	Ser	Asn	Ile	Thr	Gly	Leu	Leu	Leu		
				405					410					415			
Thr	Arg	Asp	Gly	Gly	Ser	Asp	Asn	Ser	Ser	Ser	Gly	Lys	Glu	Ile	Phe		
			420					425					430				
Arg	Pro	Gly	Gly	Gly	Asp	Met	Arg	Asp	Asn	Trp	Arg	Ser	Glu	Leu	Tyr		
		435					440			</							

Gly Cys Ser Gly Lys Leu Ile Cys Thr Thr Ala Val Pro Trp Asn Ala  
                   565                  570                  575  
 Ser Trp Ser Asn Lys Asn Leu Ser Gln Ile Trp Asp Asn Met Thr Trp  
                   580                  585                  590  
 Met Glu Trp Glu Arg Glu Ile Ser Asn Tyr Thr Glu Ile Ile Tyr Ser  
                   595                  600                  605  
 Leu Ile Glu Glu Ser Gln Asn Gln Gln Lys Asn Glu Leu Asp Leu  
                   610                  615                  620  
 Leu Gln Leu Asp Lys Trp Ala Ser Leu Cys Asp Leu Arg Ile  
                   625                  630                  635

<210> 69  
 <211> 2071  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(2071)

<400> 69  
 gct agc gcg gcc gac cgc ctg tgg gtg acc gtg tac tac ggc gtg ccc 48  
 Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
   1                  5                  10                  15  
 gtg tgg aag gac gcc acc acc acc ctg ttc tgc gcc agc gac gcc aag 96  
 Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
                   20                  25                  30  
 gcc tac gac acc gag gtg cac aac gtg tgg gcc acc cac gcg tgc gtg 144  
 Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
                   35                  40                  45  
 ccc acc gac ccc aac ccc cag gag gtg gtg ctg ggc aac gtg acc gag 192  
 Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
                   50                  55                  60  
 aac ttc aac atg ggc aag aac aac atg gtg gag cag atg cac gag gat 240  
 Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp  
                   65                  70                  75                  80  
 atc atc agc ctg tgg gac cag agc ctg aag ccc tgc gtg aag ctg acc 288  
 Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr  
                   85                  90                  95  
 ccc ctg tgc gtg acc ctg aac tgc acc aag ctg aag aac agc acc gac 336  
 Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr Asp  
                   100                  105                  110  
 acc aac aac acc cgc tgg ggc acc cag gag atg aag aac tgc agc ttc 384  
 Thr Asn Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys Ser Phe  
                   115                  120                  125  
 aac atc agc acc agc gtg cgc aac aag atg aag cgc gag tac gcc ctg 432  
 Asn Ile Ser Thr Ser Val Arg Asn Lys Met Lys Arg Glu Tyr Ala Leu  
                   130                  135                  140  
 ttc tac agc ctg gac atc gtg ccc atc gac aac gac aac acc agc tac 480  
 Phe Tyr Ser Leu Asp Ile Val Pro Ile Asp Asn Asp Asn Thr Ser Tyr

145	150	155	160	
cgc ctg cgc agc tgc aac aca tcg atc atc acc cag gcc tgc ccc aag				528
Arg Leu Arg Ser Cys Asn Thr Ser Ile Ile Thr Gln Ala Cys Pro Lys	165	170	175	
gtg agc ttc gag ccc atc ccc atc cac ttc tgc gcc ccc gcc ggc ttc				576
Val Ser Phe Glu Pro Ile Pro Ile His Phe Cys Ala Pro Ala Gly Phe	180	185	190	
gcc atc ctg aag tgc aac aac aag acc ttc aac ggc acc ggc ccc tgc				624
Ala Ile Leu Lys Cys Asn Asn Lys Thr Phe Asn Gly Thr Gly Pro Cys	195	200	205	
acc aac gtg agc acc gtg cag tgc acc cac gga att cgc ccc gtg gtg				672
Thr Asn Val Ser Thr Val Gln Cys Thr His Gly Ile Arg Pro Val Val	210	215	220	
agc acc cag ctg ctg ctg aac ggc agc ctg gcc gag gag gag gtg gtg				720
Ser Thr Gln Leu Leu Asn Gly Ser Leu Ala Glu Glu Glu Val Val	225	230	235	240
atc aga tct gag aac ttc acc aac aac gcc aag acc atc atc gtg cag				768
Ile Arg Ser Glu Asn Phe Thr Asn Asn Ala Lys Thr Ile Ile Val Gln	245	250	255	
ctg aac gag agc gtg gag atc aac tgc acc cgc ccc aac aac aac acc				816
Leu Asn Glu Ser Val Glu Ile Asn Cys Thr Arg Pro Asn Asn Asn Thr	260	265	270	
cgc aag agc atc cac atc ggc cct ggc cgc gcc ttc tac acc acc ggc				864
Arg Lys Ser Ile His Ile Gly Pro Gly Arg Ala Phe Tyr Thr Thr Gly	275	280	285	
gac atc atc ggc gac atc cgc cag gcc cac tgc aac atc tct aga acc				912
Asp Ile Ile Gly Asp Ile Arg Gln Ala His Cys Asn Ile Ser Arg Thr	290	295	300	
aac tgg acc aac acc ctg aag cgc gtg gcc gag aag ctg cgc gag aag				960
Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu Arg Glu Lys	305	310	315	320
ttc aac aac acc acc atc gtg ttc aac cag agc tcc ggc ggc gac ccc				1008
Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser Ser Gly Gly Asp Pro	325	330	335	
gag atc gtg atg cac agc ttc aac tgc ggc ggc gag ttc ttc tac tgc				1056
Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe Phe Tyr Cys	340	345	350	
aac acc acc cag ctg ttc aac agc acc tgg aac gag acc aac agc gag				1104
Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr Asn Ser Glu	355	360	365	
ggc aac atc act agt ggc acc atc acc ctg ccc tgc cgc atc aag cag				1152
Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln	370	375	380	
atc atc aac atg tgg cag gag gtg ggc aag gcc atg tac gcc ccc ccc				1200

Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro	385	390	395	400
atc ggc ggc cag atc aag tgc ctg agc aac atc acc ggc ctg ctg ctg	1248			
Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu		405	410	415
acc cgc gac ggc ggc agc gac aac tgc agc agc ggc aag gag att ttc	1296			
Thr Arg Asp Gly Gly Ser Asp Asn Ser Ser Ser Gly Lys Glu Ile Phe		420	425	430
cgc ccc ggc ggc ggc gac atg cgc gac aac tgg cgc agc gag ctg tac	1344			
Arg Pro Gly Gly Gly Asp Met Arg Asp Asn Trp Arg Ser Glu Leu Tyr		435	440	445
aag tac aag gtg gtg aag atc gag ccc ctg ggc atc gcc ccc acc aag	1392			
Lys Tyr Lys Val Val Lys Ile Glu Pro Leu Gly Ile Ala Pro Thr Lys		450	455	460
gcc aag cgc cgc gtg gtg cag cgc gag aag cgc gcc gtg ggc atc ggc	1440			
Ala Lys Arg Arg Val Val Gln Arg Glu Lys Arg Ala Val Gly Ile Gly		465	470	475
gct atg ttc ctc ggc ttc ctg ggc gct gca ggc agc acc atg ggc gcc	1488			
Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly Ser Thr Met Gly Ala		485	490	495
gcc agc ctg acc ctg acc gtg cag gcc cgc cag ctg ctg agc ggc atc	1536			
Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln Leu Leu Ser Gly Ile		500	505	510
gtg cag cag cag aac aac ctg ctg cgc gcc atc gag gcc cag cag cac	1584			
Val Gln Gln Gln Asn Asn Leu Leu Arg Ala Ile Glu Ala Gln Gln His		515	520	525
ctg ctc cag ctg acc gtg tgg ggc atc aag cag ctc cag gcc cgc gtg	1632			
Leu Leu Gln Leu Thr Val Trp Gly Ile Lys Gln Leu Gln Ala Arg Val		530	535	540
ctg gct cta gag cgc tac ctc cag gac cag cgc ttc ctg ggc atg tgg	1680			
Leu Ala Leu Glu Arg Tyr Leu Gln Asp Gln Arg Phe Leu Gly Met Trp		545	550	555
ggc tgc tcc ggc aag ctg atc tgc acc acg gcc gtg ccc tgg aac gcc	1728			
Gly Cys Ser Gly Lys Leu Ile Cys Thr Thr Ala Val Pro Trp Asn Ala		565	570	575
agc tgg agc aac aag aac ctg agc cag att tgg gac aac atg acc tgg	1776			
Ser Trp Ser Asn Lys Asn Leu Ser Gln Ile Trp Asp Asn Met Thr Trp		580	585	590
atg gag tgg gag cgc gag atc agc aac tac acc gag atc atc tac agc	1824			
Met Glu Trp Glu Arg Glu Ile Ser Asn Tyr Thr Glu Ile Ile Tyr Ser		595	600	605
ctg atc gag gag agc cag aac cag cag gag aag aac gag ctg gac ctg	1872			
Leu Ile Glu Glu Ser Gln Asn Gln Gln Glu Lys Asn Glu Leu Asp Leu		610	615	620



ctc cag ctg gac aag tgg gca agc ttg tgg aac tgg ttc aac atc acc 1920  
 Leu Gln Leu Asp Lys Trp Ala Ser Leu Trp Asn Trp Phe Asn Ile Thr  
 625 630 635 640  
  
 aac tgg ctg tgg tac atc aag att ttc atc atg atc gtg ggc ggc ctg 1968  
 Asn Trp Leu Trp Tyr Ile Lys Ile Phe Ile Met Ile Val Gly Gly Leu  
 645 650 655  
  
 atc ggc ctg cgc atc gtg ttc acc gtg ctg agc atc gtg aac cgc gtg 2016  
 Ile Gly Leu Arg Ile Val Phe Thr Val Leu Ser Ile Val Asn Arg Val  
 660 665 670  
  
 cgc cag gga tgc agc ccc ctg agc ttc cag acc cgc ctg ccc gtg tga 2064  
 Arg Gln Gly Cys Ser Pro Leu Ser Phe Gln Thr Arg Leu Pro Val \*  
 675 680 685  
  
 cgg atc c 2071  
 Arg Ile

<210> 70  
 <211> 689  
 <212> PRT  
 <213> Artificial Sequence

<400> 70  
 Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
 1 5 10 15  
 Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
 20 25 30  
 Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
 35 40 45  
 Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
 50 55 60  
 Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp  
 65 70 75 80  
 Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr  
 85 90 95  
 Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr Asp  
 100 105 110  
 Thr Asn Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys Ser Phe  
 115 120 125  
 Asn Ile Ser Thr Ser Val Arg Asn Lys Met Lys Arg Glu Tyr Ala Leu  
 130 135 140  
 Phe Tyr Ser Leu Asp Ile Val Pro Ile Asp Asn Asp Asn Thr Ser Tyr  
 145 150 155 160  
 Arg Leu Arg Ser Cys Asn Thr Ser Ile Ile Thr Gln Ala Cys Pro Lys  
 165 170 175  
 Val Ser Phe Glu Pro Ile Pro Ile His Phe Cys Ala Pro Ala Gly Phe  
 180 185 190  
 Ala Ile Leu Lys Cys Asn Asn Lys Thr Phe Asn Gly Thr Gly Pro Cys  
 195 200 205  
 Thr Asn Val Ser Thr Val Gln Cys Thr His Gly Ile Arg Pro Val Val  
 210 215 220  
 Ser Thr Gln Leu Leu Leu Asn Gly Ser Leu Ala Glu Glu Glu Val Val  
 225 230 235 240  
 Ile Arg Ser Glu Asn Phe Thr Asn Asn Ala Lys Thr Ile Ile Val Gln  
 245 250 255

Leu Asn Glu Ser Val Glu Ile Asn Cys Thr Arg Pro Asn Asn Asn Thr  
 260 265 270  
 Arg Lys Ser Ile His Ile Gly Pro Gly Arg Ala Phe Tyr Thr Thr Gly  
 275 280 285  
 Asp Ile Ile Gly Asp Ile Arg Gln Ala His Cys Asn Ile Ser Arg Thr  
 290 295 300  
 Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu Arg Glu Lys  
 305 310 315 320  
 Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser Ser Gly Gly Asp Pro  
 325 330 335  
 Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe Phe Tyr Cys  
 340 345 350  
 Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr Asn Ser Glu  
 355 360 365  
 Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln  
 370 375 380  
 Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro  
 385 390 395 400  
 Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu  
 405 410 415  
 Thr Arg Asp Gly Gly Ser Asp Asn Ser Ser Ser Gly Lys Glu Ile Phe  
 420 425 430  
 Arg Pro Gly Gly Gly Asp Met Arg Asp Asn Trp Arg Ser Glu Leu Tyr  
 435 440 445  
 Lys Tyr Lys Val Val Lys Ile Glu Pro Leu Gly Ile Ala Pro Thr Lys  
 450 455 460  
 Ala Lys Arg Arg Val Val Gln Arg Glu Lys Arg Ala Val Gly Ile Gly  
 465 470 475 480  
 Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly Ser Thr Met Gly Ala  
 485 490 495  
 Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln Leu Leu Ser Gly Ile  
 500 505 510  
 Val Gln Gln Gln Asn Asn Leu Leu Arg Ala Ile Glu Ala Gln Gln His  
 515 520 525  
 Leu Leu Gln Leu Thr Val Trp Gly Ile Lys Gln Leu Gln Ala Arg Val  
 530 535 540  
 Leu Ala Leu Glu Arg Tyr Leu Gln Asp Gln Arg Phe Leu Gly Met Trp  
 545 550 555 560  
 Gly Cys Ser Gly Lys Leu Ile Cys Thr Thr Ala Val Pro Trp Asn Ala  
 565 570 575  
 Ser Trp Ser Asn Lys Asn Leu Ser Gln Ile Trp Asp Asn Met Thr Trp  
 580 585 590  
 Met Glu Trp Glu Arg Glu Ile Ser Asn Tyr Thr Glu Ile Ile Tyr Ser  
 595 600 605  
 Leu Ile Glu Glu Ser Gln Asn Gln Gln Glu Lys Asn Glu Leu Asp Leu  
 610 615 620  
 Leu Gln Leu Asp Lys Trp Ala Ser Leu Trp Asn Trp Phe Asn Ile Thr  
 625 630 635 640  
 Asn Trp Leu Trp Tyr Ile Lys Ile Phe Ile Met Ile Val Gly Gly Leu  
 645 650 655  
 Ile Gly Leu Arg Ile Val Phe Thr Val Leu Ser Ile Val Asn Arg Val  
 660 665 670  
 Arg Gln Gly Cys Ser Pro Leu Ser Phe Gln Thr Arg Leu Pro Val Arg  
 675 680 685  
 Ile

&lt;210&gt; 71

&lt;211&gt; 2469

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)...(2469)

&lt;400&gt; 71

gct agc gcg gcc gac cgc ctg tgg gtg acc gtg tac tac ggc gtg ccc	48
Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro	
1 5 10 15	
gtg tgg aag gac gcc acc acc acc ctg ttc tgc gcc agc gac gcc aag	96
Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys	
20 25 30	
gcc tac gac acc gag gtg cac aac gtg tgg gcc acc cac gcg tgc gtg	144
Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val	
35 40 45	
ccc acc gac ccc aac ccc cag gag gtg gtg ctg ggc aac gtg acc gag	192
Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu	
50 55 60	
aac ttc aac atg ggc aag aac aac atg gtg gag cag atg cac gag gat	240
Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp	
65 70 75 80	
atc atc agc ctg tgg gac cag agc ctg aag ccc tgc gtg aag ctg acc	288
Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr	
85 90 95	
ccc ctg tgc gtg acc ctg aac tgc acc aag ctg aag aac agc acc gac	336
Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr Asp	
100 105 110	
acc aac aac acc cgc tgg ggc acc cag gag atg aag aac tgc agc ttc	384
Thr Asn Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys Ser Phe	
115 120 125	
aac atc agc acc agc gtg cgc aac aag atg aag cgc gag tac gcc ctg	432
Asn Ile Ser Thr Ser Val Arg Asn Lys Met Lys Arg Glu Tyr Ala Leu	
130 135 140	
ttc tac agc ctg gac atc gtg ccc atc gac aac gac aac acc agc tac	480
Phe Tyr Ser Leu Asp Ile Val Pro Ile Asp Asn Asp Asn Thr Ser Tyr	
145 150 155 160	
cgc ctg cgc agc tgc aac aca tcg atc atc acc cag gcc tgc ccc aag	528
Arg Leu Arg Ser Cys Asn Thr Ser Ile Ile Thr Gln Ala Cys Pro Lys	
165 170 175	
gtg agc ttc gag ccc atc ccc atc cac ttc tgc gcc ccc gcc ggc ttc	576
Val Ser Phe Glu Pro Ile Pro Ile His Phe Cys Ala Pro Ala Gly Phe	
180 185 190	
gcc atc ctg aag tgc aac aac aag acc ttc aac ggc acc ggc ccc tgc	624
Ala Ile Leu Lys Cys Asn Asn Lys Thr Phe Asn Gly Thr Gly Pro Cys	
195 200 205	

acc aac gtg agc acc gtg cag tgc acc cac gga att cgc ccc gtg gtg Thr Asn Val Ser Thr Val Gln Cys Thr His Gly Ile Arg Pro Val Val 210 215 220	672
agc acc cag ctg ctg ctg aac ggc agc ctg gcc gag gag gag gtg gtg Ser Thr Gln Leu Leu Leu Asn Gly Ser Leu Ala Glu Glu Glu Val Val 225 230 235 240	720
atc aga tct gag aac ttc acc aac aac gcc aag acc atc atc gtg cag Ile Arg Ser Glu Asn Phe Thr Asn Asn Ala Lys Thr Ile Ile Val Gln 245 250 255	768
ctg aac gag agc gtg gag atc aac tgc acc cgc ccc aac aac aac acc Leu Asn Glu Ser Val Glu Ile Asn Cys Thr Arg Pro Asn Asn Asn Thr 260 265 270	816
cgc aag agc atc cac atc ggc cct ggc cgc gcc ttc tac acc acc ggc Arg Lys Ser Ile His Ile Gly Pro Gly Arg Ala Phe Tyr Thr Thr Gly 275 280 285	864
gac atc atc ggc gac atc cgc cag gcc cac tgc aac atc tct aga acc Asp Ile Ile Gly Asp Ile Arg Gln Ala His Cys Asn Ile Ser Arg Thr 290 295 300	912
aac tgg acc aac acc ctg aag cgc gtg gcc gag aag ctg cgc gag aag Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu Arg Glu Lys 305 310 315 320	960
ttc aac aac acc acc atc gtg ttc aac cag agc tcc ggc ggc gac ccc Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser Ser Gly Gly Asp Pro 325 330 335	1008
gag atc gtg atg cac agc ttc aac tgc ggc ggc gag ttc ttc tac tgc Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe Phe Tyr Cys 340 345 350	1056
aac acc acc cag ctg ttc aac agc acc tgg aac gag acc aac agc gag Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr Asn Ser Glu 355 360 365	1104
ggc aac atc act agt ggc acc atc acc ctg ccc tgc cgc atc aag cag Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln 370 375 380	1152
atc atc aac atg tgg cag gag gtg ggc aag gcc atg tac gcc ccc ccc Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro 385 390 395 400	1200
atc ggc ggc cag atc aag tgc ctg agc aac atc acc ggc ctg ctg ctg Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu 405 410 415	1248
acc cgc gac ggc ggc agc gac aac tgc agc agc ggc aag gag att ttc Thr Arg Asp Gly Gly Ser Asp Asn Ser Ser Ser Gly Lys Glu Ile Phe 420 425 430	1296
cgc ccc ggc ggc ggc gac atg cgc gac aac tgg cgc agc gag ctg tac Arg Pro Gly Gly Gly Asp Met Arg Asp Asn Trp Arg Ser Glu Leu Tyr	1344

435	440	445	
aag tac aag gtg gtg aag atc gag ccc ctg ggc atc gcc ccc acc aag Lys Tyr Lys Val Val Lys Ile Glu Pro Leu Gly Ile Ala Pro Thr Lys 450 455 460			1392
gcc aag cgc cgc gtg gtg cag cgc gag aag cgc gcc gtg ggc atc ggc Ala Lys Arg Arg Val Val Gln Arg Glu Lys Arg Ala Val Gly Ile Gly 465 470 475 480			1440
gct atg ttc ctc ggc ttc ctg ggc gct gca ggc agc acc atg ggc gcc Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly Ser Thr Met Gly Ala 485 490 495			1488
gcc agc ctg acc ctg acc gtg cag gcc cgc cag ctg ctg agc ggc atc Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln Leu Leu Ser Gly Ile 500 505 510			1536
gtg cag cag cag aac aac ctg ctg cgc gcc atc gag gcc cag cag cac Val Gln Gln Gln Asn Asn Leu Leu Arg Ala Ile Glu Ala Gln Gln His 515 520 525			1584
ctg ctc cag ctg acc gtg tgg ggc atc aag cag ctc cag gcc cgc gtg Leu Leu Gln Leu Thr Val Trp Gly Ile Lys Gln Leu Gln Ala Arg Val 530 535 540			1632
ctg gct cta gag cgc tac ctc cag gac cag cgc ttc ctg ggc atg tgg Leu Ala Leu Glu Arg Tyr Leu Gln Asp Gln Arg Phe Leu Gly Met Trp 545 550 555 560			1680
ggc tgc tcc ggc aag ctg atc tgc acc acg gcc gtg ccc tgg aac gcc Gly Cys Ser Gly Lys Leu Ile Cys Thr Thr Ala Val Pro Trp Asn Ala 565 570 575			1728
agc tgg agc aac aag aac ctg agc cag att tgg gac aac atg acc tgg Ser Trp Ser Asn Lys Asn Leu Ser Gln Ile Trp Asp Asn Met Thr Trp 580 585 590			1776
atg gag tgg gag cgc gag atc agc aac tac acc gag atc atc tac agc Met Glu Trp Glu Arg Glu Ile Ser Asn Tyr Thr Glu Ile Ile Tyr Ser 595 600 605			1824
ctg atc gag gag agc cag aac cag cag gag aag aac gag ctg gac ctg Leu Ile Glu Glu Ser Gln Asn Gln Gln Glu Lys Asn Glu Leu Asp Leu 610 615 620			1872
ctc cag ctg gac aag tgg gca agc ttg tgg aac tgg ttc aac atc acc Leu Gln Leu Asp Lys Trp Ala Ser Leu Trp Asn Trp Phe Asn Ile Thr 625 630 635 640			1920
aac tgg ctg tgg tac atc aag att ttc atc atg atc gtg ggc ggc ctg Asn Trp Leu Trp Tyr Ile Lys Ile Phe Ile Met Ile Val Gly Gly Leu 645 650 655			1968
atc ggc ctg cgc atc gtg ttc acc gtg ctg agc atc gtg aac cgc gtg Ile Gly Leu Arg Ile Val Phe Thr Val Leu Ser Ile Val Asn Arg Val 660 665 670			2016
cgc cag ggc tac agc ccc ctg agc ttc cag acc cgc ctg ccc gtg ccc			2064

Arg	Gln	Gly	Tyr	Ser	Pro	Leu	Ser	Phe	Gln	Thr	Arg	Leu	Pro	Val	Pro		
		675					680					685					
cgc	ggc	ccc	gac	cgc	ccc	gag	ggc	atc	gag	gag	gag	ggc	ggc	gag	cgc		2112
Arg	Gly	Pro	Asp	Arg	Pro	Glu	Gly	Ile	Glu	Glu	Glu	Gly	Gly	Glu	Arg		
		690				695					700						
gac	cgc	gac	cgc	agc	acc	cgc	ctg	gtg	acc	ggc	ttc	ctg	ccc	ctg	atc		2160
Asp	Arg	Asp	Arg	Ser	Thr	Arg	Leu	Val	Thr	Gly	Phe	Leu	Pro	Leu	Ile		
		705			710					715					720		
tgg	gac	gac	ctg	cgc	agc	ctg	ttc	ctg	ttc	agc	tac	cat	cga	ttg	cgc		2208
<del>Trp</del>	Asp	Asp	Leu	Arg	Ser	Leu	Phe	Leu	Phe	Ser	Tyr	His	Arg	Leu	Arg		
			725					730						735			
gac	ctg	ctg	ctg	atc	gtg	gcc	cgc	atc	gtg	gag	ctg	ctg	ggc	cgg	cgc		2256
Asp	Leu	Leu	Leu	Ile	Val	Ala	Arg	Ile	Val	Glu	Leu	Leu	Gly	Arg	Arg		
			740				745						750				
ggc	tgg	gag	atc	ctg	aag	tac	tgg	tgg	aac	ctg	ctc	cag	tac	tgg	agc		2304
Gly	Trp	Glu	Ile	Leu	Lys	Tyr	Trp	Trp	Asn	Leu	Leu	Gln	Tyr	Trp	Ser		
		755					760					765					
cag	gag	ctg	aag	aac	tct	gca	gtg	agc	ctg	ctg	aac	gcc	acc	gcc	atc		2352
Gln	Glu	Leu	Lys	Asn	Ser	Ala	Val	Ser	Leu	Leu	Asn	Ala	Thr	Ala	Ile		
		770				775					780						
gcc	gtg	gcc	gag	ggc	acc	gac	cgc	gtg	atc	gag	gtg	gtg	cag	cgc	atc		2400
Ala	Val	Ala	Glu	Gly	Thr	Asp	Arg	Val	Ile	Glu	Val	Val	Gln	Arg	Ile		
		785			790				795					800			
tgg	cgc	ggc	atc	ctg	cac	atc	ccc	acc	cga	att	cgc	cag	ggc	ttc	gag		2448
Trp	Arg	Gly	Ile	Leu	His	Ile	Pro	Thr	Arg	Ile	Arg	Gln	Gly	Phe	Glu		
			805						810					815			
cgc	gcc	ctg	ctg	taa	gga	tcc											2469
Arg	Ala	Leu	Leu	*	Gly	Ser											
			820														

<210> 72  
 <211> 822  
 <212> PRT  
 <213> Artificial Sequence

<400> 72  
 Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
 1 5 10 15  
 Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
 20 25 30  
 Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
 35 40 45  
 Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
 50 55 60  
 Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp  
 65 70 75 80  
 Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr  
 85 90 95  
 Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr Asp



Ser Trp Ser Asn Lys Asn Leu Ser Gln Ile Trp Asp Asn Met Thr Trp  
 580 585 590  
 Met Glu Trp Glu Arg Glu Ile Ser Asn Tyr Thr Glu Ile Ile Tyr Ser  
 595 600 605  
 Leu Ile Glu Glu Ser Gln Asn Gln Gln Glu Lys Asn Glu Leu Asp Leu  
 610 615 620  
 Leu Gln Leu Asp Lys Trp Ala Ser Leu Trp Asn Trp Phe Asn Ile Thr  
 625 630 635 640  
 Asn Trp Leu Trp Tyr Ile Lys Ile Phe Ile Met Ile Val Gly Gly Leu  
 645 650 655  
 Ile Gly Leu Arg Ile Val Phe Thr Val Leu Ser Ile Val Asn Arg Val  
 660 665 670  
~~Arg~~ Gln Gly Tyr Ser Pro Leu Ser Phe-Gln Thr Arg Leu Pro Val Pro  
 675 680 685  
 Arg Gly Pro Asp Arg Pro Glu Gly Ile Glu Glu Glu Gly Gly Glu Arg  
 690 695 700  
 Asp Arg Asp Arg Ser Thr Arg Leu Val Thr Gly Phe Leu Pro Leu Ile  
 705 710 715 720  
 Trp Asp Asp Leu Arg Ser Leu Phe Leu Phe Ser Tyr His Arg Leu Arg  
 725 730 735  
 Asp Leu Leu Leu Ile Val Ala Arg Ile Val Glu Leu Leu Gly Arg Arg  
 740 745 750  
 Gly Trp Glu Ile Leu Lys Tyr Trp Trp Asn Leu Leu Gln Tyr Trp Ser  
 755 760 765  
 Gln Glu Leu Lys Asn Ser Ala Val Ser Leu Leu Asn Ala Thr Ala Ile  
 770 775 780  
 Ala Val Ala Glu Gly Thr Asp Arg Val Ile Glu Val Val Gln Arg Ile  
 785 790 795 800  
 Trp Arg Gly Ile Leu His Ile Pro Thr Arg Ile Arg Gln Gly Phe Glu  
 805 810 815  
 Arg Ala Leu Leu Gly Ser  
 820

<210> 73  
 <211> 1431  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> CDS  
 <222> (1)...(1431)

<400> 73  
 gct agc gcg gcc gac cgc ctg tgg gtg acc gtg tac tac ggc gtg ccc 48  
 Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
 1 5 10 15  
 gtg tgg aag gac gcc acc acc acc ctg ttc tgc gcc agc gac gcc aag 96  
 Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
 20 25 30  
 gcc tac gac acc gag gtg cac aac gtg tgg gcc acc cac gcg tgc gtg 144  
 Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
 35 40 45  
 ccc acc gac ccc aac ccc cag gag gtg gtg ctg ggc aac gtg acc gag 192  
 Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu  
 50 55 60



aac ttc aac atg ggc aag aac aac atg gtg gag cag atg cac gag gat Asn Phe Asn Met Gly Lys Asn Asn Met Val Glu Gln Met His Glu Asp 65 70 75 80	240
atc atc agc ctg tgg gac cag agc ctg aag ccc tgc gtg aag ctg acc Ile Ile Ser Leu Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr 85 90 95	288
ccc ctg tgc gtg acc ctg aac tgc acc aag ctg aag aac agc acc gac Pro Leu Cys Val Thr Leu Asn Cys Thr Lys Leu Lys Asn Ser Thr Asp 100 105 110	336
<del>acc</del> aac aac acc cgc tgg ggc acc <del>cag</del> -gag atg aag aac tgc agc ttc Thr Asn Asn Thr Arg Trp Gly Thr Gln Glu Met Lys Asn Cys Ser Phe 115 120 125	384
aac atc agc acc agc gtg cgc aac aag atg aag cgc gag tac gcc ctg Asn Ile Ser Thr Ser Val Arg Asn Lys Met Lys Arg Glu Tyr Ala Leu 130 135 140	432
ttc tac agc ctg gac atc gtg ccc atc gac aac gac aac acc agc tac Phe Tyr Ser Leu Asp Ile Val Pro Ile Asp Asn Asp Asn Thr Ser Tyr 145 150 155 160	480
cgc ctg cgc agc tgc aac aca tcg atc atc acc cag gcc tgc ccc aag Arg Leu Arg Ser Cys Asn Thr Ser Ile Ile Thr Gln Ala Cys Pro Lys 165 170 175	528
gtg agc ttc gag ccc atc ccc atc cac ttc tgc gcc ccc gcc ggc ttc Val Ser Phe Glu Pro Ile Pro Ile His Phe Cys Ala Pro Ala Gly Phe 180 185 190	576
gcc atc ctg aag tgc aac aac aag acc ttc aac ggc acc ggc ccc tgc Ala Ile Leu Lys Cys Asn Asn Lys Thr Phe Asn Gly Thr Gly Pro Cys 195 200 205	624
acc aac gtg agc acc gtg cag tgc acc cac gga att cgc ccc gtg gtg Thr Asn Val Ser Thr Val Gln Cys Thr His Gly Ile Arg Pro Val Val 210 215 220	672
agc acc cag ctg ctg ctg aac ggc agc ctg gcc gag gag gag gtg gtg Ser Thr Gln Leu Leu Leu Asn Gly Ser Leu Ala Glu Glu Glu Val Val 225 230 235 240	720
atc aga tct gag aac ttc acc aac aac gcc aag acc atc atc gtg cag Ile Arg Ser Glu Asn Phe Thr Asn Asn Ala Lys Thr Ile Ile Val Gln 245 250 255	768
ctg aac gag agc gtg gag atc aac tgc acc cgc ccc aac aac aac acc Leu Asn Glu Ser Val Glu Ile Asn Cys Thr Arg Pro Asn Asn Asn Thr 260 265 270	816
cgc aag agc atc cac atc ggc cct ggc cgc gcc ttc tac acc acc ggc Arg Lys Ser Ile His Ile Gly Pro Gly Arg Ala Phe Tyr Thr Thr Gly 275 280 285	864
gac atc atc ggc gac atc cgc cag gcc cac tgc aac atc tct aga acc Asp Ile Ile Gly Asp Ile Arg Gln Ala His Cys Asn Ile Ser Arg Thr 290 295 300	912

aac tgg acc aac acc ctg aag cgc gtg gcc gag aag ctg cgc gag aag 960  
 Asn Trp Thr Asn Thr Leu Lys Arg Val Ala Glu Lys Leu Arg Glu Lys  
 305 310 315 320

ttc aac aac acc acc atc gtg ttc aac cag agc tcc ggc ggc gac ccc 1008  
 Phe Asn Asn Thr Thr Ile Val Phe Asn Gln Ser Ser Gly Gly Asp Pro  
 325 330 335

gag atc gtg atg cac agc ttc aac tgc ggc ggc gag ttc ttc tac tgc 1056  
 Glu Ile Val Met His Ser Phe Asn Cys Gly Gly Glu Phe Phe Tyr Cys  
 340 345 350

aac acc acc cag ctg ttc aac agc acc tgg aac gag acc aac agc gag 1104  
 Asn Thr Thr Gln Leu Phe Asn Ser Thr Trp Asn Glu Thr Asn Ser Glu  
 355 360 365

ggc aac atc act agt ggc acc atc acc ctg ccc tgc cgc atc aag cag 1152  
 Gly Asn Ile Thr Ser Gly Thr Ile Thr Leu Pro Cys Arg Ile Lys Gln  
 370 375 380

atc atc aac atg tgg cag gag gtg ggc aag gcc atg tac gcc ccc ccc 1200  
 Ile Ile Asn Met Trp Gln Glu Val Gly Lys Ala Met Tyr Ala Pro Pro  
 385 390 395 400

atc ggc ggc cag atc aag tgc ctg agc aac atc acc ggc ctg ctg ctg 1248  
 Ile Gly Gly Gln Ile Lys Cys Leu Ser Asn Ile Thr Gly Leu Leu Leu  
 405 410 415

acc cgc gac ggc ggc agc gac aac tcg agc agc ggc aag gag att ttc 1296  
 Thr Arg Asp Gly Gly Ser Asp Asn Ser Ser Ser Gly Lys Glu Ile Phe  
 420 425 430

cgc ccc ggc ggc ggc gac atg cgc gac aac tgg cgc agc gag ctg tac 1344  
 Arg Pro Gly Gly Gly Asp Met Arg Asp Asn Trp Arg Ser Glu Leu Tyr  
 435 440 445

aag tac aag gtg gtg aag atc gag ccc ctg ggc atc gcc ccc acc aag 1392  
 Lys Tyr Lys Val Val Lys Ile Glu Pro Leu Gly Ile Ala Pro Thr Lys  
 450 455 460

gcc aag cgc cgc gtg gtg cag cgc gag aag cgc gcc tag 1431  
 Ala Lys Arg Arg Val Val Gln Arg Glu Lys Arg Ala \*  
 465 470 475

&lt;210&gt; 74

&lt;211&gt; 476

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;400&gt; 74

Ala Ser Ala Ala Asp Arg Leu Trp Val Thr Val Tyr Tyr Gly Val Pro  
 1 5 10 15  
 Val Trp Lys Asp Ala Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys  
 20 25 30  
 Ala Tyr Asp Thr Glu Val His Asn Val Trp Ala Thr His Ala Cys Val  
 35 40 45  
 Pro Thr Asp Pro Asn Pro Gln Glu Val Val Leu Gly Asn Val Thr Glu

50	55	60
Asn Phe Asn Met Gly Lys	Asn Asn Met Val Glu	Gln Met His Glu Asp
65	70	75
Ile Ile Ser Leu Trp Asp	Gln Ser Leu Lys Pro	Cys Val Lys Leu Thr
85	90	95
Pro Leu Cys Val Thr Leu	Asn Cys Thr Lys Leu	Lys Asn Ser Thr Asp
100	105	110
Thr Asn Asn Thr Arg Trp	Gly Thr Gln Glu Met	Lys Asn Cys Ser Phe
115	120	125
Asn Ile Ser Thr Ser Val	Arg Asn Lys Met Lys	Arg Glu Tyr Ala Leu
130	135	140
Phe Tyr Ser Leu Asp Ile	Val Pro Ile Asp Asn	Asp Asn Thr Ser Tyr
145	150	155
Arg Leu Arg Ser Cys Asn	Thr Ser Ile Ile Thr	Gln Ala Cys Pro Lys
165	170	175
Val Ser Phe Glu Pro Ile	Pro Ile His Phe Cys	Ala Pro Ala Gly Phe
180	185	190
Ala Ile Leu Lys Cys Asn	Asn Lys Thr Phe Asn	Gly Thr Gly Pro Cys
195	200	205
Thr Asn Val Ser Thr Val	Gln Cys Thr His Gly	Ile Arg Pro Val Val
210	215	220
Ser Thr Gln Leu Leu Leu	Asn Gly Ser Leu Ala	Glu Glu Glu Val Val
225	230	235
Ile Arg Ser Glu Asn Phe	Thr Asn Asn Ala Lys	Thr Ile Ile Val Gln
245	250	255
Leu Asn Glu Ser Val Glu	Ile Asn Cys Thr Arg	Pro Asn Asn Asn Thr
260	265	270
Arg Lys Ser Ile His Ile	Gly Pro Gly Arg Ala	Phe Tyr Thr Thr Gly
275	280	285
Asp Ile Ile Gly Asp Ile	Arg Gln Ala His Cys	Asn Ile Ser Arg Thr
290	295	300
Asn Trp Thr Asn Thr Leu	Lys Arg Val Ala Glu	Lys Leu Arg Glu Lys
305	310	315
Phe Asn Asn Thr Thr Ile	Val Phe Asn Gln Ser	Ser Gly Gly Asp Pro
325	330	335
Glu Ile Val Met His Ser	Phe Asn Cys Gly Gly	Glu Phe Phe Tyr Cys
340	345	350
Asn Thr Thr Gln Leu Phe	Asn Ser Thr Trp Asn	Glu Thr Asn Ser Glu
355	360	365
Gly Asn Ile Thr Ser Gly	Thr Ile Thr Leu Pro	Cys Arg Ile Lys Gln
370	375	380
Ile Ile Asn Met Trp Gln	Glu Val Gly Lys Ala	Met Tyr Ala Pro Pro
385	390	395
Ile Gly Gly Gln Ile Lys	Cys Leu Ser Asn Ile	Thr Gly Leu Leu Leu
405	410	415
Thr Arg Asp Gly Gly Ser	Asp Asn Ser Ser Ser	Gly Lys Glu Ile Phe
420	425	430
Arg Pro Gly Gly Asp Met	Arg Asp Asn Trp Arg	Ser Glu Leu Tyr
435	440	445
Lys Tyr Lys Val Val Lys	Ile Glu Pro Leu Gly	Ile Ala Pro Thr Lys
450	455	460
Ala Lys Arg Arg Val Val	Gln Arg Glu Lys Arg	Ala
465	470	475

&lt;210&gt; 75

&lt;211&gt; 1038

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)...(1038)

&lt;400&gt; 75

gcc gtg ggc atc ggc gct atg ttc ctc ggc ttc ctg ggc gct gca ggc	48
Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly	
1 5 10 15	
agc acc atg ggc gcc gcc agc ctg acc ctg acc gtg cag gcc cgc cag	96
Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln	
20 25 30	
ctg ctg agc ggc atc gtg cag cag cag aac aac ctg ctg cgc gcc atc	144
Leu Leu Ser Gly Ile Val Gln Gln Gln Asn Asn Leu Leu Arg Ala Ile	
35 40 45	
gag gcc cag cag cac ctg ctc cag ctg acc gtg tgg ggc atc aag cag	192
Glu Ala Gln Gln His Leu Leu Gln Leu Thr Val Trp Gly Ile Lys Gln	
50 55 60	
ctc cag gcc cgc gtg ctg gct cta gag cgc tac ctc cag gac cag cgc	240
Leu Gln Ala Arg Val Leu Ala Leu Glu Arg Tyr Leu Gln Asp Gln Arg	
65 70 75 80	
ttc ctg ggc atg tgg ggc tgc tcc ggc aag ctg atc tgc acc acg gcc	288
Phe Leu Gly Met Trp Gly Cys Ser Gly Lys Leu Ile Cys Thr Thr Ala	
85 90 95	
gtg ccc tgg aac gcc agc tgg agc aac aag aac ctg agc cag att tgg	336
Val Pro Trp Asn Ala Ser Trp Ser Asn Lys Asn Leu Ser Gln Ile Trp	
100 105 110	
gac aac atg acc tgg atg gag tgg gag cgc gag atc agc aac tac acc	384
Asp Asn Met Thr Trp Met Glu Trp Glu Arg Glu Ile Ser Asn Tyr Thr	
115 120 125	
gag atc atc tac agc ctg atc gag gag agc cag aac cag cag gag aag	432
Glu Ile Ile Tyr Ser Leu Ile Glu Glu Ser Gln Asn Gln Gln Glu Lys	
130 135 140	
aac gag ctg gac ctg ctc cag ctg gac aag tgg gca agc ttg tgg aac	480
Asn Glu Leu Asp Leu Leu Gln Leu Asp Lys Trp Ala Ser Leu Trp Asn	
145 150 155 160	
tgg ttc aac atc acc aac tgg ctg tgg tac atc aag att ttc atc atg	528
Trp Phe Asn Ile Thr Asn Trp Leu Trp Tyr Ile Lys Ile Phe Ile Met	
165 170 175	
atc gtg ggc ggc ctg atc ggc ctg cgc atc gtg ttc acc gtg ctg agc	576
Ile Val Gly Gly Leu Ile Gly Leu Arg Ile Val Phe Thr Val Leu Ser	
180 185 190	
atc gtg aac cgc gtg cgc cag ggc tac agc ccc ctg agc ttc cag acc	624
Ile Val Asn Arg Val Arg Gln Gly Tyr Ser Pro Leu Ser Phe Gln Thr	
195 200 205	
cgc ctg ccc gtg ccc cgc ggc ccc gac cgc ccc gag ggc atc gag gag	672
Arg Leu Pro Val Pro Arg Gly Pro Asp Arg Pro Glu Gly Ile Glu Glu	

210	215	220	
gag ggc ggc gag cgc gac cgc gac cgc agc acc cgc ctg gtg acc ggc Glu Gly Gly Glu Arg Asp Arg Asp Arg Ser Thr Arg Leu Val Thr Gly 225 230 235 240			720
ttc ctg ccc ctg atc tgg gac gac ctg cgc agc ctg ttc ctg ttc agc Phe Leu Pro Leu Ile Trp Asp Asp Leu Arg Ser Leu Phe Leu Phe Ser 245 250 255			768
tac cat cga ttg cgc gac ctg ctg ctg atc gtg gcc cgc atc gtg gag Tyr His Arg Leu Arg Asp Leu Leu Leu Ile Val Ala Arg Ile Val Glu 260 265 270			816
ctg ctg ggc cgg cgc ggc tgg gag atc ctg aag tac tgg tgg aac ctg Leu Leu Gly Arg Arg Gly Trp Glu Ile Leu Lys Tyr Trp Trp Asn Leu 275 280 285			864
ctc cag tac tgg agc cag gag ctg aag aac tct gca gtg agc ctg ctg Leu Gln Tyr Trp Ser Gln Glu Leu Lys Asn Ser Ala Val Ser Leu Leu 290 295 300			912
aac gcc acc gcc atc gcc gtg gcc gag ggc acc gac cgc gtg atc gag Asn Ala Thr Ala Ile Ala Val Ala Glu Gly Thr Asp Arg Val Ile Glu 305 310 315 320			960
gtg gtg cag cgc atc tgg cgc ggc atc ctg cac atc ccc acc cga att Val Val Gln Arg Ile Trp Arg Gly Ile Leu His Ile Pro Thr Arg Ile 325 330 335			1008
cgc cag ggc ttc gag cgc gcc ctg ctg taa Arg Gln Gly Phe Glu Arg Ala Leu Leu *			1038
340 345			

<210> 76  
 <211> 345  
 <212> PRT  
 <213> Artificial Sequence

<400> 76  
 Ala Val Gly Ile Gly Ala Met Phe Leu Gly Phe Leu Gly Ala Ala Gly  
 1 5 10 15  
 Ser Thr Met Gly Ala Ala Ser Leu Thr Leu Thr Val Gln Ala Arg Gln  
 20 25 30  
 Leu Leu Ser Gly Ile Val Gln Gln Gln Asn Asn Leu Leu Arg Ala Ile  
 35 40 45  
 Glu Ala Gln Gln His Leu Leu Gln Leu Thr Val Trp Gly Ile Lys Gln  
 50 55 60  
 Leu Gln Ala Arg Val Leu Ala Leu Glu Arg Tyr Leu Gln Asp Gln Arg  
 65 70 75 80  
 Phe Leu Gly Met Trp Gly Cys Ser Gly Lys Leu Ile Cys Thr Thr Ala  
 85 90 95  
 Val Pro Trp Asn Ala Ser Trp Ser Asn Lys Asn Leu Ser Gln Ile Trp  
 100 105 110  
 Asp Asn Met Thr Trp Met Glu Trp Glu Arg Glu Ile Ser Asn Tyr Thr  
 115 120 125  
 Glu Ile Ile Tyr Ser Leu Ile Glu Glu Ser Gln Asn Gln Gln Glu Lys  
 130 135 140

Asn Glu Leu Asp Leu Leu Gln Leu Asp Lys Trp Ala Ser Leu Trp Asn  
 145 150 155 160  
 Trp Phe Asn Ile Thr Asn Trp Leu Trp Tyr Ile Lys Ile Phe Ile Met  
 165 170 175  
 Ile Val Gly Gly Leu Ile Gly Leu Arg Ile Val Phe Thr Val Leu Ser  
 180 185 190  
 Ile Val Asn Arg Val Arg Gln Gly Tyr Ser Pro Leu Ser Phe Gln Thr  
 195 200 205  
 Arg Leu Pro Val Pro Arg Gly Pro Asp Arg Pro Glu Gly Ile Glu Glu  
 210 215 220  
 Glu Gly Gly Glu Arg Asp Arg Ser Thr Arg Leu Val Thr Gly  
 225 230 235 240  
~~Phe~~ Leu Pro Leu Ile Trp Asp Asp Leu-Arg Ser Leu Phe Leu Phe Ser  
 245 250 255  
 Tyr His Arg Leu Arg Asp Leu Leu Leu Ile Val Ala Arg Ile Val Glu  
 260 265 270  
 Leu Leu Gly Arg Arg Gly Trp Glu Ile Leu Lys Tyr Trp Trp Asn Leu  
 275 280 285  
 Leu Gln Tyr Trp Ser Gln Glu Leu Lys Asn Ser Ala Val Ser Leu Leu  
 290 295 300  
 Asn Ala Thr Ala Ile Ala Val Ala Glu Gly Thr Asp Arg Val Ile Glu  
 305 310 315 320  
 Val Val Gln Arg Ile Trp Arg Gly Ile Leu His Ile Pro Thr Arg Ile  
 325 330 335  
 Arg Gln Gly Phe Glu Arg Ala Leu Leu  
 340 345